

Effectiveness of this product will be noticeable within one hour after the Dispenser is activated. Recommended use is for 24 hours: but if less than 24 hour is required, the Dispenser should be set to start one hour before control is desired and run one hour after control is no longer needed. Factors such as airflow rate, size of space treated, humidity and room temperature affect insect activity and control. Therefore, one unit per 170 m³ (6,000 cubic feet) of unobstructed space should be considered as an average with adjustments made depending on specific conditions at usage site. As with any method of insect control, best results are achieved when sanitation and other positive insect prevention practices are followed.]

DIRECTIONS FOR USE: Remove the protective cap from the top of the can and insert can into dispenser. Place dispenser 2-3 metres above the floor at a point where airflow will carry the insecticide throughout the area. In areas where the ceiling are low, such as attics and crawl spaces, install the dispenser approximately 0.5 metres from the ceiling. Do not install directly over food or food packing surfaces. Read dispenser instruction sheet for installation assistance.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Contents under pressure. Container may explode if heated. Do not expose to temperatures above 50° C. Do not spray near open flames or sparks. Do not puncture or incinerate. Avoid contact with skin, eyes and clothing. Avoid inhaling spray mist. Do not install directly over polished, plastic or varnished surfaces. Do not use in nurseries or rooms where infants, ill or aged persons are confined. Do not install directly above or within 3.7 meters (12.1 feet) of any food handling or dispensing area. Wash after using. Content toxic to fish and birds. Do not contaminate food, feed or water. Do not install in the living areas of the home such as the kitchens, bedrooms or living room.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: This product contains petroleum distillate. Vomiting may cause aspiration pneumonia. Treat symptomatically.

DISPOSAL: When container is empty, press valve to release all pressure. Dispose of empty container in household garbage.

or

DISPOSAL: When container is empty, press button to release all pressure then dispose in household garbage. This container may be recycled in communities where **(steel) aerosol can recycling** is available. Before offering for recycling, empty the can by using the product according to the label (**DO NOT PUNCTURE OR INCINERATE**).