OPPORTUNE PTO

BIOLOGICAL HERBICIDE

Liquid COMMERCIAL

GUARANTEE.

KEEP OUT OF REACH OF CHILDREN

POTENTIAL SENSITIZER

READ THE LABEL BEFORE USING

REGISTRATION NUMBER: 31166 PEST CONTROL PRODUCTS ACT

Net Contents: XX (Batch)(Lot) No: XXXX Date of manufacture:

Manufactured by: Pro Farm Group, Inc. 1530 Drew Avenue Davis, California, USA 95618 1-877-664-4476; info@profarmgroup.com

GENERAL INFORMATION

OPPORTUNE PTO is a selective post-emergent herbicide that can provide partial suppression of dandelions infesting turfgrass. Close scouting and early attention to infestations is highly recommended. Proper timing of application targeting newly emerged weeds is important for optimal results

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. May cause sensitization. Avoid contact with eyes, skin and clothing. Avoid inhaling/breathing mists. Wear waterproof gloves, a long-sleeved shirt, long pants, a dust-mist filtering respirator/mask (NIOSH approval number prefix TC-21C) or NIOSH approved respirators (with any N-95, P-95, R-95 or HE filter for biological products), and shoes plus socks when handling, mixing/loading or applying the product and during all clean-up and repair activities. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before re-use

FIRST AID		
	 Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. 	
lf in eyes	• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.	
	Call a poison control center or doctor for treatment advice.	
	Move person to fresh air.	
lf inhaled	• If person is not breathing, call 911 or an ambulance, then give artificial respiration,	
	preferably mouth-to-mouth if possible.	
	Call poison control center or doctor for treatment advice.	
	Call poison control center or doctor immediately for treatment advice,	
If swallowed	Have person sip a glass of water if able to swallow.	
	• Do not induce vomiting unless told to do so by the poison control center or doctor.	
	 Do not give anything by mouth to an unconscious person. 	
	Take off contaminated clothing.	
lf on skin or	• Rinse skin immediately with plenty of water for 15 – 20 minutes.	
clothing	Call a poison control center or doctor for treatment advice.	

TOXICOLOGICAL INFORMATION: Treat symptomatically.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

DIRECTIONS FOR USE

The concentrate of OPPORTUNE PTO must be mixed with water and applied as a foliar spray with hand-held equipment. Recommended sprayers include hand-held 1 litre bottles with trigger sprayers and pump sprayers of 2 - 10 litre volume. Thorough coverage of weed foliage is necessary for effective control. Spray patterns should be adjusted to deliver medium sized droplets. Avoid fine droplets that are prone to drift.

Do not apply this product through any type of irrigation equipment.

OPPORTUNE PTO does not have systemic activity. Newly emerged dandelions are more easily controlled than are mature plants. For mature dandelions, use the higher label rate and/or increase the spray volume to improve coverage.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

FOR USE ON THE FOLLOWING TURF SPECIES FOR THE PARTIAL SUPPRESSION OF DANDELION:

Fescues	<i>Festuca</i> spp.
Kentucky bluegrass	Poa pratensis

MIXING INSTRUCTIONS

Mix OPPORTUNE PTO with water at a rate of 100 to 200 ml per litre of total spray volume.

Important – Do not add OPPORTUNE PTO to the mix sprayer reservoir before introducing the desired amount of water. Add water to the sprayer reservoir or tank. Add the desired volume of OPPORTUNE PTO to the sprayer reservoir and shake well to thoroughly mix spray solution.

APPLICATION INSTRUCTIONS

Apply spray mixture to individual dandelions by providing complete coverage of visible foliage. For a longer period of dandelion suppression, additional applications may be necessary after the initial application. Repeat applications at 21 day intervals as required. Use the lower rate for newly germinated dandelions and the higher rate for more mature dandelions.

Avoid overspray of surrounding turf. OPPORTUNE PTO applied at the highest rate may cause temporary discoloration of affected turf. Symptoms are more likely during periods of high temperature and/or moisture stress. To avoid excessive turf injury, do not apply to newly seeded lawns.

STORAGE

Store product in a cool, dark place using original container. Store at up to 25°C for up to two years from date of manufacture. Keep container tightly closed when not in use. Do not contaminate water, food or feed by storage or disposal.

DISPOSAL

- 1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
 - 2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.

- 3. Make the empty container unsuitable for further use.
- 4. Dispose of the container in accordance with provincial requirements.
- 5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the PEST CONTROL PRODUCTS ACT to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product