

GROUP	5	15	HERBICIDE
-------	---	----	-----------

FRONTIER® MAX PLUS

For weed control in corn (field and sweet)

HERBICIDE

SUSPENSION CONCENTRATE

COMMERCIAL (AGRICULTURAL)

DO NOT USE IN THE PROVINCE OF BRITISH COLUMBIA

GUARANTEE: Dimethenamid-P.....204 g/L
Atrazine 396 g/L

Contains 1,2-benzisothiazolin-3-one at 0.014 % as a preservative

REGISTRATION NO. 30519

PEST CONTROL PRODUCTS ACT



WARNING

POISON

**EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER**

Warning, contains the allergen soy

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**

NET CONTENTS: 1 L - 1000 L, Bulk

BASF Canada Inc.
5025 Creekbank Road
Building A, 2nd Floor
Mississauga, Ontario L4W 0B6

1-877-371-2273

FRONTIER is a registered trade-mark of BASF Corporation, used with permission by BASF Canada Inc.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. Eye irritant. May cause skin sensitization reactions in certain individuals. DO NOT get in eyes, on skin, or on clothing. Avoid inhalation of vapour or spray mist.

Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking or going to the washroom. Take a shower immediately after work.

Wear coveralls over a long-sleeved shirt and long pants, shoes plus socks and chemical resistant gloves during mixing, loading, application, clean-up and repair. In addition, wear goggles or face shield during mixing and loading.

If clothing becomes contaminated, remove immediately and wash. Store and wash all protective clothing separately from household laundry. Wash in detergent and hot water before reuse. Wear freshly laundered clothes daily.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Mixing and loading is not permitted within 30 m from aquatic habitats.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

ENVIRONMENTAL HAZARDS

Toxic to non-target terrestrial plants and aquatic organisms. Observe the buffer zones and precautionary measures specified under DIRECTIONS FOR USE.

Atrazine is persistent and will carry over. It is recommended that any products containing atrazine not be used in areas treated with this product during the previous season.

STORAGE

1. Store FRONTIER MAX PLUS in original container only.
2. Store FRONTIER MAX PLUS herbicide in a cool, dry place.
3. Do not contaminate water, food or feed by storage.

DISPOSAL

Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

Returnable-Refillable Containers

For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

GENERAL INFORMATION

FRONTIER MAX PLUS will provide control of most annual grass and broadleaf weeds in field and sweet corn. Most effective weed control is obtained when FRONTIER MAX PLUS is applied and subsequently incorporated into soil by rainfall or mechanical tillage prior to weed seedling emergence from soil.

FRONTIER MAX PLUS can be applied early pre-plant, pre-plant incorporated, pre-emergence or early post-emergence to field corn either alone and/or in tank-mix combination. FRONTIER MAX PLUS may also be applied to sweet corn pre-plant incorporated or pre-emergence.

Apply FRONTIER MAX PLUS at a labeled rate once per year.

DIRECTIONS FOR USE

WEEDS CONTROLLED

FRONTIER MAX PLUS applied as directed will control the following weeds:

- Barnyard grass
- Common purslane
- Crabgrass (smooth, large)
- Eastern black nightshade*
- Fall panicum
- Foxtail (green, yellow, giant)
- Lady's-thumb
- Lamb's-quarters
- Old witchgrass
- Ragweed
- Redroot pigweed
- Smartweed
- Volunteer clover
- Wild buckwheat
- Wild mustard
- Wild oats

- Wormseed mustard
- Yellow nutsedge**

* pre-plant incorporated or pre-emergence only (3.40 L/ha).

** pre-plant incorporated only (3.40 L/ha. Lower rates provide suppression only.)

Apply atrazine at a maximum of 1.5 kg ai/ha per year either as a pre- or post-emergent application before corn reaches 30 cm in height. Treatments can be made once or twice per year, but the maximum amount of atrazine used in a year cannot exceed 1.5 kg ai/ha. Atrazine may be applied alone or as a registered tank mix partner. Refer to the tank mix partners for further instructions.

FACTORS AFFECTING FRONTIER MAX PLUS PERFORMANCE

If drought conditions persist after a pre-plant incorporated or pre-emergence application, annual grass and broadleaf weed control may not be adequate.

Rainfall is required within 7 to 10 days of treatment to activate and move FRONTIER MAX PLUS into the soil zone. If dry conditions persist a shallow cultivation or the use of a rotary hoe is necessary to move the chemical into moist soil and control weed escapes. If cultivation is necessary because of soil crusting or compaction, tillage should be shallow to minimize dilution of the herbicide.

SPRAYING INSTRUCTIONS

Select proper nozzle to avoid spraying fine mist. For best results, use sprayers equipped with flat fan or similar nozzles to ensure coverage. Apply in a minimum spray volume of 170 L and at a pressure of 200 - 300 kPa. For applications to dense weed infestations and thick canopies, use a higher spray volume. Better coverage of the product results in enhanced control of weeds.

Keep bypass line on or near bottom of tank to minimize foaming. Use 16-mesh suction screens, 50-mesh screens elsewhere on sprayer.

Protect sprayer operators from drift or mist. Additional information on spray drift management for ground application is provided in the section "SPRAY DRIFT MANAGEMENT FOR GROUND APPLICATIONS". When low volumes of spray are applied, complete coverage and thorough application are essential for the most effective results. Schedule applications in accordance with local conditions. Consult your local agricultural authorities for specific use information.

APPLICATION INSTRUCTIONS FOR FULL SEASON WEED CONTROL

FRONTIER MAX PLUS may be used alone for control of labeled grass and broadleaf weeds in field corn when applied as a pre-plant incorporated, pre-emergence or early post-emergence treatment. For sweet corn, FRONTIER MAX PLUS may be applied as a pre-plant incorporated or pre-emergence treatment. Use rates of this product vary by soil texture and organic matter.

For burndown and season-long residual control of most annual grass and broadleaf weeds in no-till and reduced-till corn, FRONTIER MAX PLUS may be tank mixed with glyphosate and applied as an early pre-plant surface application.

As this pesticide is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT apply by air.

Pre-plant Incorporated Application

Apply FRONTIER MAX PLUS as a broadcast treatment and incorporate using a harrow, rolling cultivator or other implement capable of giving uniform shallow incorporation into the top 5 cm of soil within 7 days of planting. Avoid deeper incorporation or reduced weed control and/or crop injury may result. Immediate incorporation after application is not necessary. Use the higher label rate of FRONTIER MAX PLUS on fine textured or high organic soils and for heavier weed problems.

Rate: 3.04 to 3.40 L/ha

Pre-emergence and Early Post-emergence Application

Apply FRONTIER MAX PLUS at the higher rates illustrated in the table below on fine textured or high organic soils and for heavier weed problems. Do not apply beyond the 2-leaf stage of weeds. Apply after planting and up to and including the 4-leaf stage of corn.

Some leaf burn may occur with early post-emergence applications but yield will not be affected.

Unless otherwise specified, DO NOT mix FRONTIER MAX PLUS with fluid fertilizers, oils, oil concentrates or surfactants when applying post-emergence as severe crop injury may occur.

DO NOT apply to sweet corn as a post-emergence application.

Soil Type	Rate (L/ha) Organic Matter		
	Less than 3 %	3 to 6 %	7 to 10 %
Coarse Textured Soils	2.67	2.67	3.04
Medium Textured Soils	2.67	3.04	3.40
Fine Textured Soils	2.67	3.04	3.40

Tank Mixtures for Full Season Weed Control – Field Corn Only

DO NOT use these tank mixes on sweet corn.

Surface Application Prior to Planting or After Planting Prior to Crop Emergence

For burndown and season-long residual control of most annual grass and broadleaf weeds in no-till and reduced-till corn, FRONTIER MAX PLUS may be tank mixed at a rate of 2.67 to 3.40 L/ha with glyphosate and applied as a surface application prior to or after planting, but before crop emergence. Use the higher label rate of FRONTIER MAX PLUS on fine-textured or high-organic soils and for heavier weed problems.

Consult the glyphosate label for weeds controlled by glyphosate, as well as for further recommendations, precautions and restrictions. FRONTIER MAX PLUS is compatible with all liquid glyphosate formulations registered for use as a pre-plant treatment prior to corn in which glyphosate is present as an isopropylamine salt, di-ammonium salt or potassium salt.

FRONTIER MAX PLUS Rate	Tank Mix Partner Rate
FRONTIER MAX PLUS at 2.67 to 3.40 L/ha ¹	2.5 L/ha of glyphosate (360 g/L equivalent)

¹ Use higher label rate on fine-textured or high-organic soils and for heavier weed problems.

APPLICATION INSTRUCTIONS FOR SHORT SEASON WEED CONTROL

Reduced rates of FRONTIER MAX PLUS may be used where a reduced length of soil residual control is required, such as in post-emergence applications or pre-emergence applications where an in-crop application of another registered herbicide is planned for added control of the same targeted weed species.

Pre-Emergence Application

FRONTIER MAX PLUS may be applied as a pre-emergence treatment at a reduced rate of 2.15 L/ha for early season control or suppression of the weeds listed below when there is a planned in-crop application of another registered herbicide such as glyphosate in glyphosate tolerant corn.

FRONTIER MAX PLUS applied early season at this rate will provide early season control of redroot pigweed and early season suppression of common ragweed, crabgrass (large and smooth), foxtail (green, yellow, and giant), lamb's-quarters, barnyard grass, and Eastern black nightshade.

Early Post-Emergence Application

For season long control of late flushing weeds listed below in glyphosate tolerant field corn, FRONTIER MAX PLUS may be applied at a reduced rate of 2.15 L/ha as an early post-emergence application in tank mix combination with glyphosate. The in-crop application of glyphosate will control any emerged weeds in glyphosate tolerant corn and will allow for a lower rate of FRONTIER MAX PLUS to control the later emerging flushes of weeds during the season.

Redroot pigweed, common ragweed, lamb's-quarters, crabgrass (large and smooth), barnyard grass, foxtail (green, yellow, and giant), wild buckwheat, lady's-thumb, smartweed, and Eastern black nightshade.

WARNING: Apply glyphosate on glyphosate tolerant field corn varieties only, i.e. varieties with the Roundup Ready gene. **CORN VARIETIES WHICH ARE NOT DESIGNATED AS GLYPHOSATE TOLERANT WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.**

Consult the glyphosate label for weeds controlled by glyphosate, as well as for further recommendations, precautions and restrictions. FRONTIER MAX PLUS is compatible with all liquid glyphosate formulations registered for use as a pre-plant treatment prior to corn in which glyphosate is present as an isopropylamine salt, di-ammonium salt or potassium salt.

FRONTIER MAX PLUS Rate	Tank Mix Partner Rate
FRONTIER MAX PLUS at 2.15 L/ha	2.5 L/ha of glyphosate (360 g/L equivalent)

LIQUID FERTILIZER

FRONTIER MAX PLUS may be applied in a nitrogen solution or complete liquid fertilizer carrier as a replacement for water for pre-plant incorporated or pre-emergence application. DO NOT use fertilizer carriers for post-emergence application. Follow the tank mix partner product label for instructions regarding application in liquid fertilizers. Compatibility of FRONTIER MAX PLUS with the liquid fertilizer should be determined by mixing a small quantity of herbicide with a proportional quantity of liquid fertilizer in a jar test prior to loading the spray tank.

DRY BULK FERTILIZER

For pre-plant incorporated application, dry bulk fertilizers may be impregnated with FRONTIER MAX PLUS. The herbicide should be sprayed directly onto the fertilizer using a common dry bulk fertilizer blend. A highly absorptive powder such as a diatomaceous earth or finely powdered clay may be applied if the herbicide/fertilizer mixture is excessively wet. A minimum of 225 g per hectare of dry bulk fertilizer should be applied.

The herbicide/fertilizer mixture should be incorporated immediately after application. DO NOT impregnate FRONTIER MAX PLUS on nitrate fertilizers, superphosphates or limestone.

MIXING INSTRUCTIONS

1. When applying FRONTIER MAX PLUS always start with a clean sprayer. Thoroughly clean the sprayer by flushing the system with water containing detergent. Refer to previously applied product labels for specific cleaning instructions.
2. Fill clean spray tank one-half full of clean water and start agitation system.
3. If using a water dispersible granule tank mix partner, add correct amount to tank and continue agitation until thoroughly dispersed.
4. Add the correct amount of FRONTIER MAX PLUS and continue to agitate.

5. If required, add the correct amount of the liquid, solution or suspension tank mix partner while continuing agitation.
6. Continue agitation while filling the spray tank with the remaining amount of water.
7. Maintain continuous and constant agitation throughout application until spraying is complete. If spray mixture is allowed to settle at any time, thorough gentle agitation is required to re-suspend the mixture before spraying is resumed. Do not allow the mixture to sit over night.
8. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent. Refer to specific tank mix partner label for clean-up instructions. Dispose of all rinsings in accordance with provincial regulations.

RECROPPING GUIDELINES

In the event of crop failure in fields treated with FRONTIER MAX PLUS, only field corn can be planted as a rescue crop. Except for sugar beets and tobacco, all other crops may be planted one year after application of FRONTIER MAX PLUS at label rates. When extended periods of dry weather occur during the year of treatment, there can be some injury to succeeding crops such as white beans, onions, peas, tomatoes and turnips. Injury is most likely to occur when the seedling crop is subjected to periods of stress, such as abnormally hot, dry weather.

RESTRICTIONS AND LIMITATIONS

1. Clean sprayer thoroughly after use to avoid damage to the next crop sprayed.
2. DO NOT apply to field corn within 100 days of harvest.
3. DO NOT apply to sweet corn within 60 days of harvest.
4. DO NOT graze or feed the treated immature corn to livestock prior to 40 days after application of FRONTIER MAX PLUS.
5. DO NOT contaminate irrigation/drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
6. DO NOT enter or allow worker entry into treated areas for 12 hours after application.

For workers incorporating atrazine into dry bulk fertilizers, the following conditions must be met:

- This activity is restricted to commercial facilities (to prohibit on-farm impregnation).
- The facilities must employ a closed mix/load system.
- A maximum of 1500 kg active ingredient can be incorporated per day.
- This activity must occur for no more than 30 days per year.

SPRAY DRIFT MANAGEMENT FOR GROUND APPLICATIONS

General Information

Use good pesticide practices and apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and parks is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings used for application.

For the protection of non-target habitats, over-spray or drift to any body of water or other environmentally sensitive habitat must be avoided. The interaction of equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

Field Sprayer Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the weed canopy or ground.

DO NOT apply by air.

Buffer Zones

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands), and estuarine/marine habitats.

Application Method	Buffer Zone for Protection of:	
	Terrestrial Habitats (m)	Aquatic Habitats (m)
Field sprayer*	10	10

*For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labeled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labeled buffer zone can be reduced by 30%.

When a tank mix is used, consult the labels of the tank mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

SURFACE RUNOFF

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil and poorly draining soil (e.g. soils that are compacted, fine textured or low in organic matter such as

clay). Avoid applying this product when heavy rain is forecast. Potential for contamination of aquatic areas as a result of runoff may be reduced by including an untreated vegetative strip (buffer zone) between the treated area and the edge of the water body.

LEACHING

The use of this chemical may result in contamination of groundwater, particularly in areas where soils are permeable (e.g. sand, loamy sand and sandy loam soils) and/or the depth to the water table is shallow.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, FRONTIER MAX PLUS is both a Group 5 and a Group 15 herbicide. Any weed population may contain or develop plants naturally resistant to FRONTIER MAX PLUS and other Group 5 and 15 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of FRONTIER MAX PLUS or other Group 5 and 15 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information or to report suspected resistance, contact BASF at 1-877-371-2273 or at www.agsolutions.ca.

GROUP	5	15	HERBICIDE
-------	---	----	-----------

FRONTIER® MAX PLUS

For weed control in corn (field and sweet)

HERBICIDE

SUSPENSION CONCENTRATE

COMMERCIAL (AGRICULTURAL)

DO NOT USE IN THE PROVINCE OF BRITISH COLUMBIA

GUARANTEE: Dimethenamid-P.....204 g/L
Atrazine 396 g/L

Contains 1,2 – benzisothiazolin-3-one at 0.014 % as a preservative

REGISTRATION NO. 30519

PEST CONTROL PRODUCTS ACT



WARNING

POISON

**EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER**

Warning, contains the allergen soy

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**

NET CONTENTS: 1 L - 1000 L, Bulk

BASF Canada Inc.
5025 Creekbank Road
Building A, 2nd Floor
Mississauga, Ontario L4W 0B6

1-877-371-2273

FRONTIER is a registered trademark of BASF Corporation, used with permission by BASF Canada Inc.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. Eye irritant. May cause skin sensitization reactions in certain individuals. DO NOT get in eyes, on skin, or on clothing. Avoid inhalation of vapour or spray mist.

Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking or going to the washroom. Take a shower immediately after work.

Wear coveralls over a long-sleeved shirt and long pants, shoes plus socks and chemical resistant gloves during mixing, loading, application, clean-up and repair. In addition, wear goggles or face shield during mixing and loading.

If clothing becomes contaminated, remove immediately and wash. Store and wash all protective clothing separately from household laundry. Wash in detergent and hot water before reuse. Wear freshly laundered clothes daily.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Mixing and loading is not permitted within 30 m from aquatic habitats.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

ENVIRONMENTAL HAZARDS

Toxic to non-target terrestrial plants and aquatic organisms. Observe the buffer zones and precautionary measures specified under DIRECTIONS FOR USE.

Atrazine is persistent and will carry over. It is recommended that any products containing atrazine not be used in areas treated with this product during the previous season.

STORAGE

1. Store FRONTIER MAX PLUS in original container only.
2. Store FRONTIER MAX PLUS herbicide in a cool, dry place.
3. Do not contaminate water, food or feed by storage.

DISPOSAL

Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

Returnable-Refillable Containers

For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

2022-0975
2022-03-04

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.