

Ed's Turf Protector

(pre-emergent weed seed germinator inhibitor with corn gluten)

For the inhibition of large crabgrass, white clover and dandelion seed germination in residential lawns where established perennial ryegrass or established perennial ryegrass or established Kentucky bluegrass are the predominant grass species.

DOMESTIC

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

GUARANTEE: com gluten meal 100%

REG. NO. 28653 PEST CONTROL PRODUCTS ACT

CAUTION EYE: IRRITANT, POTENTIAL SENSITIZER

NET CONTENTS: 20 Kg

ED'S Purely Organic Lawn and Garden Products Inc.

101 Duncan Mill Road, Suite 400

Toronto, Ontario

M3B 1Z3

1-800-777-LAWN (5296)

Ed's Turf Protector inhibits the germination of large crabgrass, white clover and dandelion in residential landscapes where the predominant grass species is established perennial ryegrass or established perennial ryegrass or established Kentucky bluegrass.

Application Rate:

SPRING APPLICATION:

Apply to established perennial ryegrass or established Kentucky bluegrass turf in early spring application 2 weeks between application and before weed seed germination at a rate of 9.7 kg/100 m² (20lbs. /1000ft²).

LATE SUMMER/EARLY FALL APPLICATION:

Apply to established perennial ryegrass or established Kentucky bluegrass turf in late summer or early fall after heat stress has passed at a rate of 9.7 kg/100 m² (20lbs. /1000ft²).

Directions:

1. Measure the area to be applied by taking the length multiplied by the width of the total area in metres. One normal step is approximately one metre. Measure the size of any buildings, swimming pool or driveway in the same manner and subtract this amount from the total area. The remaining total will represent the grass area to be treated.
2. Apply Ed's Turf Protector (pre-emergent weed seed germinator inhibitor) using a rotary spreader for large areas. For smaller areas, apply Ed's Turf Protector (pre-emergent weed seed germinator inhibitor) using a small hand spreader.

3. Apply Ed's Turf Protector (pre-emergent weed seed germinator inhibitor) in a criss-cross pattern. Use one half the rate in each direction. Apply when soil is moist and when rain is forecasted within 2 days of treatment. If rainfall does not occur within 2 days of treatment, irrigation is required. Excessive moisture at time of treatment may reduce the effectiveness of the product.

Ed's Turf Protector must not be applied to newly seeded grass until after the first mowing when the root systems are established, since the product inhibits seeds from germinating. For best results: Apply to established perennial ryegrass or established Kentucky bluegrass twice a year; once in the early spring, and again in late summer/early fall.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. May cause sensitization. Ed's Turf Protector (pre-emergent weed seed germination inhibitor) should not be applied if a member of the household has a sensitivity or allergy to corn.

Avoid contact with skin, eyes or clothing.

Avoid inhaling dusts.

Do not apply under windy conditions

For good hygiene practice, wear a long-sleeved shirt, long pants, closed footwear and gloves when handling the product. In addition, it is recommended that a dust mask be worn when transferring the product to the spreader.

NOTE: Do not apply the product on newly seeded grass as it may inhibit seeds from germinating, wait until after first mowing when root systems are established. Do not apply if allergic to corn. If over-seeding or sodding in the spring, do not apply the product in the spring. If over-seeding or re-sodding in the fall, do not apply the product in the fall.

FIRST AID:

IF SWALLOWED:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

GENERAL: Seek medical attention immediately if irritation or signs of toxicity occur and persist or is severe. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE: Keep in cool, dry conditions, away from seed, fertilizer and other pesticides.

DISPOSAL: Do not re-use the empty container. Wrap and dispose with household garbage.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.