

KONK 416 Flying Insect Killer

COMMERCIAL

Metered Pressurized Insecticide Spray

TO CONTROL FLIES, MOSQUITOES, GNATS AND SMALL FLYING MOTHS.

READ THE LABEL BEFORE USING

GUARANTEE:

Pyrethrins..... 1.76%
Piperonyl Butoxide..... 5.00%

See Back Panel for Use Areas and Additional Precautionary Statements.

REGISTRATION NO. 28478 PEST CONTROL PRODUCTS ACT

CAUTION

WARNING



EXPLOSIVE



FLAMMABLE

Acuity Holdings, Inc.
11627 - 178 STREET
EDMONTON AB T5S 1N6
905-669-9876

NET CONTENTS 80 - 200 g

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. CAUSES SKIN IRRITATION. POTENTIAL SKIN SENSITIZER.

Do not use in nurseries or rooms where infants, ill or aged persons are confined. Do not install directly above or within 3.7 metres (12.1 feet) of any food handling or dispensing area. IN MEAT PACKING AND FOOD PROCESSING PLANTS, only spray when plant is not in operation. No exposed meat products, ingredients or packaging material to be left in room. Cover food handling equipment. Any treated surface that may contact meat food products or other food products should be washed and rinsed thoroughly with potable water after spraying and before re-use for food packaging. Do not spray over milk or milking equipment. Avoid contact with skin and inhalation of spray mist. Wash after using. Contents toxic to fish and birds. Do not contaminate food, feed, or water.

PHYSICAL OR CHEMICAL HAZARDS: CONTENTS UNDER PRESSURE. Container may explode if heated. Do not expose to temperatures above 50°C. Do not puncture or incinerate. Do not allow spray to contact plastic, painted or varnished surfaces.

FIRST AID: In case of contact with eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. In case of contact with skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. If accidentally swallowed, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give *any* liquid to the person. Do not give anything by mouth to an unconscious person. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

DISPOSAL: When container is empty, press valve to release all remaining pressure. Dispose of empty container in accordance with provincial requirements. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

DIRECTIONS FOR USE

This product is engineered for use in automatic metered dispensers. The dispenser automatically activates the special metered valve designed to release a metered dose of either 27.5 mg of insecticide every 7 ½ minutes, 55mg of insecticide every 15 minutes, 110 mg of insecticide every 30 minutes or 165 mg of insecticide every 45 minutes. Within 24 hours the area is relatively free of flies, mosquitoes, gnats, and small flying moths and with continued use reduces annoyance from them. Effectiveness of this product will be noticeable within one hour after the Dispenser is activated. Recommended use is for 24 hours: but if less than 24 hour use is required, the Dispenser should be set to start one hour before control is desired and run one hour after control is no longer needed. Factors such as airflow rate, size of space treated, humidity and room temperature affect insect activity and control. Therefore, one unit per 170 m³ (6,000 cubic feet) of unobstructed space should be considered as an average with adjustments made depending on specific conditions at usage site. As with any method of insect control, best results are achieved when sanitation and other positive insect prevention practices are followed. It is recommended that the container be changed as required if the pest problem still exists.

FOR USE IN

Food service and preparation areas, food processing plants, restaurants, clubs, stores, supermarkets, bottling plants, warehouses, buildings, factories, offices, laboratories, lavatories, ships, planes, hotels, motels, hospitals, nursing homes, mud rooms, garages, basements, porches & patios, utility rooms, storage sheds, workshops, lanais, schools, day-care centers, funeral homes, kennels, farms, barns, stables, dairies, milk and egg handling areas of dairies and chicken houses.

OPERATION: Remove protective cap from top of can and insert can in dispenser. Suspend dispenser 2-3 meters from the floor and at a point where air flow will carry the insecticide throughout the room.