

2024-02-23
2023-2408

GROUP	28	INSECTICIDE
--------------	-----------	--------------------

HGW86 200SC Potato Seed Treatment Insecticide

SUSPENSION

FOR SALE FOR USE ON POTATOES

COMMERCIAL-AGRICULTURAL

READ THE LABEL AND ACCOMPANYING BOOKLET BEFORE USING

ACTIVE INGREDIENT: Cyantraniliprole 200 g/L

Contains 1,2-Benzisothiazolin-3-one at 0.0193 % as a preservative

-OR-

Contains 5-Chloro-2-methyl-2H-isothiazol-3-one at 0.00085%, 2-Methyl-2H-isothiazol-3-one at 0.00028%
and Bronopol at 0.0088% as a preservative

REGISTRATION NO 32270 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 0.5, 3.79, 100 LITRES

FMC of Canada Limited
6755 Mississauga Road, Suite 204
Mississauga, ON L5N 7Y2
1-833-362-7722

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Wear long sleeves, long pants and chemical resistant gloves while mixing /loading, applying and during clean up and repair.
- Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IN CASE OF EMERGENCY, CALL TOLL FREE, DAY OR NIGHT: 1-800-331-3148

TOXICOLOGICAL INFORMATION: Treat symptomatically.

ENVIRONMENTAL HAZARDS:

- Toxic to aquatic organisms.
- **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
- Toxic to bees. This product is systemic and bees can be exposed to product residues in flower, leaves, pollen and/or nectar resulting from soil applications. However, when this product is applied and used according to label directions, risk to bees is expected to be negligible.
- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

STORAGE:

Store product in original container only, away from other pesticides and fertilizer. Not for use or storage in or around the home. Keep container closed. Store this product away from food or feed.

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

<<DFU Booklet>>

GROUP	28	INSECTICIDE
--------------	-----------	--------------------

HGW86 200SC POTATO SEED TREATMENT Insecticide

SUSPENSION

FOR SALE FOR USE ON POTATOES

COMMERCIAL-AGRICULTURAL

READ THE LABEL AND ACCOMPANYING BOOKLET BEFORE USING

ACTIVE INGREDIENT: Cyantraniliprole 200 g/L

Contains 1,2-Benzisothiazolin-3-one at 0.0193 % as a preservative

-OR-

Contains 5-Chloro-2-methyl-2H-isothiazol-3-one at 0.00085%, 2-Methyl-2H-isothiazol-3-one at 0.00028%
and Bronopolat 0.0088% as a preservative

REGISTRATION NO 32270 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 0.5, 3.79, 100 LITRES

FMC of Canada Limited
6755 Mississauga Road, Suite 204
Mississauga, ON L5N 7Y2
1-833-362-7722

GENERAL INFORMATION

HGW86 200SC POTATO SEED TREATMENT insecticide is a suspension concentrate that can be applied as an in-furrow spray, transplant water treatment, banded surface application, or potato seed-piece treatment to control many important insect pests. HGW86 200SC POTATO SEED TREATMENT insecticide is specially formulated for maximum performance by soil and seed-piece treatment application methods. HGW86 200SC POTATO SEED TREATMENT insecticide is mixed with water for application.

HGW86 200SC POTATO SEED TREATMENT insecticide provides early season control of listed insect pests. The length of control will vary depending on the application rate and local conditions. Fields should be scouted regularly. If insect pest populations exceed local threshold levels, apply a registered foliar insecticide.

HGW86 200SC POTATO SEED TREATMENT insecticide is a member of the anthranilic diamide class of insecticides which act on insect ryanodine receptors. After exposure to HGW86 200SC POTATO SEED TREATMENT insecticide, affected insects will rapidly stop feeding, become paralyzed, and typically die within 1 - 3 days. When pest populations are expected to be high, use the highest listed application rate for that pest.

DIRECTIONS FOR USE

POTATOES

SITE	PEST	RATE HGW86 200SC POTATO SEED TREATMENT insecticide	SPECIFIC DIRECTIONS
POTATOES	<p>Colorado potato beetle</p> <p>Potato flea beetle (early season control of spring adults)</p>	<p>Seed-Piece Treatment 45 mL/100 kg of seed pieces</p> <p>Or</p> <p>1000 ml/ha (based on 2200 kg/ha planting rate)</p>	<p><u>Potato seed-piece treatment:</u> Apply only in areas with adequate ventilation or areas equipped to remove spray mist or dust. Good coverage of the seed-piece is required for optimal control. Following application of HGW86 200SC POTATO SEED TREATMENT insecticide, application of a fungicide potato seed treatment or inert dust is recommended.</p> <p>Colorado potato beetle resistance management: DO NOT apply any Group 28 insecticide for Colorado potato beetle control if VERIMARK™ Insecticide was used at planting as an in-furrow or seed-piece treatment.</p>

RESTRICTIONS AND PRECAUTIONS: Potatoes

- Do not make more than 1 seed-piece application per season.
- Do not exceed a total of 1.5 litres HGW86 200SC POTATO SEED TREATMENT insecticide per ha per season. Use only a closed treatment system.
- DO NOT use open cab planting equipment when planting potato seed pieces treated with HGW86 200SC POTATO SEED TREATMENT insecticide. Use only closed cab planting equipment. Workers working outside the closed cab must wear chemical-resistant coveralls over a long-sleeved shirt, long pants, a dust mask, goggles, chemical-resistant gloves, socks and chemical-resistant boots.
- Do not make a subsequent foliar application of any Group 28 insecticide for a minimum of 60 days following an in-furrow application or planting of seed pieces treated with HGW86 200SC Potato Seed Treatment insecticide.

TANK MIXES:

This product may be tank mixed with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact FMC of Canada Limited at 1-833-362-7722 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

Application equipment must be clean and free of previous pesticide deposits before applying HGW86 200SC POTATO SEED TREATMENT insecticide. Always add the HGW86 200SC POTATO SEED TREATMENT insecticide to water, never put HGW86 200SC POTATO SEED TREATMENT insecticide into a dry tank or other mixing equipment without first adding water. Mix thoroughly to fully disperse the insecticide; once dispersed continued agitation is required. Use mechanical or hydraulic means; do not use air agitation. Spray mix should not be stored overnight in spray tank.

Acidification of Spray Tank:

All soil applications of HGW86 200SC POTATO SEED TREATMENT insecticide should be adjusted to approximately pH 4 - 6. For at-plant soil applications, adjust the pH of the application spray tank using a commercially available acidifier. Adjust the pH of the application solution after all products being applied have been added to the tank. Once prepared, an application solution may be held for up to 8 hours before starting the application. Do not store the application solution overnight.

SPRAYER CLEANUP:

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.

Drain spray equipment. Thoroughly rinse sprayer and flush hoses and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Wear long sleeves, long pants and chemical resistant gloves while mixing /loading, applying and during clean up and repair.
- Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so

by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IN CASE OF EMERGENCY, CALL TOLL FREE, DAY OR NIGHT: 1-800-331-3148

TOXICOLOGICAL INFORMATION: Treat symptomatically.

ENVIRONMENTAL HAZARDS:

- Toxic to aquatic organisms.
- **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
- Toxic to bees. This product is systemic and bees can be exposed to product residues in flower, leaves, pollen and/or nectar resulting from soil applications. However, when this product is applied and used according to label directions, risk to bees is expected to be negligible.
- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

ROTATIONAL CROP RESTRICTIONS:

Recommended Plant-Back Intervals (PBI) for Rotational Crops.

PBI	CROPS
0 days	Crop Group 1C (Tuberous and corm vegetables); Crop Group 3-07 (Bulb vegetables); Crop Group 4 (Leafy vegetables; except <i>Brassica</i> vegetables); Crop Group 5 (<i>Brassica</i> (Cole) leafy vegetables); Crop Group 8-09 (Fruiting vegetables); Crop Group 9 (Cucurbit vegetables); Crop Group 13-07B (Bushberry subgroup); Crop Group 20 (Oilseed); Crop Group 1B: Root vegetables
30 days	Crop Group 6 (Legume vegetables (succulent or dried); Crop Group 7 (Foliage of legume vegetables); Crop Group 13-07G (Lowgrowing berry subgroup); Crop Group 15 (Cereal grains); Crop Group 16 (Forage, fodder, and straw of cereal grains); Crop Group 17 (Grass forage, fodder, and hay); Crop Group 18 (Nongrass animal feeds: Forage, fodder, straw and hay)
365 days	Other crops

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, please note that HGW86 200SC POTATO SEED TREATMENT™ insecticide contains a Group 28 insecticide. Any insect population may contain individuals naturally resistant to HGW86 200SC POTATO SEED TREATMENT insecticide and other Group 28 insecticides. The resistant individuals may dominate the insect population if this group of insecticides is used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of HGW86 200SC POTATO SEED TREATMENT insecticide or other Group 28 insecticides with different groups that control the same pests.
- Avoid using the same mode of action (same IRAC group number) on consecutive generations of insect pests.
- For short cycle crops (< 50 days), alternate to different modes of action during subsequent planting at the same farm location.
- Use tank mixtures with insecticides from a different group that is effective on the target pest when such use is permitted.
- Insecticide use should be based on an IPM program that includes scouting and, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact FMC at 1-833-362-7722.

STORAGE:

Store product in original container only, away from other pesticides and fertilizer. Not for use or storage in or around the home. Keep container closed. Store this product away from food or feed.

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

®/TM Registered trademarks/trademarks of FMC Corporation or an affiliate.

*All other products mentioned are trademarks of their respective companies.