

<b>GROUP</b>	<b>U15</b>	<b>FUNGICIDE</b>
--------------	------------	------------------

## **ORONDIS™ ULTRA B Fungicide**

### **OIL DISPERSION**

### **COMMERCIAL**

FOR USE ON BULB VEGETABLES, BRASSICA (COLE) LEAFY VEGETABLES, CUCURBIT VEGETABLES, GREENHOUSE CUCUMBER, TOMATO (FIELD, GREENHOUSE), PEPPER (FIELD, GREENHOUSE), EGGPLANT, LEAFY VEGETABLES, CROP SUBGROUP 4-13B: LEAFY BRASSICA GREENS, BASIL, SUCCULENT PEAS, AND POTATOES

### **GUARANTEE:**

Oxathiapiprolin ..... 100 g/L

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**

**WARNING – SKIN IRRITANT  
POTENTIAL SKIN SENSITIZER**

REGISTRATION NO.: **32146**  
PEST CONTROL PRODUCTS ACT

**NET CONTENTS:** 0.5 L – 1,500 LITRES

**Syngenta Canada Inc.**  
140 Research Lane, Research Park  
Guelph, Ontario  
N1G 4Z3  
Telephone: 1-877-964-3682

**Label**

## **NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

## **FIRST AID**

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**IF SWALLOWED:** Call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

## **TOXICOLOGICAL INFORMATION**

Treat symptomatically.

## **PRECAUTIONS**

- **KEEP OUT OF REACH OF CHILDREN.**
- Causes skin irritation. DO NOT get on skin.
- Potential skin sensitizer.
- During mixing/loading/application/clean-up and repair, workers must wear coveralls over long-sleeved shirt, long pants, socks & chemical-resistant footwear and chemical-resistant gloves.
- Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.
- Do not apply by air.

- Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.
- If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

## **ENVIRONMENTAL PRECAUTIONS**

- This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.
- As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
- **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
- May be toxic to certain beneficial insects when applied at more than 2 L/ha (200 g a.i./ha) per season. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application site such as hedgerows and woodland.
- Oxathiapiprolin is persistent and may carryover. It is recommended that any products containing oxathiapiprolin not be used in areas treated with this product during the previous season.

## **STORAGE**

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. To prevent contamination, store this product away from food or feed.

## **DECONTAMINATION AND DISPOSAL**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

### **CONTAINER DISPOSAL:**

#### **For returnable containers:**

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**For refillable containers:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

**For recyclable containers:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,  
CALL 1-800-327-8633 (FASTMED)***

ORONDIS™ is a trademark of a Syngenta Group Company.

GROUP	<b>U15</b>	FUNGICIDE
-------	------------	-----------

## ORONDIS™ ULTRA B Fungicide

### OIL DISPERSION

### COMMERCIAL

FOR USE ON BULB VEGETABLES, BRASSICA (COLE) LEAFY VEGETABLES, CUCURBIT VEGETABLES, GREENHOUSE CUCUMBER, TOMATO (FIELD, GREENHOUSE), PEPPER (FIELD, GREENHOUSE), EGGPLANT, LEAFY VEGETABLES, CROP SUBGROUP 4-13B: LEAFY BRASSICA GREENS, BASIL, SUCCULENT PEAS, AND POTATOES

### GUARANTEE:

Oxathiapiprolin ..... 100 g/L

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**

**WARNING – SKIN IRRITANT  
POTENTIAL SKIN SENSITIZER**

REGISTRATION NO.: **32146**  
PEST CONTROL PRODUCTS ACT

**Syngenta Canada Inc.**  
140 Research Lane, Research Park  
Guelph, Ontario  
N1G 4Z3  
Telephone: 1-877-964-3682

**Pamphlet**

## GENERAL INFORMATION

ORONDIS™ ULTRA B Fungicide is active against selective oomycete diseases listed on this label and has preventive and antispore activity. ORONDIS ULTRA B Fungicide is locally systemic, translaminar, and moves systemically in the xylem. ORONDIS ULTRA B Fungicide must be applied in a regularly scheduled protective spray program in rotation with other fungicides. See directions below for specific crop/disease recommendations.

Different application methods (foliar and soil) must not be combined when protecting a crop during the growing season.

ORONDIS ULTRA B Fungicide rapidly penetrates into plant tissues and is rainfast within 30 minutes after spray residues have dried.

ORONDIS ULTRA B Fungicide can be applied with specified ground equipment using sufficient water to obtain thorough coverage of plants. The minimum ground application volume is 110 L/ha.

## DIRECTIONS FOR USE

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) fine classification. Boom height must be 60 cm or less above the crop or ground.

### Cultivar/Varietal Crop Safety

Not all crops within a crop group, and not all varieties, cultivars or hybrids of crops, have been individually tested for crop safety. It is not possible to evaluate for crop safety all applications of ORONDIS ULTRA B on all crops within a crop group, on all varieties, cultivars, or hybrids of those crops, or under all environmental conditions and growing circumstances. To test for crop safety, apply the product in accordance with the label instructions to a small area of the target crop to ensure that a phytotoxic response will not occur, especially where the application is a new use of the product by the applicator.

### Buffer zones:

Use of the following spray methods or equipment **DO NOT** require a buffer zone: hand-held or backpack sprayer, spot treatment, inter-row hooded sprayer, soil drench and soil incorporation.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of application	Crop	Buffer Zones (metres) Required for the Protection of Freshwater Habitat of Depths:	
		Less than 1 m	Greater than 1 m
Field sprayer	All field crops	1	0

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most

restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

#### **Tank Mixtures:**

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Syngenta Canada Inc. at 1-877-SYNGENTA (1-877-964-3682) for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by Syngenta Canada Inc.

When applied as a tank-mix combination, read and observe all label directions, including rates, personal protective equipment, restrictions and precautions for each product used in the tank-mix. Always use in accordance with the most restrictive label restrictions and precautions.

**ORONDIS ULTRA B Fungicide will control the indicated diseases unless stated otherwise.**

**Do not apply in greenhouses unless specified below.**

#### **HEAD AND STEM BRASSICA VEGETABLES – Crop subgroup 5A**

<b>SITE</b>	<b>PEST</b>	<b>RATE ORONDIS ULTRA B Fungicide/ha</b>	<b>SPECIFIC DIRECTIONS</b>
<b>HEAD AND STEM BRASSICA VEGETABLES</b> Broccoli Broccoli, Chinese (gai lon) Brussels sprouts Cabbage Cabbage, Chinese (napa) Cabbage, Chinese, mustard (gai choy) Cauliflower Cavolo broccolo Kohlrabi	Downy Mildew ( <i>Peronospora parasitica</i> )	0.175 to 0.35 L/ha	Begin applications prior to disease development and continue on a 5 to 10 day interval. Use higher rate and shorter interval when disease pressure is high.

#### **RESTRICTIONS AND PRECAUTIONS: Head and Stem Brassica Vegetables**

- Make no more than 2 sequential applications of ORONDIS ULTRA B Fungicide before switching to a fungicide with a different mode of action. Do not use ORONDIS ULTRA B Fungicide for more than 33% of the total foliar fungicide applications or make more than 4 applications of product per crop per year, whichever is more restrictive. Where multiple crop cycles are produced in the same year do not exceed 6 applications of ORONDIS ULTRA B Fungicide per hectare per year.
- Minimum spray interval is 5 days.
- Maximum seasonal use rate is 1.40 L/ha.

- Minimum time (PHI) between application and harvest is 0 days.
- For any of the pests listed above, use the high rate under heavy pest pressure.

### BULB VEGETABLES – Crop Group 3

SITE	PEST	RATE ORONDIS ULTRA B Fungicide/ha	SPECIFIC DIRECTIONS
<b>BULB VEGETABLES</b> Chive, fresh leaves; chive, Chinese, fresh leaves; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; kurrat; lady's leek; leek; leek, wild; onion, Beltsville bunching; onion, bulb; onion, Chinese, bulb; onion, fresh; onion, green; onion, macrostem; onion, pearl; onion, potato, bulb; onion, tree, tops; onion, Welsh, tops; shallot, bulb; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these.	Downy Mildew ( <i>Peronospora destructor</i> )	0.0875 to 0.35 L/ha	Begin applications prior to disease development and continue on a 5 to 10 day interval. Use higher rate and shorter interval when disease pressure is high.

### RESTRICTIONS AND PRECAUTIONS: Bulb Vegetables

- Make no more than 2 sequential applications of ORONDIS ULTRA B Fungicide before switching to a fungicide with a different mode of action. Do not use ORONDIS ULTRA B Fungicide for more than 33% of the total foliar fungicide applications or make more than 4 applications of product per crop per year, whichever is more restrictive. Where multiple crop cycles are produced in the same year do not exceed 6 applications of ORONDIS ULTRA B Fungicide per hectare per year.
- Maximum seasonal use rate is 1.40 L/ha.
- Minimum time (PHI) between application and harvest is 0 days.
- For any of the pests listed above, use the high rate under heavy pest pressure.



**CUCURBIT VEGETABLES – Crop Group 9**

SITE	PEST	RATE ORONDIS ULTRA B Fungicide/ha	SPECIFIC DIRECTIONS
<b>Cucurbit vegetables</b> Chinese waxgourd (Chinese preserving melon); citron melon; cucumber (field + greenhouse foliar only); gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); <u>Momordica</u> spp (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon (includes cantaloupe); pumpkin; squash, summer; squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon	Downy Mildew ( <i>Pseudoperonospora cubensis</i> )	0.0875 to 0.35 L/ha	Begin applications prior to disease development and continue on a 5 to 14 day interval. Use higher rate and shorter interval when disease pressure is high or for susceptible varieties.
	Phytophthora Blight – foliar phase ( <i>Phytophthora capsici</i> )	0.175 to 0.35 L/ha	Begin applications prior to disease development and continue on a 5 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.
	Phytophthora Blight – soil phase ( <i>Phytophthora capsici</i> )	0.7 to 2.8 L/ha	Apply at plant, in furrow, drip or in transplant water. See specific soil application directions in this label.

**RESTRICTIONS AND PRECAUTIONS: Cucurbit Vegetables**

- Minimum time (PHI) between application and harvest for all uses is 0 days.
- Do not follow soil applications of ORONDIS ULTRA B Fungicide or OXTP 200SC fungicide with foliar applications of ORONDIS ULTRA B Fungicide or other oxathiapiprolin-containing fungicides. Use either soil applications or foliar applications but not both for the control of relevant disease. Do not make more than 4 applications of product per crop per year by either method. Where multiple crop cycles are produced in the same year do not exceed 6 foliar applications of ORONDIS ULTRA B Fungicide per hectare per year.
- Annual maximums for specific use patterns are listed below.

**Foliar Applications**

- Make no more than 2 sequential applications of ORONDIS ULTRA B Fungicide before switching to a fungicide with a different mode of action. Do not use ORONDIS ULTRA B Fungicide for more than 33% of the total foliar fungicide applications.
- Maximum seasonal use rate for foliar use is 1.40 L/ha.
- For any of the pests listed above, use the high rate under heavy pest pressure.

**Soil Applications**

- Make no more than 2 sequential applications of ORONDIS ULTRA B Fungicide before switching to a fungicide with a different mode of action. Do not use ORONDIS ULTRA B Fungicide for more than 33% of the total soil fungicide applications.
- The minimum soil application interval is 7 days.

- Maximum seasonal soil use rate is 5.6 L/ha.
- Do not make soil applications in greenhouses.

#### LEAFY VEGETABLES – Crop Group 4

SITE	PEST	RATE ORONDIS ULTRA B Fungicide/ha	SPECIFIC DIRECTIONS
<b>LEAFY VEGETABLES</b> Amaranth leafy, Arugula, Cardoon, Celery, Celery (Chinese), Celtuce, Chevril, Chrysanthemum (edible-leaved), Chrysanthemum (garland), Corn Salad, Cress (garland), Cress (upland), Dandelion leaves, Dock, Endive, Florence Fennel, Lettuce (head & leaf), Orach, Parsley leaves, Purslane (garden), Purslane (winter), Radicchio, Rhubarb, Spinach, Spinach (vine), Spinach (New Zealand), Swiss Chard	Downy Mildew ( <i>Peronospora</i> spp, <i>Bremia</i> <i>Latacae</i> )	0.175 to 0.35 L/ha	Begin applications prior to disease development and continue on a 5 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.

#### RESTRICTIONS AND PRECAUTIONS: Leafy Vegetables

- Make no more than 2 sequential applications of ORONDIS ULTRA B Fungicide before switching to a fungicide with a different mode of action. Do not use ORONDIS ULTRA B Fungicide for more than 33% of the total foliar fungicide applications or make more than 4 applications of product per crop per year, whichever is more restrictive. Where multiple crop cycles are produced in the same year do not exceed 6 applications of ORONDIS ULTRA B Fungicide per hectare per year.
- Maximum seasonal use rate is 1.40 L/ha.
- Minimum time (PHI) between application and harvest is 0 days.
- For any of the pests listed above, use the high rate under heavy pest pressure.

**TOMATO, PEPPER, EGGPLANT**

SITE	PEST	RATE ORONDIS ULTRA B Fungicide/ha	SPECIFIC DIRECTIONS
Tomato (field, greenhouse); Pepper, bell, non-bell (field, greenhouse); Eggplant (field)	Late Blight ( <i>Phytophthora infestans</i> )	0.175 to 0.35 L/ha	Begin applications prior to disease development and continue on a 5 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.
	Phytophthora Blight – foliar phase ( <i>Phytophthora capsici</i> )	0.175 to 0.35 L/ha	Begin applications prior to disease development and continue on a 5 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.
	Phytophthora Blight – soil phase ( <i>Phytophthora capsici</i> )	0.7 to 2.8 L/ha	Apply at plant, in furrow, drip or in transplant water. See specific soil application directions in this label.

**RESTRICTIONS AND PRECAUTIONS: Tomato, Pepper, Eggplant**

- Minimum time (PHI) between application and harvest is 0 days.
- Do not follow soil applications of ORONDIS ULTRA B Fungicide or OXTP 200SC fungicide with foliar applications of ORONDIS ULTRA B Fungicide or other oxathiapiprolin-containing fungicides. Use either soil applications or foliar applications but not both for the control of relevant disease. Do not make more than 4 applications of product per crop per year by either method. Where multiple crop cycles are produced in the same year do not exceed 6 foliar applications of ORONDIS ULTRA B Fungicide per hectare per year.
- Annual maximums for specific use patterns are listed below.

**Foliar Applications**

- Make no more than 2 sequential applications of ORONDIS ULTRA B Fungicide before switching to a fungicide with a different mode of action. Do not use ORONDIS ULTRA B Fungicide for more than 33% of the total foliar fungicide applications.
- Maximum seasonal foliar use rate is 1.40 L/ha.
- For any of the pests listed above, use the high rate under heavy pest pressure.

**Soil Applications**

- Make no more than 2 sequential applications of ORONDIS ULTRA B Fungicide before switching to a fungicide with a different mode of action. Do not use ORONDIS ULTRA B Fungicide for more than 33% of the total soil fungicide applications.
- Maximum seasonal soil use rate is 5.6 L/ha.
- Do not make soil applications in greenhouses. Do not apply to hydroponic solution.

**POTATO**

SITE	PEST	RATE ORONDIS ULTRA B Fungicide/ha	SPECIFIC DIRECTIONS
Potato	Late Blight ( <i>Phytophthora infestans</i> )	0.12 – 0.35 L/ha	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.

**RESTRICTIONS AND PRECAUTIONS: Potato**

- Make no more than 2 sequential applications of ORONDIS ULTRA B Fungicide before switching to a fungicide with a different mode of action. Do not use ORONDIS ULTRA B Fungicide for more than 33% of the total foliar fungicide applications or make more than 4 applications of product per crop per year, whichever is more restrictive. Where multiple crop cycles are produced in the same year do not exceed 6 applications of ORONDIS ULTRA B Fungicide per hectare per year.
- Maximum seasonal use rate is 1.40 L/ha.
- Minimum time (PHI) between application and harvest is 5 days.
- For any of the pests listed above, use the high rate under heavy pest pressure.

**SUCCULENT PEAS**

SITE	PEST	RATE ORONDIS ULTRA B Fungicide /ha	SPECIFIC DIRECTIONS
Peas, succulent <i>Pisum</i> spp., includes dwarf pea, edible-pod pea, snow pea, sugar snap pea, English pea, garden pea, green pea.	Pea Downy Mildew ( <i>Peronospora viciae</i> )	0.175 to 0.35 L/ha	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.

**RESTRICTIONS AND PRECAUTIONS: Pea subgroup**

- Make no more than 2 sequential applications of ORONDIS ULTRA B Fungicide before switching to a fungicide with a different mode of action. Do not use ORONDIS ULTRA B Fungicide for more than 33% of the total foliar fungicide applications or make more than 4 applications of product per crop per year, whichever is more restrictive. Where multiple crop cycles are produced in the same year do not exceed 6 applications of ORONDIS ULTRA B Fungicide per hectare per year.
- Maximum seasonal use rate is 1.40 L/ha.
- Minimum time (PHI) between application and harvest is 0 days.
- For any of the pests listed above, use the high rate under heavy pest pressure.

**CROP ROTATION**

For all legume crops except succulent peas, a plant-back interval of 180 days is required. All other crops may be planted immediately following the last application.

**NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE APPLICATIONS:**

The DIRECTIONS FOR USE for this product for the uses described below were developed by persons other than the Syngenta Canada Inc. and accepted for registration by Health Canada. The Syngenta Canada Inc. itself makes no representation or warranty with respect to product performance (efficacy) and crop tolerance (phytotoxicity) of this product when used on the crops listed below. Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold the Syngenta Canada Inc. harmless from any claims based on efficacy or phytotoxicity in connection with the uses described below.

**Crop Subgroup 4-13B: Leafy Brassica Greens**

SITE	PEST	RATE ORONDIS Fungicide/ha	SPECIFIC DIRECTIONS
Arugula, Broccoli raab, Broccoli, Chinese, Cabbage, Abyssinian, Cabbage, seakale, Chinese cabbage, bok choy, Collards, Cress, garden, Cress, upland, Hanover salad, Kale, Maca, Mizuna, Mustard greens ( <i>Brassica juncea</i> subsp. including <i>Brassica juncea</i> subsp. <i>integrifolia</i> and <i>Brassica juncea</i> subsp. <i>tsatsai</i> ), Radish leaves, Rape greens, Rocket, wild, Shepherd's purse, Turnip greens, Watercress, Cultivars, varieties and/or hybrids of these	Downy Mildew ( <i>Peronospora parasitica</i> )	0.175 to 0.35 L/ha	Begin foliar applications prior to disease development, and continue on a 5 to 10 day interval. Use higher rate and shorter interval when disease pressure is high. Apply in a minimum of 100L of water, increasing spray volume as needed to ensure complete canopy coverage.

**RESTRICTIONS AND PRECAUTIONS:**

- Make no more than 2 sequential applications of ORONDIS ULTRA B fungicide before switching to a fungicide with a different mode of action. Do not use ORONDIS ULTRA B fungicide for more than 33% of the total foliar fungicide applications.
- Do not make more than 2 applications (high rate) or more than 4 applications (low rate) per year
- Maximum seasonal use rate is 1.4 L/ha.
- Minimum time (PHI) between application and harvest is 0 days.
- For any of the pests listed above, use the high rate under heavy pest pressure.

**Basil, field and greenhouse (*Ocimum basilicum*)**

SITE	PEST	RATE ORONDIS Fungicide/ha	SPECIFIC DIRECTIONS
Basil, field and greenhouse ( <i>Ocimum basilicum</i> )	Downy Mildew ( <i>Peronospora belbahrii</i> )	0.175 to 0.35 L/ha	Begin foliar applications prior to disease development, and continue on a 5 to 10 day interval. Use higher rate and shorter interval when disease pressure is high. Apply in a minimum of 100L of water, increasing spray volume as needed to ensure complete canopy coverage.

**RESTRICTIONS AND PRECAUTIONS:**

- Make no more than 2 sequential applications of ORONDIS ULTRA B fungicide before switching to a fungicide with a different mode of action. Do not use ORONDIS ULTRA B fungicide for more than 33% of the total foliar fungicide applications.
- Do not make more than 2 applications (high rate) or more than 4 applications (low rate) per year
- Maximum seasonal use rate is 1.4 L/ha.
- Minimum time (PHI) between application and harvest is 0 days.
- For any of the pests listed above, use the high rate under heavy pest pressure.

**APPLICATION INFORMATION****FOLIAR APPLICATION****Mixing Instructions For Product Used Alone**

1. Fill clean spray tank 1/4 - 1/2 full of water.
2. While agitating, add the required amount of ORONDIS ULTRA B Fungicide, continuing agitation until the product is completely dispersed.
3. Continue filling the tank, with agitation.

Mix thoroughly to fully disperse the fungicide; once dispersed continued agitation is required. Use mechanical or hydraulic means; do not use air agitation.

**SPRAYER CLEANUP:**

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.

Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

**SOIL APPLICATIONS**

For suppression or control of soil borne diseases, as recommended in this label, ORONDIS ULTRA B Fungicide must be applied in a manner that ensures the product solution adequately

saturates the target crop root/crown zone.

For soil application, apply ORONDIS ULTRA B Fungicide using drip application, transplant water application (water wheel or continuous stream transplanters), surface band or directed application, or in-furrow application. If the application method does not move the product to the target root/crown disease zone, the application must be followed with irrigation or cultivation to correctly place the product for disease control.

### **Transplant Water application**

Transplants should be adequately watered before transplanting. Ensure transplant water volume is sufficient to thoroughly wet the rootzone.

Use 125-250 mL of transplant water per transplant in sandy soil, and 175-250 mL of transplant water per transplant in silty soil.

- For Water Wheel transplanters, use the plant population to determine the rate per plant.

### **Surface Band or Directed application**

Apply in a 10-30 cm (4-12 inch) band. Follow application with cultivation or irrigation 1.25 - 2.5 cm (0.5 - 1 inch) to move ORONDIS ULTRA B Fungicide to the target disease zone.

### **Transplant Tray Application**

Apply as a foliar spray to the transplants in the transplant tray 24-48 hrs prior to transplanting. Immediately water the spray off the foliage into the transplant tray soil, not watering past soil saturation. Wait 24 hours to transplant.

- For this application, the acre of transplants receive the full recommended acre rate for transplant tray application, applied as a foliar spray, and then washed into the transplant cubes. Transplant cubes should be on the dry side at the beginning of this treatment.

### **SPRAY TANK CLEANOUT**

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.

Drain application equipment. Thoroughly rinse and flush all application equipment with clean water. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

### **PRECAUTIONS**

- **KEEP OUT OF REACH OF CHILDREN.**
- Causes skin irritation. DO NOT get on skin.
- Potential skin sensitizer
- During mixing/loading/application/clean-up and repair, workers must wear coveralls over long-sleeved shirt, long pants, socks & chemical-resistant footwear and chemical-resistant gloves.

- Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.
- Do not apply by air.
- Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.
- If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

## **ENVIRONMENTAL PRECAUTIONS**

- This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.
- As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
- **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
- May be toxic to certain beneficial insects when applied at more than 2 L/ha (200 g a.i./ha) per season. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application site such as hedgerows and woodland.
- Oxathiapiprolin is persistent and may carryover. It is recommended that any products containing oxathiapiprolin not be used in areas treated with this product during the previous season.

## **INTEGRATED PEST MANAGEMENT**

Syngenta Canada Inc. recommends the use of Integrated Pest Management (IPM) programs to control pests.

ORONDIS ULTRA B Fungicide may be used as part of an IPM program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring,



and treating when disease forecasting models reach locally determined action levels. Consult your provincial extension service, professional consultants, or other qualified authorities to determine the appropriate management, cultural practice and treatment threshold levels for the specific crop, geography and diseases.

## **RESISTANCE MANAGEMENT**

ORONDIS ULTRA B Fungicide contains the active ingredient oxathiapiprolin which is listed as Group U15 by the Fungicide Resistance Action Committee (FRAC). Oxathiapiprolin inhibits the oxysterol-binding protein (OSBP). Any fungal population may contain individuals naturally resistant to ORONDIS ULTRA B Fungicide. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

### **To delay fungicide resistance:**

Where possible, rotate the use of ORONDIS ULTRA B Fungicide or other fungicides containing Oxathiapiprolin with different mode of action groups that control the same pathogens.

Use tank mixtures with fungicide from a different group that is effective on the target pathogen when such use is permitted. Do not tank mix ORONDIS ULTRA B Fungicide with any fungicide for which resistance to the target disease has developed.

Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for resistance development. Notify Syngenta Canada Inc. if reduced sensitivity of the pathogen to ORONDIS ULTRA B Fungicide is suspected.

If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different site of action, if available.

Make no more than 2 sequential applications of ORONDIS ULTRA B Fungicide before rotating to a fungicide with a different mode of action.

Do not follow soil applications of ORONDIS ULTRA B Fungicide or OXTP 200SC Fungicide with foliar applications of ORONDIS ULTRA B Fungicide or other oxathiapiprolin-containing fungicides. Different application methods (foliar and soil) must not be combined when protecting a crop during a growing season.

Do not use ORONDIS ULTRA B Fungicide for more than 33% of the total fungicide applications

per season per crop.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information or to report suspected resistance, contact company representatives at 1-877-SYNGENTA (1-877-964-3682) or at [www.syngenta.ca](http://www.syngenta.ca).

## **STORAGE**

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. To prevent contamination, store this product away from food or feed.

## **DECONTAMINATION AND DISPOSAL**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

### **CONTAINER DISPOSAL:**

#### **For returnable containers:**

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

#### **For refillable containers:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

#### **For recyclable containers:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

## **FIRST AID**

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**IF SWALLOWED:** Call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,  
CALL 1-800-327-8633 (FASTMED)***

## **TOXICOLOGICAL INFORMATION**

Treat symptomatically.

## **NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

ORONDIS™ is a trademark of a Syngenta Group Company.