

THOR

ACTICIDE® RS MICROBIOCIDE LIQUID

An Industrial Microbiocide for use in Polymer Latices, paints, coatings, inks, adhesives, building materials (caulks), printing fluids, industrial and household consumer products, metalworking fluids, recirculating electrodeposition systems, process water systems, and pesticide products

Solution

COMMERCIAL

ACTIVE INGREDIENTS:

5-Chloro-2-methyl-4-isothiazolin-3-one 1.11%
2-Methyl-4-isothiazolin-3-one 0.37%

REGISTRATION NO.30128 PEST CONTROL PRODUCTS ACT

DANGER

CORROSIVE

KEEP OUT OF REACH OF

CHILDREN READ THE LABEL

BEFORE USING POTENTIAL

SKIN SENSITIZER

CORROSIVE TO EYES AND SKIN

NET CONTENTS: kg

BATCH NO.:

MANUFACTURING

DATE:

ACTICIDE® RS MICROBIOCIDE LIQUID weighs 1.01 kg per litre ACTICIDE® is a registered trademark of Thor Specialties, Inc.

Thor GmbH
Landwehrstrasse 1
D-67346, Speyer, Germany

Thor Specialties, Inc.
50 Waterview Drive
Shelton, CT, 06484 U.S.A.
Tel: (203) 516-6980
Fax: (203) 954-0005

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. POTENTIAL SKIN SENSITIZER. MAY CAUSE ALLERGIC SKIN REACTION. CORROSIVE TO EYES AND SKIN. Do not swallow. Do not get in eyes, on skin or on clothing. Avoid breathing vapours or spray mists. Wash skin thoroughly with soap and water after handling. Remove contaminated clothing and wash separately from other laundry before reuse.

Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and a full-face respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides, or a NIOSH-approved canister approved for pesticides, during mixing, loading, application, clean-up and repair.

ENVIRONMENTAL PRECAUTIONS

This product is toxic to fish and other aquatic organisms. It is not to be used in circumstances that would cause or allow it to enter lakes, streams, ponds, estuaries, oceans or other waters in contravention of federal or provincial regulatory requirements. The requirements of applicable laws should be determined before using the product. Do not contaminate water by cleaning of equipment or disposal of waste. Use only as specified on this label.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take the container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsions may be required.

DIRECTIONS FOR USE

PRESERVATIVE FOR LATICES:

ACTICIDE® RS MICROBIOCIDE LIQUID is recommended for the control of bacteria in the manufacture and storage of synthetic and natural polymer latices – acrylics,

styrene/butadiene, carboxylated styrene/butadiene, ethylene/vinyl acetate, starches, derived polymers.

Add 430 grams to 3.3 kg of ACTICIDE® RS MICROBIOCIDE LIQUID to each 1000 kg of emulsion to provide 6.5 to 50 ppm active isothiazolones.

NOTE: The actual concentration of ACTICIDE® RS MICROBIOCIDE LIQUID required will depend upon such factors as specified substance to be treated, frequency of repeated microbial contamination expected and level of protection required. To ensure uniform mixing and in order to prevent localized shock to latex materials, add the ACTICIDE® RS MICROBIOCIDE LIQUID slowly to the formulation with agitation as early as possible during the production process.

PRESERVATIVE FOR PAINTS AND COATINGS:

ACTICIDE® RS MICROBIOCIDE LIQUID is recommended for the control of bacteria and fungi in water-based coatings, dispersed pigments and aqueous paints used for architectural purposes.

Add 430 grams to 1.65 kg of ACTICIDE® RS MICROBIOCIDE LIQUID to each 1000 kg of emulsion to provide 6.5 to 25 ppm active isothiazolone ingredients. A higher dosage rate of up to 3.3 kg per 1000 kg, providing up to 50 ppm active ingredients, may be required for storage during extremely high temperatures and humidity.

PRESERVATIVE FOR ADHESIVES, INKS, AND BUILDING MATERIALS:

ACTICIDE® RS MICROBIOCIDE LIQUID is also recommended as an in-container preservative in industrial products – mastics, caulks, joint cements, concrete additives, inks, slurries.

Add 430 grams to 3.3 kg of ACTICIDE® RS MICROBIOCIDE LIQUID to each 1000 kg of product to provide 6.5 to 50 ppm active isothiazolones.

PRESERVATIVE FOR PRINTING FLUIDS:

ACTICIDE® RS MICROBIOCIDE LIQUID is recommended for the control of bacteria and fungi in aqueous printing inks such as flexographic, gravure, screen and ink jet printing inks. Add 400 g – 1670 g of ACTICIDE® RS MICROBIOCIDE LIQUID to each 1000 kg of fluid to provide 400 to 1670 ppm product.

ACTICIDE® RS MICROBIOCIDE LIQUID is recommended for the control of bacteria and fungi in the manufacture and storage of printing ink components such as resins, plasticizers, water soluble dyes, pigments, gelling agents, waxes, surfactants, and thickeners. Add 400 g – 1670 g of ACTICIDE® RS MICROBIOCIDE LIQUID to each 1000 kg of fluid to provide 400 to 1670 ppm product.

ACTICIDE® RS MICROBIOCIDE LIQUID is recommended for the control of bacteria and

fungi in fountain solutions used for printing operations. ACTICIDE® RS MICROBIOCIDE LIQUID should only be used in those fountain solution concentrates, which are automatically diluted prior to use. ACTICIDE® RS MICROBIOCIDE LIQUID is also recommended as a preservative for photo processing chemicals such as stabilizer solutions. ACTICIDE® RS MICROBIOCIDE LIQUID is added to the fountain solution concentrate or processing chemical concentrate at a level to ensure that the final use dilution fluid will contain between 400 – 1670 ppm product.

PRESERVATIVE FOR HOUSEHOLD AND CONSUMER PRODUCTS:

ACTICIDE® RS MICROBIOCIDE LIQUID is recommended as an in-container preservative for the control of bacteria and fungi in industrial and household consumer products in all-purpose cleaners, polishes and anti-tarnish products, solid air freshener, carpet shampoos and pre- spotters.

Add 377 grams to 1.5 kg ACTICIDE® RS MICROBIOCIDE LIQUID to each 1000 kg of product to be treated.

ACTICIDE® RS MICROBIOCIDE LIQUID is recommended for the control of bacteria and fungi in liquid dishwashing detergents, automotive appearance products such as cleaners, wax and polishes and liquid laundry products.

Add 377 grams to 1.0 kg ACTICIDE® RS MICROBIOCIDE LIQUID to each 1000 kg of product to be treated.

To ensure uniform distribution, slowly disperse ACTICIDE® RS MICROBIOCIDE LIQUID into product with agitation. Mix thoroughly until evenly dispersed throughout the product.

ACTICIDE® RS MICROBIOCIDE LIQUID is recommended for the control of bacteria and fungi in packaged products, premoistened sponges and mops.

Add 377 grams to 1.5 kg ACTICIDE® RS MICROBIOCIDE LIQUID to each 1000 kg of water used to premoisten sponges and mops.

PRESERVATIVE FOR PESTICIDE PRODUCTS:

ACTICIDE® RS MICROBIOCIDE LIQUID is recommended for the control of bacteria and fungi in pesticide products. Add 400 grams to 3.04 kg ACTICIDE® RS MICROBIOCIDE LIQUID to each 1000 kg of product to be treated to provide 6 to 45 ppm active isothiazolones.

PRESERVATIVE FOR METALWORKING FLUIDS AND CONVEYER LUBRICANTS:

For non-food plant industrial use only:

ACTICIDE® RS MICROBIOCIDE LIQUID is recommended for the control of bacteria and

fungi in soluble and emulsifiable type aqueous metalworking fluids, such as emulsified petroleum oils, and formulations containing alcoholic fatty acid, sulfonated red oil, or naphthalene sulfonate and water based conveyor lubricants, These fluids are usually prepared by diluting the product concentrates 1:50 with water.

Metal Working Fluids: For the maintenance of a non-fouled system, use the diluted ACTICIDE® RS MICROBIOCIDE LIQUID at 22 ml per 100 L of emulsion every 4 weeks, or 22-100 ml per 100 L of emulsion every 8-12 weeks. For a noticeably fouled system use initial dose of 43-100 ml per 100 L of emulsion to be followed by subsequent dosages depending upon the treatment interval noted above. The higher dosage or increased frequency of treatment may be required depending on rate of dilution of the preservative with makeup fluid, the nature of severity of contamination, level of control required, filtration effectiveness, system design, etc.

The preservative should be dispensed into the use-dilution of the metalworking fluid by a metering pump and uniformly dispersed throughout the system.

Conveyor Lubricant: These fluids are prepared by diluting the product concentrated 1:50 to 1:400 with water. The concentration of ACTICIDE® RS MICROBIOCIDE LIQUID added to ensure final use dilution will contain 200-1000 ppm of ACTICIDE® RS MICROBIOCIDE LIQUID. Grossly contaminated systems must be cleared before treatment is begun. For initial dose in a notably fouled system meter 1 L of ACTICIDE® RS MICROBIOCIDE LIQUID per 1000 L of use dilution conveyor lubricant. If necessary, continue metering 400 ml to 1 L of ACTICIDE® RS MICROBIOCIDE LIQUID per 1000 L of conveyor lubricant until control is achieved and maintain the final use dilution at 400 – 1000 ppm. After establishing microbial control or to prevent fouling of a clean system, maintain ACTICIDE® RS MICROBIOCIDE LIQUID at 3-15 ppm active ingredient by metering 200 ml to 1 L per 1000 L of conveyor lubricant.

Avoid contamination of food in the application and storage of the product.

RECIRCULATING ELECTRODEPOSITION SYSTEMS:

ACTICIDE® RS MICROBIOCIDE LIQUID is recommended for the control of bacteria, fungi and algae in recirculating electrodeposition systems and associated rinse systems. ACTICIDE® RS MICROBIOCIDE LIQUID should be dispensed into the circulating rinse system, ultrafilter permeate or final distilled water rinse at a point to ensure uniform mixing.

Initial Dose: When the system is noticeably fouled, add 66.7 to 233.3 ppm of ACTICIDE® RS MICROBIOCIDE LIQUID (6.7 to 23.3 Litres) per 10,000 Litres of fluid in the system. Repeat until control is achieved.

Subsequent Dose: When microbial control is evident, add 33.3 to 100 ppm of ACTICIDE® RS MICROBIOCIDE LIQUID (3.3 to 10 Litres) per 10,000 Litres of fluid in the system weekly or as needed to maintain the system. A change of frequency of treatment may be required depending upon the rate of dilution of the preservative with the makeup fluid, the nature and severity of contamination, level of control required,

filtration effectiveness, system design, etc. Badly fouled systems must be cleaned before treatment is begun.

PROCESS WATER SYSTEMS

Pulp & Paper (non-food contact paper): ACTICIDE® RS MICROBIOCIDE LIQUID is recommended for the control of bacterial and fungal slime in the production of non-food contact paper.

ACTICIDE® RS MICROBIOCIDE LIQUID should be added at a point in the system to ensure uniform mixing such as the Beater, Hydropulper Fan or Broke Storage Pump.

Add 0.3 to 1.1 kg of ACTICIDE® RS MICROBIOCIDE LIQUID as supplied per tonne (dry basis) of pulp or paper produced, as a slug dose. If needed repeat daily. Badly fouled systems should be cleaned before initial treatment.

Cooling Towers: ACTICIDE® RS MICROBIOCIDE LIQUID is recommended for the removal of slime from fouled towers; add 0.33 to 8.67 kg of ACTICIDE® RS MICROBIOCIDE LIQUID as supplied per 10 tonnes of water in the system. This dosage should be repeated 2 to 3 times per week until fouling has been reduced to an acceptable level thus establishing microbial control.

ACTICIDE® RS MICROBIOCIDE LIQUID should be added to the basin, to the distribution boxes, or at any other point where it will be rapidly and uniformly dispersed throughout the system.

Air Washer Systems: Add to the air washer sump or chill water sump to ensure uniform mixing. Add 4.20 to 16.43 kg of ACTICIDE® RS MICROBIOCIDE LIQUID per 10,000 litres of water in the system, depending on the severity of contamination, to control bacteria, fungi and algae, which cause fouling in industrial air washer systems. ***Badly fouled systems must be cleaned before initial treatment. For use only in industrial air washing systems that maintain effective mist eliminating components.***

Intermittent or slug method:

Initial Dose: When the system is noticeably fouled, add 1.51 to 8.95 kg of ACTICIDE® RS MICROBIOCIDE LIQUID per 10,000 litres of water in the system. Repeat until control is achieved.

Subsequent Dose: When microbial control is evident, add 0.36 to 2.23 kg of ACTICIDE® RS MICROBIOCIDE LIQUID per 10,000 litres of water weekly or as needed to maintain control.

Continuous Feed Method:

Initial Dose: When the system is just noticeably fouled, add 1.51 to 8.95 kg of ACTICIDE® RS MICROBIOCIDE LIQUID per 10,000 litres of water in the system.

Subsequent Dose: Maintain this treatment level by adding a continuous feed of 0.36 to 2.23 kg of ACTICIDE® RS MICROBIOCIDE LIQUID per 10,000 litres of makeup water.

Oil Field Injection Waters: For the control of slime-forming bacteria and sulphate-reducing (sulphide-forming) bacteria in oil field injection waters. Slug treat with 70 to 360 ppm of ACTICIDE® RS MICROBIOCIDE LIQUID as supplied in accordance with the severity of bacterial contamination.

Add 180-360 ppm of ACTICIDE® RS MICROBIOCIDE LIQUID as supplied (17.8 to 35.6 litres per tonnes of water) at any point from which the material mixes uniformly throughout the system. Repeat the treatment after 3 days or as needed to achieve control. After achieving control, add 70 – 180 ppm of ACTICIDE® RS MICROBIOCIDE LIQUID as supplied (7.1 to litres per 100 tonnes of water) every seven days or as needed to maintain control.

STORAGE

Store away from children, food, drinking water or animal feed.

DISPOSAL

1. Triple or pressure-rinse the empty container. Add the rinsings to the treatment site.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills.

For Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.