

GROUP	9	12	FUNGICIDES
-------	---	----	------------

BUTTON Fungicide

**WETTABLE GRANULES
COMMERCIAL (AGRICULTURAL)**

Fungicide for control or suppression of listed diseases on listed crops.

ACTIVE INGREDIENT:

Cyprodinil	37.5%
Fludioxonil	25.0%

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF THE REACH OF CHILDREN**

CAUTION – EYE IRRITANT

REGISTRATION NO.: 34439
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **794 g – 100 kg**

SHARDA Cropchem Limited
2nd Floor, Prime Business Park
Dashrathlal Joshi Road
Vile Parle (West)
Mumbai - 400056, India

Canadian Agent:
SHARDA Cropchem Limited
63 Kingsview Blvd
Etobicoke, Ontario, CA
M9R 1V1
1-844-810-5720
1-416-840-5639

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FIRST AID

IN CASE OF POISONING, call a physician or poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes or prolonged contact with skin or clothing. Avoid inhalation of dust. May cause eye irritation.

DO NOT apply using handheld mist blower/airblast or handheld fogging equipment.

DO NOT enter or allow worker entry into treated areas during the intervals specified in the following table:

Crop	Post-application Activity	REI and/or PHI
Saskatoon berries	Hand harvesting, pruning and thinning	10 days
	Mechanical harvesting	1 day
	All other activities	12 hours
Small fruit vine climbing (crop subgroup 13-07F)	Cane turning, girdling and harvesting	7 days
	Training, thinning, hand pruning, tying and leaf pulling	2 days
	All other activities	12 hours
Carrots, root vegetables (crop subgroup 1B), ginseng, turnip greens, cabbage, leafy <i>Brassica</i> greens (crop subgroup 5B), succulent beans	Harvesting	7 days

(crop subgroup 6B), dried shelled peas and beans (crop subgroup 6C)	All other activities	12 hours
Bulb vegetables (crop group 3-07)	Harvesting	7 days
	Hand weeding	3 days
	All other activities	12 hours
Spinach	Harvesting	3 days
	All other activities	12 hours
Caneberries (crop subgroup 13-07A), bushberries, (crop subgroup 13-07B), low growing berries (crop subgroup 13-07G)	Harvesting	1 day
	All other activities	12 hours
Celery, field peppers, field-grown tomatoes	All activities, including harvesting	12 hours

DO NOT eat, drink or smoke while mixing or during application. Change and wash clothing immediately after use. Wash hands and face before eating, drinking, smoking or using the toilet. Store and wash all protective clothing separately from household laundry. Wash clothing in detergent and hot water before reuse.

Apply to agricultural crops only when there is low risk of drift to areas of human habitation or activity such as houses, cottages, schools and recreational areas. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact SHARDA Cropchem Limited at 1-844-810-5720.

PERSONAL PROTECTIVE EQUIPMENT

Wear coveralls over a long-sleeved shirt and long pants, socks, shoes and chemical-resistant gloves (such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, or Viton) during mixing and loading. Wear a long-sleeved shirt, long pants, socks and shoes during application, clean-up and repair. In addition, wear protective eyewear (goggles or faceshield) during open-cab application.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.

Fludioxonil is persistent and may carry over. It is recommended that this product not be used in areas treated with any products containing fludioxonil during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid applications to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

STORAGE

To prevent contamination, store this product away from food or feed.

DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

For Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

GROUP	9	12	FUNGICIDES
--------------	----------	-----------	-------------------

BUTTON Fungicide

WETTABLE GRANULES COMMERCIAL (AGRICULTURAL)

Fungicide for control or suppression of listed diseases on listed crops.

ACTIVE INGREDIENT:

Cyprodinil 37.5%
Fludioxonil 25.0%

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF THE REACH OF CHILDREN**

CAUTION – EYE IRRITANT

REGISTRATION NO.: 34439
PEST CONTROL PRODUCTS ACT

SHARDA Cropchem Limited
2nd Floor, Prime Business Park
Dashrathlal Joshi Road
Vile Parle (West)
Mumbai - 400056, India

Canadian Agent:
SHARDA Cropchem Limited
63 Kingsview Blvd
Etobicoke, Ontario, CA
M9R 1V1
1-844-810-5720
1-416-840-5639

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FIRST AID

IN CASE OF POISONING, call a physician or poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes or prolonged contact with skin or clothing. Avoid inhalation of dust. May cause eye irritation.

DO NOT apply using handheld mist blower/airblast or handheld fogging equipment.

DO NOT enter or allow worker entry into treated areas during the intervals specified in the following table:

Crop	Postapplication Activity	REI and/or PHI
Saskatoon berries	Hand harvesting, pruning and thinning	10 days
	Mechanical harvesting	1 day
	All other activities	12 hours
Small fruit vine climbing (crop subgroup 13-07F)	Cane turning, girdling and harvesting	7 days
	Training, thinning, hand pruning, tying and leaf pulling	2 days
	All other activities	12 hours
Carrots, root vegetables (crop subgroup 1B), ginseng, turnip greens, cabbage, leafy <i>Brassica</i> greens (crop subgroup 5B), succulent beans (crop subgroup 6B), dried shelled peas	Harvesting	7 days
	All other activities	12 hours

and beans (crop subgroup 6C)		
Bulb vegetables (crop group 3-07)	Harvesting	7 days
	Hand weeding	3 days
	All other activities	12 hours
Spinach	Harvesting	3 days
	All other activities	12 hours
Caneberries (crop subgroup 13-07A), bushberries, (crop subgroup 13-07B), low growing berries (crop subgroup 13-07G)	Harvesting	1 day
	All other activities	12 hours
Celery, field peppers, field-grown tomatoes	All activities, including harvesting	12 hours

Do not eat, drink or smoke while mixing or during application. Change and wash clothing immediately after use. Wash hands and face before eating, drinking, smoking or using the toilet. Store and wash all protective clothing separately from household laundry. Wash clothing in detergent and hot water before reuse.

Apply to agricultural crops only when there is low risk of drift to areas of human habitation or activity such as houses, cottages, schools and recreational areas. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Sharda Cropchem Limited at 1-844-810-5720.

PERSONAL PROTECTIVE EQUIPMENT

Wear coveralls over a long-sleeved shirt and long pants, socks, shoes and chemical-resistant gloves (such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, or Viton) during mixing and loading. Wear a long-sleeved shirt, long pants, socks and shoes during application, clean-up and repair. In addition, wear protective eyewear (goggles or faceshield) during open-cab application.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Fludioxonil is persistent and may carry over. It is recommended that this product not be used in areas treated with any products containing fludioxonil during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid applications to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

STORAGE

To prevent contamination, store this product away from food or feed.

DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

For Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

PRODUCT INFORMATION

BUTTON Fungicide is a broad-spectrum fungicide for the control or suppression of listed diseases.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Spray Equipment

Thorough coverage is necessary to provide good disease control. Applications using sufficient water volume to provide thorough and uniform coverage generally provide the most effective disease control.

For ground application equipment, a minimum of 200 L of water per hectare is recommended.

To avoid spray drift, do not apply when conditions favor drift beyond the target area. Avoid spray overlap, as crop injury may occur.

Equip sprayers with nozzles that provide accurate and uniform application. Calibrate sprayer before use.

DIRECTIONS FOR USE

General Information

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Feeding Restriction: **DO NOT** graze or feed treated field pea hay, field pea vines/silage, cowpea forage and cowpea hay to livestock.

DO NOT apply this product through any type of irrigation system.

DO NOT APPLY BY AIR.

Do not exceed 2.9 kg of product per hectare or 3 applications per year.

Within the stated range, apply the highest rate and the shortest interval under conditions of high disease pressures.

Make no more than two (2) sequential applications of BUTTON Fungicide before alternating with another registered fungicide with a different mode of action.

Ground Application

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) fine classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. DO NOT apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

DO NOT apply by air.

Mixing Procedures

Prepare only the amount of spray mixture that is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

1. Add 1/2 of the required amount of water to the mix tank.

2. With the agitator running, add the BUTTON Fungicide to the tank.
3. Continue agitation while adding the remainder of the water.
4. Begin application of the solution after the BUTTON Fungicide has completely dispersed into the mix water.
5. Maintain agitation until all of the mixture has been applied.

Spray Buffer Zones

A spray buffer zone is NOT required for uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands) and estuarine/marine habitats.

Method of application	Crop	Spray Buffer Zones (meters) Required for the Protection of:				
		Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths:		
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Turnip greens, Cabbage, Carrot, Crop Group 5B, Crop Group 3-07, Field Pepper, Tomatoes (field grown), Crop Subgroup 13-07A, Crop Subgroup 13-07B, Crop Subgroup 13-07G, Crop Subgroup 6C, The Following Succulent Beans Only: Bean (<i>Phaseolus</i> spp.) (includes lima bean, snap bean and wax bean), Bean (<i>Vigna</i> spp.) (includes blackeyed pea, asparagus bean), Broad bean (fava bean) (<i>Vicia faba</i>)	5	4	3	1	
	Celery Spinach Crop Group 1B (excluding carrots)	5	3	3	1	
Airblast	Crop subgroup 13-07F	Early airblast	20	10	10	4
		Late airblast	10	5	5	2

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

Rotational Crop Restrictions

Do not plant any other crop for a period of 30 days after harvest or crop failure unless BUTTON Fungicide is registered for that use.

MINOR USES

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than SHARDA Cropchem Limited under the User Requested Minor Use Label Expansion program. For these uses, SHARDA Cropchem Limited has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

Crop	Diseases	Product Rate in g/ha	Application Timing
Onions: Dry bulb, green, and onions grown for seed	Control of Botrytis Leaf Blight or Blast (<i>Botrytis squamosa</i>) Suppression of Purple Blotch (<i>Alternaria porri</i>)	775 to 975	Begin applications when conditions become favourable for disease but before infection. If favourable conditions persist make additional applications on 7 to 14 day intervals. Do not make more than 3 applications per year.
Strawberries	Control of Grey Mold (<i>Botrytis cinerea</i>)	975	Begin applications at or before bloom and continue on 7 to 10 day intervals. Do not make more than 3 applications per year.
	Suppression of powdery mildew (<i>Sphaerotheca macularis</i>) Control of anthracnose (<i>Colletotrichum acutatum</i>)	775 to 975	Begin applications at or before bloom and continue on 7 to 10 day intervals. Apply in sufficient water volume to obtain thorough coverage; a minimum of 200 L/ha is recommended. Do not make more than 3 applications per year. In the absence of a viable registered fungicide alternative for the specific crop to be treated, the maximum number of applications is limited to 2 per year. Resistance Management: After 2 applications of BUTTON Fungicide, alternate with another fungicide with a different mode of action for 2 applications.
Blueberries: highbush and lowbush	Control of Anthracnose (<i>Colletotrichum acutatum</i>) Botrytis Fruit Rot (<i>Botrytis cinerea</i>) Control of the primary	775 to 975	Make the first application during early bloom. A second application may be made 7 to 10 days later. A third application can be made if conditions remain favourable for disease development. Apply in sufficient water volume to obtain thorough coverage; a minimum of 200 L/ha

	<p>phase infection of mummy berry/Monilinia (<i>Monilinia vaccinia-corymbosi</i>)</p> <p>Suppression of sclerotinia berry drop (<i>Sclerotinia sclerotiorum</i>) on lowbush blueberry only</p>		<p>is recommended.</p> <p>Do not make more than 3 applications per year. In the absence of a viable registered fungicide alternative for the specific crop to be treated, the maximum number of applications is limited to 2 per year.</p>
Caneberries: Raspberries, Blackberries	Control of Botrytis Fruit Rot (<i>Botrytis cinerea</i>)	775 to 975	Make the first application during early bloom. A second application may be made 7 to 10 days later. A third application can be made if conditions remain favourable for disease development.
Carrots	Control of Alternaria Leaf Blight (<i>Alternaria dauci</i>)	775 to 975	<p>The first application should be made when the disease first appears; a second application should be made 7-10 days later. One more application may be made a minimum of 7 days later if conditions remain favourable for disease development.</p> <p>Do not make more than 3 applications per year.</p> <p>Apply in sufficient water volume to obtain thorough coverage; a minimum spray volume of 200 L/ha is recommended.</p>
Mustard Greens, Turnip Greens	Control of Powdery Mildew (<i>Erysiphe polygoni</i> , <i>E. heraclei</i>)	775 to 975	<p>The first application should be made when disease first appears and continue on 7 to 10 day intervals. Apply in sufficient water volume to obtain thorough coverage; a minimum spray volume of 200 L/ha is recommended.</p> <p>Do not make more than 3 applications per year.</p>
Cabbage	Suppression of Alternaria Leaf Blight (<i>Alternaria brassicicola</i>)	775 to 975	<p>The first application should be made when disease first appears and continue on 7 to 10 day intervals. Apply in sufficient water volume to obtain thorough coverage; a minimum spray volume of 200 L/ha is recommended.</p> <p>Do not make more than 3 applications per year.</p>
Includes: Broccoli raab (rapini), Chinese cabbage (bok choy), collards, kale, mizuna, mustard greens, mustard spinach, rape greens	Suppression of Alternaria leaf blight (<i>Alternaria brassicicola</i>)	775 to 975	The first application should be made when disease first appears and continue on 7 to 10 day intervals. Apply in sufficient water volume to obtain thorough coverage; a minimum spray volume of 200 L/ha is recommended.

			Do not make more than 3 applications per year.
<p>Crop Group 3-07 Complete list of crops: Garlic (bulb and greatheaded), Leek, Onion (bulb, green, Welsh, potato, Chinese, tree, Beltsville bunching, fresh, macrostem, pearl), Shallot (bulb and fresh leaves), Chive (fresh leaves, Chinese), Daylily (bulb), Elgans hosta, Fritillaria, Garlic (serpent, bulb), Kurrat, Lady's Leek, Lily, Wild Leek</p>	<p>Control of Botrytis Leaf Blight or Blast (<i>Botrytis squamosa</i>)</p> <p>Suppression of Purple Blotch (<i>Alternaria porri</i>)</p>	775 to 975	<p>Begin applications when conditions become favourable for disease but before infection. If favourable conditions persist make additional applications on 7 to 14 day intervals.</p> <p>Apply in sufficient water volume to obtain thorough coverage.</p> <p>Do not make more than 3 applications per year. Make no more than two (2) sequential applications of BUTTON Fungicide before alternating with another registered fungicide with a different mode of action.</p>
<p>Saskatoon berries</p>	<p>Suppression of Entomosporium leaf spot (<i>Entomosporium mespili</i>)</p>	775 to 975	<p>The first application should be made during early bloom. A second application may be made 7 to 10 day later. A third application can be made if conditions remain favourable for disease development.</p> <p>Use the highest rate and the shortest application interval under conditions that promote rapid disease development or if there is a history of high disease pressure in the field.</p> <p>Apply in sufficient water to ensure thorough coverage (187-935 L/ha).</p> <p>Do not make more than 3 applications per year.</p>

<p>CROP SUBGROUP 6C: Dried shelled pea and bean (except soybean) Includes: chickpea (garbanzo bean) (<i>Cicer arietinum</i>); beans (<i>Lupinus</i> spp. including grain lupin, sweet lupin, white lupin, white sweet lupin); beans (<i>Phaseolus</i> spp. including field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); broad bean (fava bean) (<i>Vicia faba</i>); beans (<i>Vigna</i> spp. adzuki bean, black-eyed pea, catjang, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); Guar (<i>Cyamopsis tetragonoloba</i>); Lablab bean or Hyacinth bean (<i>Lablab purpureus</i>); Lentil (<i>Lens esculenta</i>); Pea (<i>Pisum</i> spp.) (includes field pea); and Pigeon pea (<i>Cajanus cajan</i>)</p> <p>The Following Succulent Beans: Bean (<i>Phaseolus</i> spp.) (includes lima bean, snap bean and wax bean); Bean (<i>Vigna</i> spp.) (includes blackeyed pea, asparagus bean); Broad bean (fava bean) (<i>Vicia faba</i>)</p>	<p>Control of White mold (<i>Sclerotinia sclerotiorum</i>) and Grey mold (<i>Botrytis cinerea</i>)</p>	<p>775 - 975</p>	<p>Begin applications prior to or at the onset of disease and repeat applications at 7 day intervals if conditions remain favourable for disease development.</p> <p>For white mold control, make the first application at 10-20% bloom. In some locations, a single application at this timing will provide adequate disease control.</p> <p>Apply the highest application rate under conditions of high disease pressure.</p> <p>Apply in sufficient water to ensure thorough coverage:</p> <p>Apply in 175 – 225 L of water /ha. Do not apply more than 1.1 kg ai/ha of cyprodinil products and 0.73 kg ai/ha of fludioxonil containing products per crop.</p> <p>Do not make more than 3 applications of BUTTON fungicide per year.</p>
<p>Crop Group 1B: Includes: garden beet, burdock, carrot, celeriac, turnip-rooted chervil, chicory, ginseng, horseradish, turnip-rooted parsley, parsnip, radish, oriental radish, rutabaga, salsify, black salsify, Spanish salsify, skirret, turnip</p>	<p>Botrytis grey mold (<i>Botrytis cinerea</i>)</p> <p>White mold (<i>Sclerotinia sclerotiorum</i>)</p>	<p>775 - 975</p>	<p>Begin applications prior to or at the onset of disease and repeat applications on a 7-10 day interval if conditions remain favorable for disease development.</p> <p>Do not make more than 2 applications of BUTTON Fungicide per year.</p> <p>Do not allow cattle or other livestock to feed upon the leaves of root vegetables.</p>
<p>Celery</p>	<p>Anthraco­nose (<i>Colletotrichum acutatum</i>)</p>	<p>775-975</p>	<p>Begin applications prior to or at the onset of disease and repeat applications on a 7-10 day interval if conditions remain favorable for disease development.</p> <p>Do not make more than 2 applications per year.</p>

Ginseng	Alternaria leaf blight (<i>Alternaria panax</i>)	975	<p>Begin applications prior to or at the onset of disease and repeat applications on a 7-10 day interval if conditions remain favorable for disease development.</p> <p>Do not make more than 2 applications of BUTTON Fungicide per year.</p> <p>Do not allow cattle or other livestock to feed upon the leaves of root vegetables.</p>
	<p>Cylindrocarpon root rot (<i>Cylindrocarpon destructans</i>) (Suppression)</p> <p>Rhizoctonia root rot (<i>Rhizoctonia solani</i>)</p>	775 - 975	<p>Begin applications prior to or at the onset of disease and repeat applications on a 14-21 day interval if conditions remain favorable for disease development.</p> <p>While treating for other ginseng diseases listed on the label, BUTTON Fungicide will also provide control of Rhizoctonia and suppression of Cylindrocarpon if present. If only treating for Rhizoctonia and/or Cylindrocarpon, use SCHOLAR® 230SC FUNGICIDE or other fungicides registered for the targeted diseases.</p> <p>Apply by broadcast ground application in sufficient water volume to obtain thorough coverage and penetration to the soil and root zone.</p> <p>Do not make more than 2 applications of BUTTON Fungicide per year.</p> <p>Do not allow cattle or other livestock to feed upon the leaves of root vegetables.</p>
Grape	Control of botrytis bunch rot caused by <i>Botrytis cinerea</i>	775-975	<p>Begin applications at early bloom. One additional application may be made at berry touch, veraison, or preharvest. Botrytis bunch rot is most effectively controlled by ground application, using sufficient water volume to provide thorough coverage. Thorough coverage of bunches is essential.</p> <p>Application interval is 21 days.</p> <p>Do not make more than 2 applications of BUTTON Fungicide per year.</p> <p>Resistance Management: After a maximum of 2 applications of BUTTON Fungicide, alternate with another registered fungicide with a different mode of action for 2 applications.</p>

			<p>Do not use an adjuvant.</p> <p>Do not apply more than 0.75 kg ai/ha of cyprodinil products and 0.50 kg ai/ha fludioxonil containing products per crop.</p>
Spinach	<p>Anthracnose (<i>Colletotrichum dematium</i> f. sp. <i>spinaciae</i>) and grey mold (<i>Botrytis cinerea</i>)</p>	775-975	<p>Begin applications prior to or at the onset of disease and repeat applications on a 7-10 day interval if conditions remain favorable for disease development.</p> <p>Apply the highest application rate under conditions of high disease pressure.</p> <p>Ground Application: Apply in 175 – 225 L of water per hectare.</p> <p>DO NOT make more than 2 applications of BUTTON Fungicide per year.</p>
Field Pepper	<p>Anthracnose (<i>Colletotrichum</i> spp.) Grey mould (<i>Botrytis cinerea</i>)</p>	775-975	<p>Begin applications when conditions become favorable for disease but before infection. Apply the highest rate under conditions of high disease pressure. If favorable conditions persist, make additional applications on a 7 to 10 day interval. Make no more than two sequential applications before using another registered fungicide with a different mode of action.</p> <p>A minimum spray volume of 200 L water/ha is recommended.</p> <p>DO NOT make more than 3 applications of BUTTON Fungicide per year.</p>
Tomatoes (field grown)	<p>Botrytis grey mold (<i>Botrytis cinerea</i>)</p>	775-975	<p>Begin applications prior to or at the onset of disease and repeat applications at 7 to 10 day intervals if conditions remain favourable for disease development.</p> <p>Apply the highest application rate under conditions of high disease pressure.</p> <p>Apply in sufficient water to ensure thorough coverage: Ground: 200 L of water /ha After 2 applications of BUTTON Fungicide, alternate with another fungicide with a different mode of action for 1 application.</p> <p>Do not make more than 3 applications of BUTTON Fungicide per year.</p>

<p>Crop subgroup 13-07A - Caneberry Subgroup Raspberry, Blackberry, Loganberry (Including cultivars, varieties and/or hybrids of these)</p>	<p>Control of Botrytis Fruit Rot (<i>Botrytis cinerea</i>)</p>	<p>775 to 975</p>	<p>Make the first application during early bloom. A second application may be made 7 to 10 days later. A third application can be made if conditions remain favourable for disease development</p> <p>Do not make more than 3 applications of BUTTON Fungicide per year. In the absence of a viable registered fungicide alternative for the specific crop to be treated, the maximum number of applications is limited to 2 per year.</p> <p>Resistance Management: After 2 applications of BUTTON Fungicide, alternate with another fungicide with a different mode of action for 2 applications.</p> <p>Apply in sufficient water volume to obtain thorough coverage; a minimum of 200 L/ha is recommended.</p>
<p>Crop subgroup 13-07B - Bushberry Subgroup Blueberry (highbush and lowbush), Saskatoon berry, Currant (black and red), Elderberry, Gooseberry, Huckleberry, Aronia berry, Buffalo currant, Chilean guava, European barberry, Highbush cranberry, Honeysuckle (edible), Jostaberry, Lingonberry, Native currant, Salal, Sea buckthorn (Including cultivars, varieties and/or hybrids of these)</p>	<p>Control of Anthracnose (<i>Colletotrichum acutatum</i>) on blueberries</p> <p>Suppression of Entomosporium leaf spot (<i>Entomosporium mespili</i>) on Saskatoon berries</p> <p>Control of Botrytis Fruit Rot (<i>Botrytis cinerea</i>) on bushberries</p>	<p>775 to 975</p>	<p>Make the first application during early bloom. A second application may be made 7 to 10 days later. A third application can be made if conditions remain favourable for disease development.</p> <p>For Saskatoon Berry: Use the highest rate and shortest application interval under conditions that promote rapid disease development or if there is a history of high disease pressure in the field.</p> <p>Apply in sufficient water to ensure thorough coverage (187-935 L/ha).</p> <p>Do not make more than 3 applications of BUTTON Fungicide per year. In the absence of a viable registered fungicide alternative for the specific crop to be treated, the maximum number of applications is limited to 2 per year.</p> <p>Resistance Management: After 2 applications of BUTTON Fungicide, alternate with another fungicide with a different mode of action for 2 applications.</p>
<p>Crop subgroup 13-07F - Small Fruit Vine Climbing Subgroup Grape, Armur river grape, Hardy kiwifruit, Maypop, Schisandra berry (Including cultivars,</p>	<p>Control of Bunch Rot (<i>Botrytis cinerea</i>)</p>	<p>775 to 975</p>	<p>Begin applications at early bloom. One additional application may be made at berry touch, veraison, or preharvest. Botrytis Bunch Rot is most effectively controlled by ground application, using sufficient water volume to provide thorough coverage. Thorough coverage of bunches is essential.</p>

varieties, and/or hybrids of these, excluding fuzzy kiwifruit)			<p>Application Interval 21 days. Do not use an adjuvant.</p> <p>Do not make more than two applications of BUTTON Fungicide per year.</p> <p>Do not apply more than 0.73 kg ai/ha of cyprodinil products and 0.49 kg ai/ha of fludioxonil containing products per crop.</p> <p>Resistance Management: After 2 applications of BUTTON Fungicide, alternate with another fungicide with a different mode of action for 2 applications.</p>
<p>Crop subgroup 13-07G - Low Growing Berry Subgroup Bearberry, Bilberry, Blueberry (lowbush), Cloudberry, Lingonberry, Muntries, Partridgeberry, Strawberry (Including cultivars, varieties, and/or hybrids of these, excluding cranberries)</p>	<p>Control of Grey Mould (<i>Botrytis cinerea</i>)</p>	<p>775 to 975</p>	<p>Begin applications at or before bloom and continue on 7 to 10 day intervals.</p> <p>Apply in sufficient water volume to obtain thorough coverage; a minimum of 200 L/ha is recommended.</p> <p>Do not make more than 3 applications of BUTTON Fungicide per year. In the absence of a viable registered fungicide alternative for the specific crop to be treated, the maximum number of applications is limited to 2 per year.</p> <p>Resistance Management: After 2 applications of BUTTON Fungicide, alternate with another fungicide with a different mode of action for 2 applications.</p>

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, BUTTON Fungicide contains a Group 9 and Group 12 Fungicide. Any fungal population may contain individuals naturally resistant to BUTTON Fungicide and other Group 9 and Group 12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of BUTTON Fungicide or other Group 9 and Group 12 fungicides with different groups that control the same pathogens.

Avoid application of more than the maximum number of sprays listed in the label and do not apply more than 2 consecutive sprays of BUTTON Fungicide or other fungicides in the same group in a season.

Use tank mixtures with fungicide from a different group that is effective on the target pathogen when such use is permitted.

Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for sign of resistance development. Notify Sharda Cropchem Limited if reduced sensitivity of the pathogen to BUTTON Fungicide is suspected.

If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information or to report suspected resistance, contact company representatives at 1-844-810-5720.