

Bag Label

GROUP	3	8	HERBICIDES
--------------	----------	----------	-------------------

YIELDSHIELD RESILIENT XP SOILSORB

Granular herbicide for Fall and Spring treatment to control wild oats and wild millet (green and yellowfoxtail) in Spring and Durum Wheat, Spring Barley, Rapeseed (including Canola), Flax (not including low linolenic acid varieties), and Mustard.
Also provides suppression of several annual broadleaved weeds.

COMMERCIAL (AGRICULTURAL)

Active Ingredients: TRIALLATE.....10%
TRIFLURALIN.....4%

Granular

REGISTRATION NO. 34566 PEST CONTROL PRODUCTS ACT

READ THIS LABEL AND ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

POTENTIAL SKIN SENSITIZER

NET CONTENTS: 22.7-1000 kg

IMPORTANT NOTE: Before use, recalibrate application equipment.

Read NOTICE before buying or using. If notice terms are not acceptable, return at once unopened.

Rath Holdco Inc.
346 Brighton Gate
Saskatoon, Saskatchewan
S7V 0P9 Canada
1-888-466-0056

NOTE: Populations of green foxtail tolerant to trifluralin have developed in a number of fields in Western Canada which have a long history of repeated trifluralin use. Trifluralin or ethalfluralin containing products (i.e. RIVAL®, TREFLAN®, TRIFLUREX®, YIELDSHIELD RESILIAN XP SOILSORB and EDGE®) will not control trifluralin-tolerant green foxtail. To delay selection or reduce the spread of trifluralin-tolerant green foxtail, avoid use of these products repeatedly in the same field, or use a separate non-trifluralin herbicide application for control of trifluralin-tolerant green foxtail.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.
MAY CAUSE SKIN AND EYE IRRITATION.
MAY BE HARMFUL IF SWALLOWED.
POTENTIAL SKIN SENSITIZER
Wash thoroughly after handling.
Do not take internally.
Do not get in eyes or on skin or on clothing.
Launder clothes before reuse.

Wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested when handling. For application to stored grain or when using a dust blower, wear a NIOSH-approved supplied-air respirator (SAR) operated in a pressure-demand or other positive-pressure mode during mixing, loading and application.

All handlers must wear chemical-resistant coveralls over long-sleeved shirt and long pants, chemical-resistant gloves, socks, and chemical-resistant footwear during mixing, loading, application, clean-up and repair.

Restricted-Entry Interval

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

ENVIRONMENTAL PRECAUTIONS:

Runoff

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

Leaching

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Volatilization

The active ingredient contained in this product is known to volatilize. To reduce the atmospheric loading of triallate, effort should be made to reduce the volatilization such as the following:

- Application should occur when soil temperatures are less than 4°C or less.
- To minimize the release of trifluralin into the environment due to volatilization, trifluralin should only be applied on cool mornings and evenings when air temperatures are 15°C or lower. To further reduce volatilization to the atmosphere, incorporation into the soil should occur concurrently with application.

CAUTION: Do not graze the treated crops or cut for hay; there are insufficient data available to support such uses.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies involving this product, call CANUTEC at 1-613-996-6666.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

Keep YieldShield RESILIENT XP Soilsorb bag closed to prevent spills and contamination.

Store this product away from food or feed.

DISPOSAL

Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.

2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/ territorial requirements.

Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer and the provincial/ territorial regulatory agency. Contact the manufacturer and the provincial/ territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

BOOKLET

GROUP	3	8	HERBICIDES
--------------	----------	----------	-------------------

YIELDSHIELD RESILIAN T XP SOILSORB

Granular herbicide for Fall and Spring treatment to control wild oats and wild millet (green and yellowfoxtail) in Spring and Durum Wheat, Spring Barley, Rapeseed (including Canola), Flax (not including low linolenic acid varieties), and Mustard.

Also provides suppression of several annual broadleaved weeds.

COMMERCIAL (AGRICULTURAL)

Active Ingredients: TRIALLATE.....10%
TRIFLURALIN.....4%

Granular

REGISTRATION NO. 34566 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND THIS BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

POTENTIAL SKIN SENSITIZER

NET CONTENTS: 22.7-1000 kg

IMPORTANT NOTE: Before use, recalibrate application equipment.

Read NOTICE before buying or using. If notice terms are not acceptable, return at once unopened.

Rath Holdco Inc.
346 Brighton Gate
Saskatoon, Saskatchewan
S7V 0P9 Canada
1-888-466-0056

NOTE: Populations of green foxtail tolerant to trifluralin have developed in a number of fields in Western Canada which have a long history of repeated trifluralin use. Trifluralin or ethalfluralin containing products (i.e. RIVAL(R), TREFLAN(R), TRIFLUREX(R), YIELDSHIELD RESILIENT XP SOILSORB and EDGE(R)) will not control trifluralin-tolerant green foxtail. To delay selection or reduce the spread of trifluralin-tolerant green foxtail, avoid use of these products repeatedly in the same field, or use a separate non-trifluralin herbicide application for control of trifluralin-tolerant green foxtail.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.
MAY CAUSE SKIN AND EYE IRRITATION.
MAY BE HARMFUL IF SWALLOWED.
POTENTIAL SKIN SENSITIZER
Wash thoroughly after handling.
Do not take internally.
Do not get in eyes or on skin or on clothing.
Launder clothes before reuse.

Wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested when handling. For application to stored grain or when using a dust blower, wear a NIOSH-approved supplied-air respirator (SAR) operated in a pressure-demand or other positive-pressure mode during mixing, loading and application.

All handlers must wear chemical-resistant coveralls over long-sleeved shirt and long pants, chemical-resistant gloves, socks, and chemical-resistant footwear during mixing, loading, application, clean-up and repair.

Restricted-Entry Interval

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

ENVIRONMENTAL PRECAUTIONS:

Runoff

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

Leaching

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Volatilization

The active ingredient contained in this product is known to volatilize. To reduce the atmospheric loading of triallate, effort should be made to reduce the volatilization such as the following:

- Application should occur when soil temperatures are less than 4°C or less.
- To minimize the release of trifluralin into the environment due to volatilization, trifluralin should only be applied on cool mornings and evenings when air temperatures are 15°C or lower. To further reduce volatilization to the atmosphere, incorporation into the soil should occur concurrently with application.

CAUTION: Do not graze the treated crops or cut for hay; there are insufficient data available to support such uses.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies involving this product, call CANUTEC at 1-613-996-6666.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

GENERAL INFORMATION

This granular herbicide is recommended for wild oat and wild millet (green and yellow foxtail) control in spring and durum wheat, spring barley, rapeseed (including canola), flax (not including low linolenic acid varieties) and mustard only.

This product will also provide suppression of the following annual broadleaved weed species: lamb's quarters, kochia, redroot pigweed, Russian thistle and wild buckwheat. Other crops should not be treated with this product as injury may occur.

This product should be applied according to the recommendations on this label. See the "**Directions For Use**" section for additional information.

Application equipment must be properly calibrated. Application of an excessive rate of granular YieldShield RESILIAN XP Soilsorb may injure the crop, whereas application of too low a rate may result in poor weed control. Before applying this product, be sure the soil is in good working condition. Clay soils may require additional tillage to obtain good tilth. Application to a field which is wet, lumpy, rough or ridged will result in reduced weed control and promote crop thinning. See the "**Field Preparation**" section for additional information.

The initial incorporation of this product into the soil must be completed within 24 hours after application, except for the Fall surface treatment. For Fall surface guidelines, see the "**Fall Treatment (surface)**" section of this label. The incorporation should not be deeper than 5 cm as deeper incorporation tends to dilute the product, thus decreasing weed control and increasing the risk of crop injury. See the "**How To Incorporate**" section for additional information.

When using granular YieldShield RESILIAN XP Soilsorb, an untreated strip should be left for proof of results. Wild oat control may be evaluated by removing a surface 2.5 to 5 cm of soil at the time of germination to inspect the number of wild oats that were killed before emergence. This product is primarily absorbed by wild oat shoots from the treated layer of the soil. Wild oats are usually controlled before they emerge through the soil. Occasionally, and particularly under dry conditions and/or cool soil temperatures, some wild oats may emerge and reach the 3 to 4 leaf stage before dying. Wild Millet seedlings will generally be controlled as they germinate.

Rainfall of at least 1.5 cm within 2 weeks after seeding in the spring is required to ensure maximum performance.

Under conditions of prolonged cool soil temperatures at the time of germination, or extreme drought in spring, this product may not maintain the usual high standard of weed control.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, YieldShield RESILIAN XP Soilsorb is both a Group 3 and a Group 8 herbicide. Any weed population may contain or develop plants naturally resistant to YieldShield RESILIAN XP Soilsorb and other Group 3 or Group 8 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of YieldShield RESILIAN XP Soilsorb or other Group 3 or Group 8 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is

permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.

- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Rath Holdco Inc. at 1-888-466-0056.

HOW TO APPLY YIELDSHIELD RESILIAN XP SOILSORB AND CALIBRATE APPLICATION EQUIPMENT

Granular YieldShield RESILIAN XP Soilsorb must be applied through a ground broadcast applicator such as a grass seed attachment or a specially designed ground broadcast applicator. This product can also be applied by an airplane capable of applying small quantities of granules evenly.

GROUND EQUIPMENT: It is important that the applicator be calibrated properly to deliver the desired amount of this product to avoid applying too little or too much material. To give even distribution, scatter plates (similar to those used for applying granules in band) must be attached to each delivery tube or outlet in such a manner to give overall coverage. To calibrate, attach cloth or strong plastic bag over each spreader plate or delivery tube. Operate over normal terrain to be treated at 6 to 8 km/h. Collect the granules from all outlets after covering the desired distance.

For example: If a 4.25 m applicator is being used, stake off a distance of 200 m in the field to be treated. After attaching a bag to each outlet, collect the granules while driving the distance. Check to see that each outlet disperses the same amount of granules. Combine all samples and weigh. For the above set of conditions, the quantities of granules that should be collected for the following recommended rates for broadcast treatment are:

Recommended Rate

Amount to be collected	11 kg/ha 0.9 kg	14 kg/ha 1.2 kg	17 kg/ha 1.5 kg
---------------------------	--------------------	--------------------	--------------------

If more or less than the desired quantity is collected adjust opening accordingly and again collect the granules while driving the staked off distance. Continue this procedure until the proper amount is delivered.

To calibrate applicators using air pressure to disperse the granules, follow directions as outlined by the manufacturer.

AERIAL APPLICATION

Do not use human flaggers.

Directions for Use

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices, or equivalent electronic positioning systems (GPS).

GRANULAR YIELDSHIELD RESILIAN XP SOILSORB RATES (kg/ha) AERIAL APPLICATION					
Crop	Less than 2% OM	2-4% OM	4-6% OM	6-8% OM	Greater than 8% OM
Spring and Durum Wheat	-	11	14	14	17
Spring Barley	11	14	14	17	17

Rapeseed, (including Canola) Flax**, Mustard	14	14	14	17	17
----------------------------------------------------------	----	----	----	----	----

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even target coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Do not apply to any body of water. Avoid drifting of granules onto any body of water or other non-target areas.

Do not apply during periods of dead calm or when wind velocity and direction pose a risk of drift. Do not spread when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of pre-mixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-888-466-0056 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

For aerial applications, attachments designed for applying low volumes of granules must be used. Appropriate modifications should be made to equipment to ensure precise application. It is also necessary to properly calibrate equipment to ensure uniformity of application and properrate.

See the “**Directions for Use**” section of this label for application rates and additional information.

FIELD PREPARATION

Before applying this product, be sure the soil is in good working condition. All deep tillage by cultivation or double disc implements must be completed prior to application of this product. If excessive trash is present, additional tillage or other appropriate trash management practice will be necessary to reduce the trash cover to an acceptable level before application as too much trash can make uniform incorporation difficult.

If the soil is excessively wet or lumpy, making proper incorporation difficult, cultivation with suitable tillage equipment may be required before application and incorporation to improve the soil conditions. If ridging is a problem, suitable tillage is recommended prior to application to eliminate the ridging and ensure uniform distribution of the herbicide. Fall tillage is not necessary with the Fall surface treatment.

HOW TO INCORPORATE

This product must be incorporated into the soil, using suitable tillage equipment, in order to provide weed control. Two incorporation operations at right angles to one another are necessary for thorough mixing of this product in the soil. The first incorporation should be completed within 24 hours of application, except for the Fall Surface application. The second incorporation may then be done during the fall or delayed until the spring.

Do not incorporate this product more than 5 cm. This can be accomplished by setting the tillage equipment to work the soil not deeper than 7.5 cm to 10 cm. Shallow incorporation is necessary to prevent dilution of the product, thus decreasing weed control and increasing the risk of crop injury.

Harrowing does not provide effective incorporation if compact soil prevents penetration of harrow teeth or if trash accumulates in the harrow section or if harrows bounce.

FALL INCORPORATION

This product must be incorporated as described in the "**How To Incorporate**" section of this label.

Only one incorporation is required in the fall, except for the fall surface treatment. The second incorporation may be done either in the fall (prior to soil freeze-up) or in the spring. If the second incorporation is done in the spring it may be done any time until crop emergence. If the second incorporation is conducted after seeding it should be done with a harrow or other suitable tillage equipment adjusted to a depth so as not to disturb the seed. If ridging is a problem after seeding, one harrowing is recommended to eliminate the ridging and ensure uniform distribution of the herbicide.

SPRING INCORPORATION

Before Seeding

This product must be incorporated as described in the "**How To Incorporate**" section of this label.

This product may be incorporated prior to the seeding of any labelled crop. See the "**Directions For Use**" section for exceptions. The second incorporation may then be done any time until crop emergence. If the second incorporation is conducted after seeding it should be done with a harrow or other suitable tillage equipment adjusted to a depth so as not to disturb the seed. If ridging is a problem after seeding, one harrowing is recommended to eliminate the ridging and ensure uniform distribution of the herbicide.

NOTE

FOR THE BEST RESULTS ON SPRING APPLICATIONS, DELAY THE SECOND INCORPORATION FOR AT LEAST 48 HOURS FOLLOWING THE FIRST. Under conditions of prolonged cool soil temperatures or extreme drought a longer delay may be utilized to maximize control.

DIRECTIONS FOR USE

DO NOT apply more than one application per year.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Discard clothing or other absorbent materials that have been heavily contaminated with the product. DO NOT reuse them.

Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

To minimize the release of trifluralin into the environment due to volatilization, trifluralin should only be applied on cool mornings and evenings when air temperatures are 15°C or lower. To further reduce volatilization to the atmosphere, incorporation into the soil should occur concurrently with application.

This product can be applied prior to the seeding of Spring and Durum wheat, spring barley, rapeseed (including canola), mustard and flax (not including low linolenic acid varieties).

Fall Treatment (Incorporated)

Fall applications of this product must be made after September 15 and until soil freeze-up. Applications made before September 15 may result in reduced effectiveness of the herbicide. See the "**Field Preparation**", "**How to Incorporate**" and "**Important Seeding Requirements**" sections of this label for additional information.

GRANULAR YIELDSHIELD RESILIAN XP SOILSORB RATES (kg/ha) (Incorporated)						
Crop	Less than 2% OM	2-4% OM	4-6% OM	6-8% OM	Greater than 8% OM	Seeding Depth
Spring and Durum Wheat	-	11	14	14	17	6-7.5cm*
Spring Barley	11	14	14	17	17	6-7.5cm
Rapeseed, (including Canola) Flax**, Mustard	14	14	14	17	17	As desired

*This seeding depth will ensure placement below the treated layer.

DO NOT SEED DEEPER THAN 7.5 cm.

** Does not include low linolenic acid varieties.

OM = Organic Matter.

Fall Treatment (Surface)

Where fields are prone to water and/or wind erosion, and fall tillage is therefore undesirable, Fall Surface applications should be made when average soil temperature at the 5 cm depth is 4°C or less and within 3 weeks of soil freeze up. This situation generally occurs by October 1 across the prairies.

Apply at rates shown in the table below. Applications should be made to standing stubble, chemicalfallow or summerfallow fields in a state of low soil erodibility. Avoid smooth, hard-packed soil conditions in summerfallow which may allow granules to drift.

Incorporate twice at right angles in spring. Early seeding and a delayed second incorporation of 48 hours or more will provide best results. Surface applications should not be made to fields covered with snow or excessive crop residue, which will not allow granule contact with soil. Under excessively warm and/or wet conditions between application and crop emergence, control may be reduced.

For best results on heavy wild oat infestations, use the incorporated treatment only.

Soil colour may not be a precise indication of soil organic matter content. Ensure that the application rate chosen from the table below is appropriate for your soil type.

FALL GRANULAR YIELDSHIELD RESILIAN XP SOILSORB RATES (kg/ha) (Surface)					
Crop	Less than 2% OM	2-4% OM	4-6% OM	Greater than 6% OM	Seeding Depth
Spring and Durum Wheat	-	11	14	14	6-7.5cm*
Spring Barley	11	14	14	17	6-7.5cm
Rapeseed, (including Canola) Flax**, Mustard	14	14	14	17	As desired

*This seeding depth will ensure placement below the treated layer.

DO NOT SEED DEEPER THAN 7.5 cm.

** Does not include low linolenic acid varieties.

OM = Organic Matter.

Note: Weed control may be slightly reduced with a fall surface treatment

Spring Treatment

This product can be applied before seeding labelled crops, except as noted in the table below.

See the "Field Preparation", "How To Incorporate" and "Important Seeding Requirements" sections of this label for additional information.

SPRING GRANULAR YIELDSHIELD RESILIAN XP SOILSORB RATES (kg/ha)					
Crop	Application Timing	Less than 2% OM	2-4% OM	4-6% OM	Greater than 6% OM
Spring and Durum Wheat	Before Seeding	-	-	11	14
Spring Barley	Before Seeding	-	11	14	17
Rapeseed, (including Canola) Flax*, Mustard	Before Seeding	14	14	17	17

* Does not include low linolenic acid varieties.

OM = Organic Matter

IMPORTANT SEEDING REQUIREMENTS

1. For Wheat or Barley, hoe drills and double disc press drills can be used on all soil types that are indicated in the rate charts.
2. Wheat or barley may be seeded into an application of this product using a disc seeder or air seeder, only in the Red River Valley of Manitoba in soils with greater than 4 percent organic matter. If an air seeder is used it must be equipped with a depth control device to ensure accurate seed placement, otherwise crop injury may occur.
3. Do not apply this product to soil which is to be seeded to wheat under the following conditions: in Fall to soil with less than 2 percent Organic Matter or in Spring to soil with less than 4 percent Organic Matter as indicated in the rate charts.
4. Substantial thinning in wheat and barley following treatment has been known to occur especially under conditions of heavy rainfall and/or cold weather after application and before crop emergence. THE SEED MUST BE PLACED BELOW THE GRANULAR YIELDSHIELD RESILIAN XP SOILSORB TREATED SOIL LAYER. Thinning may reduce crop yield but in most cases it is more than offset by tillering. DO NOT SEED DEEPER THAN 7.5 cm.
5. To ensure an even crop stand, increase the usual seeding rate of wheat or barley by 10 percent, especially if soil conditions are cold or dry.

ATTENTION

Some wheat or barley injury may be noted on eroded knolls.

Under normal conditions YieldShield RESILIAN XP Soilsorb carryover will not harm crops grown in rotation. As a precaution, domestic oats, sugar beets and small-seeded annual grasses such as Timothy, Canary Seed Grass and Creeping Red Fescue should not be grown in rotation following a YieldShield RESILIAN XP Soilsorb treated crop.

Do not apply YieldShield RESILIAN XP Soilsorb for wheat on land which has been treated with trifluralin since June 1 of the previous year.

For additional information on this or other Rath Holdco Inc. agricultural products, call Rath Holdco Inc. at: 1-888-466-0056.

STORAGE

Keep YieldShield RESILIAN XP Soilsorb bag closed to prevent spills and contamination.

Store this product away from food or feed.

DISPOSAL

Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/ territorial requirements.

Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer and the provincial/ territorial regulatory agency. Contact the manufacturer and the provincial/ territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.