(Container)



Stellar[™] UNPACKED Herbicide

GROUP 2 4 HERBICIDES

For postemergent control or suppression of annual broadleaved weeds in spring wheat, durum wheat, spring barley, oats, grass grown for seed production, new grass pastures and established grass pastures.

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING PEACE RIVER REGION) ONLY

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS: florasulam 2.5 g/L fluroxypyr 100 g/L (present as 1-methylheptyl ester)

Contains 1,2-benzisothiazoline-3-one at 0.01% as a preservative Suspension

REGISTRATION NUMBER 34516 PEST CONTROL PRODUCTS ACT

WARNING EYE AND SKIN IRRITANT POTENTIAL SKIN SENSITIZER HARMFUL IF SWALLOWED

NET CONTENTS: 1 L - bulk

Corteva Agriscience Canada Company

2450, 215-2nd Street SW Calgary, Alberta T2P 1M4 1-800-667-3852

[™]Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS KEEP OUT OF REACH OF CHILDREN DO NOT APPLY USING AERIAL APPLICATION EQUIPMENT

Harmful if swallowed. Causes eye and skin irritation. Do not get in eyes, on skin or on clothing. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab. Wash contaminated clothing before reuse. Destroy contaminated shoes and leather articles.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12-hours.

DO NOT use in residential areas. Residential areas are defined as any use site where bystanders, including children, could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public, including children, could be exposed.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store near heat or open flame.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on the judgment of the physician in response to reactions of the patient. This product contains a **PETROLEUM DISTILLATE**. **DO NOT INDUCE VOMITING.** Vomiting may cause aspiration pneumonia. If burn is present, treat as any thermal burn after decontamination. If swallowed, aspiration may cause chemical pneumonia. When considering emptying the stomach, the danger of chemical pneumonia must be weighed against toxicity. If lavage is performed, a cuffed endotrachael tube should be considered

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This product contains an active ingredient and aromatic petroleum distillate which are toxic to aquatic organisms.

STORAGE

Store in original containers in a secure, dry heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. Store this product away from food or feed. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recycle Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

- 1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
- 2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Containers

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

(Booklet)



Stellar[™] UNPACKED Herbicide

GROUP 2 4 HERBICIDES

For postemergent control or suppression of annual broadleaved weeds in spring wheat, durum wheat, spring barley, oats, grass grown for seed production, new grass pastures and established grass pastures.

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING PEACE RIVER REGION) ONLY

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND BOOKLET BEFORE USING KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS: florasulam 2.5 g/L fluroxypyr 100 g/L (present as 1-methylheptyl ester)

Contains 1,2-benzisothiazoline-3-one at 0.01% as a preservative Suspension

REGISTRATION NUMBER 34516 PEST CONTROL PRODUCTS ACT

WARNING EYE AND SKIN IRRITANT POTENTIAL SKIN SENSITIZER HARMFUL IF SWALLOWED

NET CONTENTS: 1 L - Bulk

Corteva Agriscience Canada Company

2450, 215-2nd Street SW Calgary, Alberta T2P 1M4 1-800-667-3852

[™]Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS KEEP OUT OF REACH OF CHILDREN DO NOT APPLY USING AERIAL APPLICATION EQUIPMENT

Harmful if swallowed. Causes eye and skin irritation. Do not get in eyes, on skin or on clothing. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab. Wash contaminated clothing before reuse. Destroy contaminated shoes and leather articles.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12-hours.

DO NOT use in residential areas. Residential areas are defined as any use site where bystanders, including children, could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public, including children, could be exposed.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store near heat or open flame.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on the judgment of the physician in response to reactions of the patient. This product contains a **PETROLEUM DISTILLATE. DO NOT INDUCE VOMITING.** Vomiting may cause aspiration pneumonia. If burn is present, treat as any thermal burn after decontamination. If swallowed, aspiration may cause chemical pneumonia. When considering emptying the stomach, the danger of chemical pneumonia must be weighed against toxicity. If lavage is performed, a cuffed endotrachael tube should be considered

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This product contains an active ingredient and aromatic petroleum distillate which are toxic to aquatic organisms.

STORAGE

Store in original containers in a secure, dry heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. Store this product away from food or feed. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

- 1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
- 2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Containers

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

GENERAL INFORMATION

Stellar[™] UNPACKED Herbicide is a selective postemergence herbicide for the control of hard-to-kill annual broadleaved weeds in spring wheat, durum wheat, spring barley, oats, grass grown for seed production, new grass pastures and established grass pastures. Stellar UNPACKED Herbicide is mixed with water and applied as a uniform broadcast spray.

Stellar UNPACKED Herbicide MUST be applied early postemergence, to the main flush of actively growing broadleaved weeds. Warm, moist growing conditions promote active weed growth and enhance the activity of Stellar UNPACKED Herbicide by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds. See "DIRECTIONS FOR USE" section of this label for complete use details.

Stellar UNPACKED Herbicide stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 2 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

MODE OF ACTION

Stellar UNPACKED Herbicide contains a Group 2 and a 4 mode of action herbicide. The Group 2 mode of action herbicide inhibits the production of the ALS enzyme in plants. This enzyme is essential for the production of certain amino acids required for plant growth. The Group 4 mode of action herbicide disrupts normal plant growth regulation resulting in death of susceptible plants.

GENERAL USE PRECAUTIONS

Sensitive Plants

Do not apply Stellar UNPACKED Herbicide directly to, or otherwise permit it to come in direct contact with susceptible crops or desirable plants including alfalfa, edible beans, canola, flowers and ornamentals, lentils, lettuce, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes or tobacco.

Non-Target Sites

Do not apply where proximity of susceptible crops or other desirable plants is likely to result in exposure to spray or spray drift. See Environmental Precautions section of the label.

Crop Rotation

Fields previously treated with Stellar UNPACKED Herbicide can be seeded the following year to barley, canola, corn, dry common beans (*Phaseolus vulgaris*), flax, lentils, mustard (brown, oriental, and/or yellow), oats, peas, potatoes (except see potatoes), soybeans, sunflowers or wheat or fields can be summerfallowed.

This product has potential to leach. Do not apply excessive irrigation. Do not use in successive years at the same site.

Tank Mixtures

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Corteva Agriscience Canada Company at 1-800-667-3852 or <u>www.corteva.ca</u> for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

Spray Equipment Precaution

Do not apply through any type of irrigation system.

To Reduce Spray Drift:

- 1. Use nozzles delivering higher volumes and coarser droplets.
- 2. Use low pressures (200 to 275 kPa).
- 3. Use 100 L/ha of spray solution.
- 4. Spray when the wind velocity is 15 km/hr or less.
- 5. Spot treatments should only be applied with a calibrated boom to prevent over-application.

Sprayer clean-out

To avoid injury to desirable plants, thoroughly clean equipment used to apply this product before re-use or using it to apply other chemicals.

- 1. Immediately after spraying, completely drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
- 2. First rinse:
 - Spray the inside of tank with clean water and fill the sprayer with at least one tenth of the spray tank volume.
 - Agitate and circulate for 15 minutes, and flush through booms and hoses.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.
- 3. Second rinse:

- Fill the tank with clean water.
- Add All Clear Spray Tank Decontaminator, or Clean-Out Spray Tank Cleaner, or 1 L of household ammonia (containing a minimum of 3 % ammonia) per 100 L of water, or similar tank cleaning agent as per manufacturer's recommendations while filling the tank with clean water.
- Agitate and then flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full. Allow to stand for 15 minutes with agitation. Flush the solution out of the spray tank through the spray booms. Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
- If possible, let the solution stand in the sprayer tank and booms for an extended period of time, overnight if possible.
- After flushing the boom and hoses, drain tank completely.
- Remove nozzles and screens and clean separately with a cleaning agent or an ammonia solution (100 mL in 10 L water).
- 4. Third rinse:
 - Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.

Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour which may cause eye, nose, throat, and lung irritation. Do not clean equipment in an enclosed area.

DIRECTIONS FOR USE

READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE. DO NOT APPLY TO CROPS UNDERSEEDED WITH LEGUMES.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Crops

Spring wheat, durum wheat, spring barley and oats

Application Rate

For control of a wide spectrum of broadleaved weeds, apply a uniform spray containing 1.0 L/ha of Stellar UNPACKED Herbicide in a minimum of 100 L of water per hectare to thoroughly cover the weeds. Apply to actively growing wheat, barley and oats from the 2 leaf expanded to 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds.

Mixing Instructions

- 1. Fill sprayer tank 1/2 full of water
- 2. Start sprayer tank agitation
- 3. Add the required amount of Stellar UNPACKED Herbicide
- 4. Complete filling the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare
- 5. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes)
- 6. Follow sprayer clean-up directions

Note: Do not add a surfactant to this product unless instructed to do so (see tank mix combinations – Table 3).

Table 1: Weeds Controlled by Stellar UNPACKED Herbicide at 1.0 L/ha

cleavers	volunteer flax
common chickweed	wild buckwheat
kochia•	

including ALS resistant biotypes

Grass Grown for Seed Production (Timothy, brome, tall and creeping red fescue) or in New Grass Pastures

To control wild mustard, shepherd's purse and those weeds labelled for control with 1.0 L/ha, apply Stellar UNPACKED Herbicide in the spring or fall, after seeding, to actively growing weeds at the rate of 1 L/ha.

Established Grass Pastures

To control cleavers, creeping buttercup, curled dock, sheep sorrel, broadleaved dock and dandelion (season long control) in established grass pastures, apply Stellar UNPACKED Herbicide in the spring to actively growing weeds at the rate of 1.6 L/ha.

Pre-Harvest/Grazing Intervals

- 1. Do not cut the treated crop for hay or graze treated crop within 7 days after application.
- 2. Do not harvest the treated crop within 60 days after application.

TANK-MIX COMBINATIONS WITH STELLAR UNPACKED HERBICIDE

Crops Registered

spring wheat, durum wheat and spring barley (see Table 3 for exclusions)

Tank mixtures of Stellar UNPACKED Herbicide with other herbicides will provide control of additional broadleaved weeds and specified annual grasses. Apply when crops and weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds. Follow all precautions, minimum interval to harvest and directions for use on the Stellar UNPACKED Herbicide and tank-mix partner labels.

Mixing Instructions

- 1. Fill sprayer tank 1/2 full of water.
- 2. Start sprayer tank agitation.
- 3. Add the required amount of Stellar UNPACKED Herbicide and continue to agitate.
- 4. Add the required amount of tank-mix partner.
- 5. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.

Note: Add only the adjuvant recommended below. Follow tank-mix partner label for order of mixing.

Herbicide Tank Mix Partner	Crops Registered	Rate (Product/ha)	Additional Weeds Controlled or Suppressed (S) Include
2,4-D 700 spring wheat, Ester [†] durum wheat &		424 mL/ha (280 g ae/ha)	lamb's-quarters, volunteer canola•
	spring barley	530-636 mL/ha (350-420 g ae/ha)	flixweed, redroot pigweed, ragweed, (S), shepherd's-purse, stinkweed, stork's-bill, sunflower (annual), wild mustard
MCPA Ester 600 [†]	spring wheat, durum wheat &	467 mL/ha (280 g ae/ha)	lamb's-quarters, volunteer canola•
	spring barley -	580-700 mL/ha (350-420 g ae/ha)	flixweed, hempnettle, ragweed, redroot pigweed, shepherd's-purse, smartweed, stork's-bill (S), sunflower (annual), wild mustard
Express Toss- N-Go Dry Flowable 75% Herbicide	spring wheat, durum wheat & spring barley	10 g/ha	lamb's-quarters, redroot pigweed (S), Russian thistle (S)

Table 2: Stellar UNPACKED Herbicide Tank Mix Combinations for Broadleaf Weed Con	trol
Table 2. Otenai Otti Aoneb Herbiciae Tank ink oombinations for Broadical Weed oon	

[†] Use higher rates for weeds in the bud stage, dry or cold weather or heavy infestations

*Including imidazolinone tolerant volunteer canola

Table 3: Stellar UNPACKED Broadleaf Tank Mix Combinations plus Other Herbicides for Annual
Grass Control

Herbicide Tank Mix Partner	Crops Registered	Rate Product/h a	Adjuvant and Rate	Additional Weeds Controlled or Suppressed (S)
Assert 300 SC Herbicide*	spring wheat, durum wheat & spring barley	1.6 L	Refer to the Assert 300 SC Herbicide label	wild oat
Axial 100 EC Herbicide*	spring wheat & spring barley	600 mL/ha	Adigor 700 mL/ha	wild oats & green foxtail
Everest 2.0 Herbicide***††‡	spring wheat & durum wheat	36-72 mL/ha	Non-ionic surfactant (see label) 0.25% v/v	wild oats & green foxtail

^{††}For conditions for when to use the higher rates, refer to the Everest 2.0 Herbicide label.

[‡]Warning: control of green foxtail may be reduced with the tank mix of Stellar UNPACKED Herbicide + Express Toss-N-Go Dry Flowable 75% Herbicide.

* Do not apply Axial 100 EC Herbicide in tank mixes containing 2,4-D.

** Wheat (including durum) exposed to water-logged or saturated soils, or temperature extremes such as heat or freezing weather, or drought, low fertility or plant disease at application time could show unacceptable injury symptoms. Weed control may also be reduced by these same conditions.

* Do not apply Everest 2.0 Herbicide or Assert 300 SC Herbicide in tank mixes containing Express Toss-N-Go Dry Flowable 75% Herbicide.

SPRAY BUFFER ZONES

Spot treatments using hand-held equipment **DO NOT** require a spray buffer zone.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Сгор	Spray Buffer Zones (metres) Required for the Protection of:				
		Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Spring wheat, durum wheat, winter wheat, spring barley, oats, seedling and established forage grasses for seed, new and established grass pastures	1	0	1	1	3

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product that are larger than 1m can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, Stellar UNPACKED Herbicide is a Group 2 and Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Stellar UNPACKED Herbicide and other Group 2 and Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Stellar UNPACKED Herbicide or other Group 2 and/or Group 4 herbicide within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.

- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Corteva Agriscience Canada Company at 1-800-667-3852.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

[™]Trademark of Corteva Agriscience and its affiliated companies.