

2023-0109
2023-09-05



{Base Container Label}

GROUP	7	FUNGICIDE
-------	---	-----------

Zeltera Fungicide
Suspension
COMMERCIAL



WARNING – POISON

ACTIVE INGREDIENT: Inpyrfluxam.....381 g/L

Contains 1,2-benzisothiazolin-3-one at 0.046 % OR 5-Chloro-2-methyl-4-isothiazolin-3-one at 0.0017% and 2-Methyl-4-isothiazolin-3-one at 0.0006% as a preservative.

REGISTRATION NUMBER 33820 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 80 mL to bulk

Valent Canada, Inc.
201-230 Halon Creek Blvd.
Guelph, ON N1C 0A1
Contact Phone Number: (519) 767-9262

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

PRECAUTIONS:

KEEP OUT OF THE REACH OF CHILDREN. KEEP TREATED SEED OUT OF REACH OF CHILDREN AND ANIMALS. Harmful or fatal if swallowed.

[OPTIONAL STATEMENT FOR SMALLER CONTAINER SIZE]

[Follow the personal protective equipment, engineering control, and restriction requirements for the appropriate mixer/loader and applicator scenario as described on the booklet label.]

Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Apply only in a way that this product will not contact workers or other persons, either directly or through drift. Only workers wearing the personal protective equipment may be in the area where treating, bagging, sewing and/or stacking is occurring.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Follow the personal protective equipment and restriction requirements for the appropriate mixer/loader and applicator scenario:

Seed Type	Tasks	PPE/Engineering Controls
For Commercial Seed Treatment (including facility workers and mobile treaters)		
All labelled seeds	Treating (including mixing, loading, treating, calibrating, bagging, sewing and stacking)	Closed transfer only. Closed transfer includes closed mixing, loading, calibrating and closed treatment. Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.
	Cleaning and repairing	Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.
For On-Farm Seed Treatment		
<ul style="list-style-type: none"> Legume vegetable seeds of crop group 6, including soybean seeds 	Treating (including mixing, loading, treating and calibrating)	Closed or open transfer is permitted. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.

• Labelled cereal seeds, except corn seeds	Cleaning and repairing	Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.
For Planting and Handling Treated Seed		
All label seeds	Planting	Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Use an open-cab or closed-cab tractor when planting. Gloves are not required within a closed cab.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., contact Valent Canada, Inc.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote is available. Treat the patient symptomatically.

ENVIRONMENTAL PRECAUTIONS:

[OPTIONAL STATEMENT FOR SMALLER CONTAINER SIZE]

[See booklet for full user precautions and restrictions.]

Toxic to aquatic organisms, birds, small wild mammals and non-target terrestrial plants. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE:

Keep pesticide in its original container. Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures. Store this product away from food or feed.

DISPOSAL:**RECYCLABLE CONTAINER:**

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer and the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.



{Booklet Label}

GROUP	7	FUNGICIDE
-------	---	-----------

Zeltera Fungicide

Suspension
COMMERCIAL

A FUNGICIDE SEED TREATMENT PRODUCT PROVIDING SYSTEMIC PROTECTION AGAINST SEED AND SEEDLING DISEASES CAUSED BY *RHIZOCTONIA SOLANI* AND DISEASES CAUSED BY OTHER FUNGAL PATHOGENS

WARNING – POISON



An INDIFLIN® brand

ACTIVE INGREDIENT: Inpyrfluxam.....381 g/L

Contains 1,2-benzisothiazolin-3-one at 0.046 % OR 5-Chloro-2-methyl-4-isothiazolin-3-one at 0.0017% and 2-Methyl-4-isothiazolin-3-one at 0.0006% as a preservative.

REGISTRATION NUMBER 33820 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

For all labelled seeds, commercial seed treatment (facilities and mobile treaters with closed transfer, including closed mixing, loading, calibrating, and closed treatment equipment only) is permitted.

For seeds of legume vegetables of crop group 6 (including soybeans), barley, buckwheat, pearl and proso millet, oats, rye, teosinte, triticale and wheat, on-farm treatment (open transfer, including open mixing, loading, calibrating, and open treatment equipment) is permitted.

For sugar beet seeds, commercial or on-farm treatment is NOT PERMITTED in Canada, but treated seeds can be imported for planting.

This product contains no colourant. An appropriate colourant must be added when the product is applied to seed, according to regulations pertaining to the “Seeds Act”. A baby blue colourant must be added when this product is applied to canola/rapeseed. A red colourant must be added when product is applied to grain seed.

NET CONTENTS: 80 mL to bulk

Valent Canada, Inc.
201-230 Halon Creek Blvd.
Guelph, ON N1C 0A1
Contact Phone Number: (519) 767-9262

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

PRECAUTIONS:

KEEP OUT OF THE REACH OF CHILDREN. KEEP TREATED SEED OUT OF REACH OF CHILDREN AND ANIMALS. Harmful or fatal if swallowed.

Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Apply only in a way that this product will not contact workers or other persons, either directly or through drift. Only workers wearing the personal protective equipment may be in the area where treating, bagging, sewing and/or stacking is occurring.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Follow the personal protective equipment and restriction requirements for the appropriate mixer/loader and applicator scenario:

Seed Type	Tasks	PPE/Engineering Controls
For Commercial Seed Treatment (including facility workers and mobile treaters)		
All labelled seeds	Treating (including mixing, loading, treating, calibrating, bagging, sewing and stacking)	Closed transfer only. Closed transfer includes closed mixing, loading, calibrating and closed treatment. Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.
	Cleaning and repairing	Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.
For On-Farm Seed Treatment		
<ul style="list-style-type: none"> • Legume vegetable seeds of crop group 6, including soybean seeds • Labelled cereal seeds, 	Treating (including mixing, loading, treating and calibrating)	Closed or open transfer is permitted. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.
	Cleaning and repairing	Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.

except corn seeds		
For Planting and Handling Treated Seed		
All label seeds	Planting	Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Use an open-cab or closed-cab tractor when planting. Gloves are not required within a closed cab.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., contact Valent Canada, Inc.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote is available. Treat the patient symptomatically.

ENVIRONMENTAL PRECAUTIONS:

Toxic to aquatic organisms, birds, small wild mammals and non-target terrestrial plants. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

MODE OF ACTION:

The active ingredient in Zeltera Fungicide, inpyrfluxam, belongs to FRAC Group 7, the succinate dehydrogenase inhibitor (SDHI) group of fungicides. As with other Group 7 fungicides, inpyrfluxam acts by inhibiting succinate dehydrogenase, a key enzyme in the fungal respiration chain.

SEED QUALITY AND STORAGE OF TREATED SEED:

Seed treatments applied to low vigour or poor quality seed or to mechanically damaged seed may result in loss of seed germination or reduced seedling vigour. To verify seed safety, treat a small portion of seed with the desired slurry mixture (Zeltera Fungicide alone or in combination with other commercial seed treatment products) prior to treatment of an entire seed lot. Utilize standard seed germination methodologies to determine and verify seed safety. Use proper storage conditions for treated seed. Due to seed quality and seed storage conditions beyond the control of Valent Canada, Inc., no claims are made to guarantee the germination of carry-over treated seed.

USE RESTRICTIONS:

- Do not use treated seed for food, feed, or oil processing.
- Do not graze or feed forage and hay of legume vegetables (including pea vines) to livestock.
- Due to seed quality conditions beyond the control of Valent Canada, Inc., no claims are made to guarantee germination of carry-over seed.

LABELLING OF TREATED SEED:

All containers or packages containing treated seed for sale or use in Canada must be labeled or tagged as follows:

- This seed has been treated with Zeltera Fungicide containing inpyrfluxam.
- Do not use for feed, food or oil processing.
- Excess seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practices.
- Store away from feeds and other foodstuffs.
- Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes when handling and planting treated seeds. Gloves are not required within a closed-cab tractor.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Toxic to birds and small wild mammals. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.
- Do not graze or feed forage and hay of legume vegetables (including pea vines) to livestock.
- KEEP TREATED SEED OUT OF REACH OF CHILDREN AND ANIMALS.
- DO NOT plant by hand.

USE INFORMATION:

Zeltera Fungicide is a systemic fungicide seed protection product for use on a wide range of crops. The active ingredient, inpyrfluxam, provides protection against *Rhizoctonia solani* causing seed decay/pre-emergence damping-off, post-emergence damping-off, seedling blight and root

rot on listed crops. Inpyrfluxam also controls or suppresses certain other fungal diseases as listed on this label.

For best results, use Zeltera Fungicide combined with Oomycete-active seed treatment fungicides, to broaden the spectrum of activity against seed and seedling pathogens that cause seed decay and seedling dieback.

DIRECTIONS FOR USE:

Zeltera Fungicide must be used only in accordance with use instructions on this label utilizing standard mechanical or slurry seed treating equipment that is calibrated to accurately and uniformly apply the product to seed. This product is for both commercial and on-farm application. Do not apply this product in a hopper-box or planter-box at planting time. Always mix product thoroughly before use.

Tank Mixing

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Valent Canada, Inc. at 519-767-9262 for information before applying any tank mix that is not specifically recommended on this label.

Pre-test tank mixtures with other seed treatment products to evaluate formulation compatibility and to ensure proper physical compatibility of products.

Accurate application of Zeltera Fungicide to seed is required for best performance. Thoroughly mix Zeltera Fungicide in the desired seed slurry mixture, prior to application to seed.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

The following table provides use rates of Zeltera Fungicide by labeled crop, with disease protection guidelines. Where a rate range is indicated, use up to the maximum rate in fields with a history of higher disease pressure or when conditions are predicted to be favourable for disease development (e.g., cool, wet weather and field conditions).

When applied as directed, Zeltera Fungicide can be expected to control the listed diseases, unless otherwise indicated as suppression.

Cereal grain crops: specifically barley, buckwheat, pearl millet, proso millet, oats, rye, teosinte, triticale, wheat	
Diseases Controlled	Application Rate of Zeltera Fungicide (mL/100 kg seed)

Barley: True Loose Smut (<i>Ustilago nuda</i>)	2.6 – 5.2 (1 to 2 g a.i./100 kg seed)
Wheat: Wheat Loose Smut (<i>Ustilago tritici</i>)	
Seed decay/pre-emergence damping-off, post - emergence damping-off and seedling blight (<i>Rhizoctonia solani</i>) Suppression of root rot (<i>R. solani</i>)	2.6 – 5.2 (1 to 2 g a.i./100 kg seed)
Corn (sweet and field); popcorn	
Diseases Controlled	Application Rate of Zeltera Fungicide (mL/100 kg seed)
Seed decay/pre-emergence damping-off, post- emergence damping-off and seedling blight (<i>Rhizoctonia solani</i>)	13 (5 g a.i./100 kg seed) The volume required to treat 80,000 seeds can be calculated as follows based on seed weight (# seeds/kg): # mL/80,000 seeds = 80,000 / (# seeds/100 kg) x 13 mL Field Corn: To deliver 5 g ai/100 kg seed, apply 3 mL/80,000 seed (equals 0.0141 mg ai/seed) based on 3524 seed/kg (1600 seed/lb) count. The volume required to treat 100,000 seeds can be calculated as follows based on seed weight (# seeds/kg): # mL/100,000 seeds = 100,000 / (# seeds/100 kg) x 13 mL Sweet Corn: To deliver 5 g ai/100 kg seed, apply 2 mL/100,000 seeds (equals 0.0072 mg ai/seed) based on 6938 seed/kg (3150 seed/lb) count Popcorn: To deliver 5 g ai/100 kg seed, apply 2 mL/100,000 seeds (equals 0.0074 mg ai/seed) based on 6742 seed/kg (3061 seed/lb) count

**Crop Group 6 - Legume Vegetables, Succulent or Dried
(except Soybean)**

Bean (*Lupinus*) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (*Phaseolus*) (includes field pea, cowpea, field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); bean (*Vigna*) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); broad bean (fava); chickpea (garbanzo); guar; jackbean; lablab bean; lentil; pea (*Pisum*) (includes dwarf pea, edible-podded pea, English pea, garden pea, green pea, snowpea, sugar snap pea); pigeon pea; sword bean

Diseases Controlled	Application Rate of Zeltera Fungicide (mL /100 kg seed)
Seed decay/pre-emergence damping-off, post-emergence damping-off, and seedling blight (<i>Rhizoctonia solani</i>), Suppression of root rot (<i>R. solani</i>)	6.5 – 13 (2.5 to 5 g a.i./100 kg seed)
Soybean	
Diseases Controlled	Application Rate of Zeltera Fungicide (mL / 100 kg seed)
Seed decay/pre-emergence damping-off, post-emergence damping-off and seedling blight (<i>Rhizoctonia solani</i>)	6.5 – 13 (2.5 to 5 g a.i./100 kg seed)* *The volume of product required to treat 140,000 seeds can be calculated with the following formula: mL per 140,000 seeds = labeled rate (mL/100kg seed) X [140,000 seeds / seed weight (seed/kg)] ÷ 100
Sudden Death Syndrome (<i>Fusarium virguliforme</i>)	208 (80 g a.i./100 kg seed)
In soybean, it is acceptable to apply foliar applications of Excalia Fungicide, also containing inpyrfluxam, following the use of Zeltera Fungicide. Regardless of application method, do not apply more than 180 g a.i./ha of inpyrfluxam per year.	

Rapeseed (including Canola)

Diseases Controlled	Application Rate of Zeltera Fungicide (mL/100 kg seed)
Seed decay/pre-emergence damping-off, post-emergence damping-off, seedling blight and root rot (<i>Rhizoctonia solani</i>)	13 (5 g a.i./100 kg seed)

Blackleg (<i>Leptosphaeria maculans</i>)	13-26 (5-10 g a.i./100 kg seed)
Within the rate range, use the lowest rate of Zeltera Fungicide for blackleg suppression; use the highest rate for blackleg control in fields where there is a history of disease pressure or when conditions are conducive for disease development.	

Imported Treated Seed Information

NOT FOR COMMERCIAL or ON-FARM SEED TREATMENT OF SUGAR BEET SEED IN CANADA

The following treated seed can be imported into Canada for planting:

Sugar beet	
Disease Protection Claims	Application Rate of Zeltera Fungicide (mL/100,000 seeds)
Seed decay/pre-emergence damping-off and post-emergence damping off (<i>Rhizoctonia solani</i>)	<p style="text-align: center;">0.13-0.26 (0.05 to 0.1 g a.i./100,000 seeds)*</p> <p>* The volume of product required to treat 100 kg of raw seeds can be calculated with the following formula: mL per 100 kg raw seeds = labeled rate (g a.i./100,000 seeds) X seed weight (seed/kg raw seeds) ÷381.4 g a.i./L</p> <p>A maximum of one foliar application of Excalia Fungicide, also containing inpyrfluxam, may be made following use of Zeltera Fungicide as a seed treatment.</p>

RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, please note that Zeltera Fungicide is a Group 7 fungicide. Any fungal population may contain individuals naturally resistant to Zeltera Fungicide and other Group 7 fungicides. A gradual or total loss of pest control may occur over time if this fungicide is used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of Zeltera Fungicide or other Group 7 fungicides with different chemical groups that control the same pathogens.

- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Monitor treated fungal populations for resistance development. Notify Valent Canada, Inc. if reduced sensitivity of the pathogen to Zeltera Fungicide is suspected.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance contact Valent Canada, Inc. at 1-800-682-5368.

STORAGE:

Keep pesticide in its original container. Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures. Store this product away from food or feed.

DISPOSAL:

RECYCLABLE CONTAINER:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

RETURNABLE CONTAINER:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

REFILLABLE CONTAINER:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

NON-RECYCLABLE/NON-RETURNABLE/NON-REFILLABLE CONTAINER:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer and the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.