

Brigade

Bird Repellent Granular

Solid

COMMERCIAL/AGRICULTURAL

ACTIVE INGREDIENTS:

Methyl Anthranilate 14.75% (When packed)

Crushed garlic 14.0%

REGISTRATION NO. 34098 PEST CONTROL PRODUCTS ACT

This product has been formulated from Food Grade Ingredients that are on the
“AAFCO” ‘Generally Regarded as Safe’ (GRAS) list

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION- SKIN AND EYE IRRITANT

POTENTIAL SKIN SENSITIZER

NET CONTENTS: 300 – 2,500 Grams

(6x50 – 50x50)

Norac Concepts Inc.
P.O. Box 31097
Guelph, ON, N1H 8K1
519-821-3633

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Do not swallow. May irritate the skin and eyes. Avoid contact with skin, eyes, or clothing. Avoid breathing dusts. Potential skin sensitizer. Wash thoroughly with soap and water after handling. Wear a long-sleeved shirt and long pants, chemical-resistant gloves, socks and shoes during handling.

For terrestrial uses; do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Individuals who are sensitive or allergic to garlic should avoid handling Brigade Bird Repellent Granular.

Do not use Brigade Bird Repellent Granular to repel birds in residential backyards. When used in parks, Brigade Bird Repellent Granular must not be placed in recreational areas of parks. Brigade Bird Repellent Granular must be placed out of reach of public.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air immediately. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

There is no specific antidote for this product. Apply symptomatic therapy.

STORAGE:

Store in original container, inaccessible to children, pets and domestic animals. Store apart from pesticides, fertilizers, food or feed that may cause cross-contamination from odour. Keep package closed when not in use. Store between 5 - 32 Celsius.

DISPOSAL:

1. Make the empty container unsuitable for further use.
2. Dispose of the container in accordance with provincial requirements.

3. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the provincial regulatory agency and the manufacturer.

GENERAL INFORMATION:

This product can be used in an integrated pest management program to repel birds. Brigade Bird Repellent Granular repels birds from areas where they are considered a nuisance or a hazard to property, structures, equipment, trees and plants. Brigade Bird Repellent Granular encourages birds to nest in new areas and away from sites where birds are unwanted. Do not apply directly to water, such as lakes, ponds or streams. Use Brigade Repellent Granular as part of an integrated pest management program to repel bird species causing economic and aesthetic damage only. Do not use to repel bird species considered potential vectors for human diseases. Do not use in areas where threatened or endangered birds are present. If eggs or young chicks are present, wait until the eggs hatch and the chicks leave the nest. Do not place Brigade Bird Repellent Granular on areas that may stain such as window sills and vinyl or in water bodies or areas where water has pooled.

Birds that can be repelled are Robins, Blue Jays, Cardinals, Sparrows, Grackles, European Starlings, and Blackbirds.

DIRECTIONS FOR USE:

APPLICATION DIRECTIONS: Each Granular pouch has an effective coverage area of approximately 9.3 square metres. (100 sq. ft). For best results, place or hang Granular pouches 2.4 – 3.0 metres (8 - 10 feet) apart. For increased effectiveness, place Granular pouches 1.8 metres (6 feet) or less apart.

Commercial Use Sites:

Use Brigade Bird Repellent Granular to repel birds that gather in large numbers and can damage industrial or commercial equipment and property, buildings, structures, trees and plants. Remove bird attractants such as nests and food sources before use.

Use Brigade Bird Repellent Granular to repel birds by hanging or placing Granular pouches in areas where birds congregate and can cause damage to property or equipment such as rafters, warehouses, walkway covers, industrial equipment, air conditioning and HVAC equipment.

Agricultural Use Sites:

Use Brigade Bird Repellent Granular to repel birds in crops. This product is not applied directly to crops. Use Brigade Bird Repellent Granular to repel birds by hanging or placing Granular pouches in areas where birds congregate and can cause damage to crops.

Use Brigade Bird Repellent Granular in the following crops:

Crop Group 1 - Root and Tuber vegetables	arrowroot, Chinese artichoke, Jerusalem artichoke, garden beet, sugar beet, edible burdock, edible canna, carrot, celeriac (celery root), turnip-rooted chervil, chicory, chufa, dasheen (taro), ginseng, horseradish, turnip-rooted parsley, parsnip, potato, radish, oriental radish (daikon), rutabaga, salsify, black salsify, Spanish salsify, skirret, sweet potato, turnip and true yam.
Crop Group 11-09 - Pome Fruits	apple, azarole, crabapple, mayhaw, medlar, pear, Asian pear, quince, Chinese quince, Japanese quince, tejocote, as well as cultivars, varieties and/or hybrids of these commodities.
Crop Group 12-09 -	apricot, Japanese apricot, plum, American plum, beach plum, Canada plum, cherry plum,

Stone Fruits	chicksaw plum, damson plum, Japanese plum, klamath plum, prune plum, plumcot, sloe, as well as cultivars, varieties and hybrids of these commodities.
Crop Group 13-07 - Berries and Small Fruits	Amur river grape, aronia berry, bayberry, bearberry, bilberry, blackberry (includes Andean blackberry, arctic blackberry, bingleberry, black satin berry, boysenberry, brombeere, California blackberry, Chesterberry, Cherokee blackberry, Cheyenne blackberry, common blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, evergreen blackberry, Himalayaberry, hullberry, lavacaberry, loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, mora, mures deronce, nectarberry, Northern dewberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, Southern dewberry, tayberry, youngberry, zarzamora, as well as cultivars, varieties and/or hybrids of these), highbush blueberry, lowbush blueberry, buffalo currant, buffaloberry, Chilean guava, chokecherry, cloudberry, cranberry, black currant, red currant, elderberry, European barberry, gooseberry, grape, highbush cranberry, edible honeysuckle, huckleberry, jostaberry, Juneberry (Saskatoon berry), fuzzy kiwifruit, hardy kiwifruit, lingonberry, maypop, mountain pepper berries, mulberry, muntries, native currant, partridgeberry, pincherry, raspberry (red and black), salal, schisandra berry, sea buckthorn, serviceberry, strawberry, wild raspberry, as well as cultivars, varieties and/or hybrids of these commodities.
Crop Group 15 - Cereal Grains	barley, buckwheat, corn, pearl millet, proso millet, oats, popcorn, rice, rye, sorghum (milo), teosinte, triticale, wheat and wild rice.

- For fruit-bearing trees and berries, have Granular pouches in place just as fruit or berries are starting to ripen.
- For all crops listed, have Granular pouches in place before bird damage begins to occur.
- Remove bird attractants such as nests and non-crop food sources before use.

DURATION: Granular pouches will last up to 3 months depending on environmental factors such as rain, sun, humidity and temperature. Granular in the pouches will break down and lose their form in time. As the granules degenerate, more active ingredient will be released. Reapply Granular pouches if they reduce in size by a quarter to a third of the original size to help maintain bird repellency.

EMERGENCY TELEPHONE NUMBER: All hours, (613) 787-5620 ONLY for health and environmental information.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offense under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.