

2022-4758
2022-10-06

GROUP	7	FUNGICIDE
-------	---	-----------

CABIL FUNGICIDE

WETTABLE GRANULES

For use in alfalfa grown for seed production, berries group, bulb vegetables group, carrots, cucurbit vegetables, field lettuce (head and leaf), fruiting vegetables, grapes, potatoes, mustard (oilseed and condiment variety only), stone fruits group, strawberries, succulent beans, succulent peas and sunflowers.

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: Boscalid ... 70%

REGISTRATION NO. 34638 PEST CONTROL PRODUCTS ACT

WARNING - EYE IRRITANT

READ THE LABEL AND ATTACHED BROCHURE BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 0.1 kg – 10 kg

Sharda Cropchem Limited
2nd Floor, Prime Business Park
Dashrathlal Joshi Road
Vile Parle (West)
Mumbai - 400056, India

Canadian Agent:
Sharda Cropchem Limited
63 Kingsview Blvd
Etobicoke, Ontario, CA
M9R 1V1
1-844-810-5720
1-416-840-5639

GENERAL INFORMATION

This package contains **CABIL FUNGICIDE**, a 70% water dispersible granular (WDG). The active ingredient in **CABIL FUNGICIDE** belongs to the anilid group of fungicides and has a unique mode of action.

CABIL FUNGICIDE has a protective effect because it inhibits spore germination, mycelial growth and sporulation of the fungus on the leaf surface. Optimum disease control is achieved when **CABIL FUNGICIDE** is applied in a regularly scheduled protective spray program and is used in a rotation program with other fungicides.

DIRECTIONS FOR USE (See specific sections for each crop group)

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Apply recommended rates of **CABIL FUNGICIDE** as instructed in the following series of crop application rate tables. Apply **CABIL FUNGICIDE** with ground or aerial spray equipment as specified. Equipment should be checked frequently for proper calibration.

Tank mixing: In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Sharda Cropchem Limited at 1-844-810-5720 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

Alfalfa Grown For Seed Production Application Rate Table

Ground, Aerial and Pivot or Sprinkler Irrigation (Chemigation)

Crop	Diseases Controlled	Application Rate (g/ha)
Alfalfa grown for seed production	Blossom blight (<i>Sclerotinia sclerotiorum</i> / <i>Botrytis cinerea</i>) Common leaf spot (<i>Pseudopeziza medicaginis</i>) Spring black stem (<i>Phoma medicagnis</i>) Leaf spot (<i>Leptosphaerulina briosiani</i>)	420
DO NOT GRAZE OR FEED TREATED HAY TO LIVESTOCK. Apply CABIL FUNGICIDE at 420 g/ha at 20-50% flowering to control blossom blight (<i>sclerotinia</i> , <i>botrytis</i>), common leaf spot and spring black stem. Apply every 7- to 14-days if disease persists, or weather conditions are favourable for disease development. For ground application, use a minimum spray volume of 100 L/ha. For air application, use a minimum spray volume of 40 L/ha. Apply a maximum of 3 applications per year.		

Succulent Beans Application Rate Table

Ground application only. DO NOT APPLY BY AIR. DO NOT APPLY BY PIVOT OR SPRINKLER IRRIGATION (Chemigation).

Crop	Diseases Controlled	Application Rate (g/ha)
Succulent beans <i>Phaseolus spp.</i> (includes runner bean, snap bean, wax bean, lima bean {green}, broad bean {succulent}) <i>Vigna spp.</i> (includes moth bean, yardlong bean, jackbean, asparagus bean, Chinese longbean, podded bean, blackeyed pea, southern pea)	Gray mold (<i>Botrytis cinerea</i>)	420
	White mold (<i>Sclerotinia sclerotiorum</i>)	560 - 770

Apply **CABIL FUNGICIDE** at 560-770 g/ha, at 20-50% flowering, to control white mold (*Sclerotinia sclerotiorum*), or at 420 g/ha to control gray mold (*Botrytis cinerea*). Apply a second time 7- to 14-days later if disease persists, or weather conditions are favourable for disease development.

Tank Mix with Matador® 120 EC or Warrior® Insecticide

For the control of corn borer, **CABIL FUNGICIDE** can be tank mixed with Matador 120 EC or Warrior Insecticide at 83 mL/ha. Apply before the larva bores into the plant stalk or pods. Follow the most restrictive application directions for each of the tank mix partners with respect to the maximum number of applications, pre-harvest interval and other label instructions.

Berries Group Application Rate and Timing Table

(Ground application only. DO NOT APPLY BY AIR. DO NOT APPLY BY PIVOT OR SPRINKLER IRRIGATION.)

Crop	Disease Controlled	Application Rate (g/ha)
Blackberry Raspberry (black and red) Currant Elderberry Blueberry (highbush and lowbush) Gooseberry Huckleberry Loganberry	Botrytis gray mold (<i>Botrytis cinerea</i>)	560

Begin applications prior to disease development at 560 g/ha of **CABIL FUNGICIDE** and continue on a 7- to 14-day schedule. Use the shorter interval when conditions favour the development of high disease pressure. Refer to Resistance Management instructions.

Bulb Vegetables Group Application Rate and Timing Table

(Ground application only. DO NOT APPLY BY AIR. DO NOT APPLY BY PIVOT OR SPRINKLER IRRIGATION.)

Crop	Diseases Controlled	Application Rate (g/ha)
Onions (dry bulb and green) Garlic Leek Shallot	Alternaria purple blotch (<i>Alternaria porri</i>) Botrytis leaf blight (<i>Botrytis squamosa</i>)	475
Begin applications prior to disease development at 475 g/ha of CABIL FUNGICIDE and continue on a 7- to 14-day schedule. Use the shorter interval when conditions favour the development of high disease pressure. Refer to Resistance Management instructions.		

Carrots Application Rate and Timing Table

(Ground application only. DO NOT APPLY BY AIR. DO NOT APPLY BY PIVOT OR SPRINKLER IRRIGATION.)

Crop	Disease Controlled	Application Rate (g/ha)
Carrots	Alternaria leaf blight (<i>Alternaria dauci</i>)	315
Begin applications prior to disease development at 315 g/ha of CABIL FUNGICIDE at 7- to 14-day intervals. Use the shorter interval when conditions favour the development of high disease pressure. Refer to Resistance Management instructions.		

Cucurbit Vegetables Group Application Rate and Timing Table

(Ground application only. DO NOT APPLY BY AIR. DO NOT APPLY BY PIVOT OR SPRINKLER IRRIGATION.)

Crop	Diseases Controlled	Application rate (g/ha)
Citron melon Cucumber Gherkin Pumpkin Watermelon <u>Edible gourd</u> Hyotan Cocuzza Chinese okra <u>Muskmelon</u> Cantaloupe Casaba Crenshaw melon Golden pershaw melon Honeydew melon Honey balls Mango melon Persian melon Pineapple melon Santa Claus melon Snake melon <u>Summer squash</u> Crookneck squash Scallop squash Straightneck squash Vegetable marrow Zucchini <u>Winter squash</u> Butternut squash Calabaza Hubbard squash Acorn squash Spaghetti squash	Alternaria blight (<i>Alternaria cucumerina</i>) Gummy stem blight (<i>Didymella bryoniae</i>)	460

Begin applications of **CABIL FUNGICIDE** prior to disease development and continue on a 7- to 14-day interval. Use the higher rate and the shorter interval when conditions favour the development of high disease pressure. Refer to Resistance Management instructions.

Fruiting Vegetables Group Application Rate and Timing Table

(Ground application only. DO NOT APPLY BY AIR. DO NOT APPLY BY PIVOT OR SPRINKLER IRRIGATION.)

Crop	Diseases Controlled	Application Rate (g/ha)
Eggplant Ground cherry Pepino	Early blight (<i>Alternaria solani</i>)	175 - 315
Field pepper (all varieties) Tomatillo Field tomato	Botrytis gray mold (<i>Botrytis cinerea</i>)	420

For control of early blight, begin applications of **CABIL FUNGICIDE** prior to disease development at 175-315 grams per hectare at 7- to 10-day intervals. Use the higher rate and shorter interval when conditions favour the development of high disease pressure. For control of botrytis gray mold, begin applications of **CABIL FUNGICIDE** prior to disease development at 420 g/ha followed by a second application after 7-10 days. **DO NOT** make more than 5 applications per year. Refer to Resistance Management instructions.

Grape Application Rate and Timing Table

(Ground application only. DO NOT APPLY BY AIR. DO NOT APPLY BY PIVOT OR SPRINKLER IRRIGATION.)

Crop	Disease Controlled	Application Rate (g/ha)
Grape	Powdery mildew (<i>Uncinula necator</i>)	315

For powdery mildew control, sprays may begin as of budbreak at the rate of 315 g/ha of **CABIL FUNGICIDE** on a 10- to 14-day schedule. Use the shorter interval when conditions favour the development of high disease pressure. Refer to Resistance Management instructions.

Field Lettuce (Head and Leaf) Application Rate and Timing Table

(Ground application only. DO NOT APPLY BY AIR. DO NOT APPLY BY PIVOT OR SPRINKLER IRRIGATION.)

Crop	Disease Controlled or Suppressed	Application Rate (g/ha)
Field lettuce (head and leaf)	Lettuce drop (<i>Sclerotinia minor</i>) - suppression	385
	Botrytis rot (<i>Botrytis cinerea</i>)	285

Apply **CABIL FUNGICIDE** at 385 g/ha for suppression of lettuce drop (*Sclerotinia minor*), and at 285 g/ha for control of botrytis rot (*Botrytis cinerea*). On direct seeded lettuce, make the first application immediately after thinning (within 2 days). If conditions continue to favour disease development, make a second application 10 to 20 days after the first application. On transplanted lettuce, make the first application 7-10 days after transplanting. If conditions continue to favour disease development, make a second application 10 to 20 days after the first application. Ensure thorough coverage of the lower portion of the plants and the surrounding soil surface.

Mustard Application Rate Table

Ground, Aerial and Pivot or Sprinkler Irrigation (Chemigation)

Crop	Diseases Controlled or Suppressed	Application Rate (g/ha)
Mustard (oilseed and condiment)	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>) Black spot (<i>Alternaria brassicae</i> and <i>raphani</i>) – Suppression or control	350

Apply **CABIL FUNGICIDE** at 350 g/ha at 20-50% flowering to control sclerotinia stem rot (*Sclerotinia sclerotiorum*) and suppress black spot (*Alternaria brassicae* and *raphani*). Apply a second time 7- to 14-days later up to full bloom if disease persists, or weather conditions are favourable for disease development. Apply **CABIL FUNGICIDE** at 350 g/ha at late flowering to early green pod to control black spot (*Alternaria brassicae* and *raphani*).

Succulent Pea Application Rate Table

Ground application only. DO NOT APPLY BY AIR. DO NOT APPLY BY PIVOT OR SPRINKLER IRRIGATION (Chemigation).

Crop	Diseases Controlled	Application Rate (g/ha)
Edible podded peas	Ascochyta blight (<i>Ascochyta</i> spp.) Mycosphaerella blight (<i>Mycosphaerella</i> spp.)	420
<u>Pea (<i>Pisum</i>)</u> Dwarf pea Edible-podded pea Snow pea Sugar snap pea Succulent shelled peas <u>Pea (<i>Pisum</i>)</u> English pea Garden pea Green pea Pigeon pea	White mold (<i>Sclerotinia sclerotiorum</i>)	560 - 770

For optimal disease control, begin applications of **CABIL FUNGICIDE** prior to disease development or at the beginning of flowering and repeat on a 5- to 14-day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when conditions favour the development of high disease pressure. Refer to Resistance Management instructions.

Potato Application Rate and Timing Table

(Ground, Aerial and Pivot or Sprinkler Irrigation)

Crop	Disease Controlled	Application Rate (g/ha)
Potato	Early blight (<i>Alternaria solani</i>)	175 - 315

For control of alternaria early blight, begin applications of **CABIL FUNGICIDE** prior to disease development at 175-315 g/ha and continue applications at 14 day intervals if conditions

continue to be favourable for disease development. Refer to the Resistance Management section for alternation instructions. Do not make more than 4 applications per year on potatoes.

Stone Fruit Group Application Rate and Timing Table

(Ground application only. DO NOT APPLY BY AIR. DO NOT APPLY BY PIVOT OR SPRINKLER IRRIGATION.)

Crop	Diseases Controlled	Application Rate (g/ha)
Apricots Cherry (sweet and tart) Nectarine Peach Plum Plumcot Prune	Brown rot (<i>Monilinia</i> spp.) Monilinia blossom blight (<i>Monilinia</i> spp.)	370

Begin applications at pink bud or prior to disease development. Apply **CABIL FUNGICIDE** at 370 g/ha and continue at 7- to 14-day intervals. Use the shorter interval when conditions favour the development of high disease pressure. Refer to Resistance Management instructions.

Strawberry Application Rate and Timing Table

(Ground application only. DO NOT APPLY BY AIR. DO NOT APPLY BY PIVOT OR SPRINKLER IRRIGATION.)

Crop	Disease Controlled	Application Rate (g/ha)
Strawberry	Botrytis gray mold (<i>Botrytis cinerea</i>)	560

Begin applications prior to disease development at 560 g/ha of **CABIL FUNGICIDE** and continue on a 7- to 14-day schedule. Use the shorter interval when conditions favour the development of high disease pressure. Refer to Resistance Management instructions.

Sunflower Application Rate Table

(Ground and Aerial)

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Sharda Cropchem Limited under the User Requested Minor Use Label Expansion program. For these uses, Sharda Cropchem Limited has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

Crop	Diseases Suppressed	Application Rate (g/ha)
Sunflowers	Sclerotinia head rot (<i>Sclerotinia sclerotiorum</i>) Leaf spot (<i>Alternaria helianthi</i>)	350 - 640

For optimal disease suppression, apply **CABIL FUNGICIDE** at early flower. Use the higher rate when conditions favour the development of high disease pressure or there is a history of high disease in the field. Apply a maximum of 1 application per year.
 For ground application, use a minimum spray volume of 100 L/ha.
 For aerial application, use a minimum spray volume of 50 L/ha.

Crop-Specific Restrictions and Limitations

Crop	Minimum Time from Application to Harvest (PHI)	Maximum Rate Per Hectare Per Application (kilograms)	Maximum Number of Applications Per Year	Maximum Rate Per Hectare Per Year (kilograms)
Alfalfa grown for seed production	N/A	0.420	3	1.26
Beans (succulent)	7 days	0.770	2	1.54
Berries group ¹	0 days	0.560	4	2.24
Bulb vegetables group ¹	7 days	0.475	6	2.87
Carrots	0 days	0.315	5	1.58
Cucurbit vegetables group ¹	0 days	0.460	4	1.84
Fruiting vegetable group ¹ (for early blight)	0 days	0.315	5	1.89
Fruiting vegetable group ¹ (for botrytis)	0 days	0.420	2 out of 5	1.89
Grapes (for powdery mildew)	14 days	0.315	5	1.58
Lettuce (lettuce drop)	14 days	0.385	2	0.77
Lettuce (botrytis rot)	14 days	0.285	2	0.57
Mustard (oilseed and condiment)	21 days	0.350	2	0.700
Peas - succulent (edible podded and shelled)	7 days	0.770	2	1.540
Potato (for early blight)	30 days	0.315	4	1.26
Stone fruits group ¹	0 days	0.370	5	1.85
Strawberries	0 days	0.560	5	2.80
Sunflowers	21 days	0.640	1	0.640

¹ For a complete list of the various crop groups, see DIRECTIONS FOR USE listed above.

Plant Back Interval: A plant back restriction of 14 days is required for all crops not on the label.

Ground application

Apply **CABIL FUNGICIDE** at rates listed in the Application Rate and Timing Tables (crop specific) when conditions are favourable for the development of disease. Use a minimum water volume of 100 L/ha and ensure thorough coverage of foliage. Consult nozzle manufacturer's recommendation for spray pressures for specific nozzles.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. DO NOT apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

Aerial application

Apply **CABIL FUNGICIDE** at rates listed in application rate and timing tables (crop specific) when conditions are favourable for the development of disease. Use a minimum water volume of 40 L/ha. Ensure thorough coverage of foliage. Consult nozzle manufacturer's recommendation for spray pressures for specific nozzles.

Aerial application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

Instructions for Aerial Applications

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label. Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment. Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted. It is desirable that the pilot have communication capabilities at each treatment site at the time of application. The field crew and the mixer/ loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-844-810-5720 or obtain technical advice from the distributor or your provincial agricultural representative.

Pivot or Sprinkler Irrigation

Sprayer Preparation: Chemical tank and injector system should be thoroughly cleaned. Flush System with clean water.

Application instructions: Apply **CABIL FUNGICIDE** at rates and timings described above.

Use Precautions for Sprinkler Irrigation Applications

- Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.
- Add this product to the pesticide supply tank containing sufficient water to maintain a continuous flow by the injection equipment. In continuous moving systems, inject this product-water mixture continuously, applying the labeled rate per acre for that crop. Do not exceed 0.64 cm (1/4) inch (63,500 litres) per hectare. In stationary or non-continuous moving systems, inject the product-water mixture in the last 15-30 minutes of each set allowing sufficient time for all of the required pesticide to be applied by all the sprinkler heads and applying the labeled rate per acre for that crop. Do not apply when wind speed favours drift beyond the area intended for treatment. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. Thorough coverage of foliage is required for good control. Good agitation should be maintained during the entire application period.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label- prescribed safety devices for public water systems are in place.

Specific Instructions for Public Water Systems

- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional, reduced- pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favours drift beyond the area intended for treatment.

Chemigation: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Applications MUST be conducted WITHOUT the use of end guns.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label,
- low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, rangelands, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of Application	Crop	Spray Buffer Zones (metres) Required for the Protection of:		
		Freshwater Habitat of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	

Field sprayer	Alfalfa (for seed), berries, bulb vegetables, carrots, cucurbits, fruiting vegetables, lettuce, mustard (oilseed and condiment), potato, strawberries, succulent beans, succulent peas, sunflowers		1	0	1
Chemigation	Alfalfa (for seed), mustard (oilseed and condiment)		1	0	1
Airblast	Grapes	Early growth stage	5	0	1
		Late growth stage	3	0	1
	Stone fruits	Early growth stage	10	0	2
		Late growth stage	4	0	1
Aerial	Alfalfa (for seed), potatoes, sunflowers	Fixed wing	1	0	15
		Rotary wing	1	0	10
	Mustard (oilseed and condiment)	Fixed and rotary wing	1	0	1

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under **DIRECTIONS FOR USE**.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

Boscalid is persistent and may carry over. It is recommended that this product not be used in areas treated with any products containing boscalid during the previous season.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, **CABIL FUNGICIDE** contains a Group 7 fungicide. Any fungal population may contain individuals naturally resistant to **CABIL FUNGICIDE** and other Group 7 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of **CABIL FUNGICIDE** or other Group 7 fungicides with different groups that control the same pathogens. Avoid application of consecutive sprays of **CABIL FUNGICIDE** or other fungicides in the same group in a season. **DO NOT** apply more than two (2) sequential applications of **CABIL FUNGICIDE** before alternating to a labeled fungicide with a different mode of action for at least one application. **DO NOT** exceed the total number of applications of **CABIL FUNGICIDE** per year as stated in the Crop-Specific Restrictions and Limitations.
- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor treated fungal populations for resistance development. Notify Sharda Cropchem Limited if reduced sensitivity of the pathogen to **CABIL FUNGICIDE** is suspected.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information and to report suspected resistance, contact Sharda Cropchem Limited Inc. at 1-844-810-5720.

MIXING

1. Ensure the spray tank is clean before use. Follow the clean-out recommendations stated on the label of the product that was previously used.
2. Fill the spray tank one-half full of water and start agitation.
3. Add the required amount of **CABIL FUNGICIDE** to the tank.
4. Add the required amount of the tank mix partner.
5. Continue agitation while filling the remainder of the spray tank.
6. After use, clean the spray tank according to PRECAUTIONS.

RESTRICTIONS AND LIMITATIONS

1. All crops except alfalfa can be grazed or fed to livestock.
2. A plant back restriction of 14 days is required for all crops not on the label.
3. **DO NOT** use in greenhouses.

PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. **DO NOT** take internally.
3. **DO NOT** get in eyes. Causes eye irritation. Avoid contact with skin and clothing.
4. Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking.
5. **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
6. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab and/or cockpit.
7. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
8. Clean spray equipment thoroughly after use.
9. Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.
10. When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

1. Store the leftover product in original tightly closed container.
2. **DO NOT** ship or store the product near food, feed, seed and fertilizers.
3. Store the product in cool, dry, locked, well-ventilated area without floor drain.

DISPOSAL

For Recyclable Containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

GROUP	7	FUNGICIDE
-------	---	-----------

CABIL FUNGICIDE

WETTABLE GRANULES

For use in alfalfa grown for seed production, berries group, bulb vegetables group, carrots, cucurbit vegetables, field lettuce (head and leaf), fruiting vegetables, grapes, potatoes, mustard (oilseed and condiment variety only), stone fruits group, strawberries, succulent beans, succulent peas and sunflowers.

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: Boscalid ... 70%

REGISTRATION NO. 34638 PEST CONTROL PRODUCTS ACT

WARNING - EYE IRRITANT

READ THE LABEL AND ATTACHED BROCHURE BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 0.1 kg – 10 kg

Sharda Cropchem Limited
2nd Floor, Prime Business Park
Dashrathlal Joshi Road
Vile Parle (West)
Mumbai - 400056, India

Canadian Agent:
Sharda Cropchem Limited
63 Kingsview Blvd
Etobicoke, Ontario, CA
M9R 1V1
1-844-810-5720
1-416-840-5639

PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. **DO NOT** take internally.
3. **DO NOT** get in eyes. Causes eye irritation. Avoid contact with skin and clothing.
4. Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking.
5. **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
6. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab and/or cockpit.
7. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
8. Clean spray equipment thoroughly after use.
9. Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.
10. When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.
11. As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.
12. **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

Boscalid is persistent and may carry over. It is recommended that this product not be used in areas treated with any products containing boscalid during the previous season.

STORAGE

1. Store the leftover product in original tightly closed container.
2. **DO NOT** ship or store the product near food, feed, seed and fertilizers.
3. Store the product in cool, dry, locked, well-ventilated area without floor drain.

DISPOSAL

For Recyclable Containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.