

2020-07-31
2020-2912

Container label

GROUP	3	FUNGICIDE
-------	----------	-----------

PRINCETON
FUNGICIDE

COMMERCIAL

Emulsifiable Concentrate for:

Broad-spectrum disease control in Wheat, Barley, Oats, Canola, Corn and Soybeans (grown for seed), Dry Edible Beans and for the suppression of Septoria Leaf Mottle on Canary Seed and Peaches, Nectarines, Plums, Sweet Cherries, Sour Cherries, Apricots, Highbush and Lowbush Blueberries, Saskatoon Berries, Cranberries, Caneberries, Strawberries, Rutabagas, Asparagus, Western Red Cedar, and Kentucky Bluegrass Grown for Seed.

ACTIVE INGREDIENT: Propiconazole 418 g/L

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

REGISTRATION NO.: 33840 PEST CONTROL PRODUCTS ACT

WARNING



POISON

EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

Net Contents: 1-1050 litres

SHARDA Cropchem Limited
2nd Floor, Prime Business Park
Dashrathlal Joshi Road
Vile Parle (West)
Mumbai - 400056, India

Canadian Agent:
SHARDA Cropchem Limited
63 Kingsview Blvd
Etobicoke, Ontario, CA
M9R 1V1
1-844-810-5720
1-416-840-5639

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take the container label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Apply symptomatic therapy.

PRECAUTIONS:

KEEP OUT OF THE REACH OF CHILDREN. Do not get in eyes, on skin or on clothing. Avoid breathing spray mist or vapours. Harmful or fatal if swallowed, inhaled or absorbed through the skin. Causes eye and skin irritation. This product may cause skin-sensitization reactions in certain individuals.

Wear coveralls over a long-sleeved shirt and long pants, socks, chemical-resistant footwear and chemical-resistant gloves during mixing, loading, application, clean-up and repair activities. Wear protective goggles or faceshield when handling the concentrated product. The wearing of neoprene gloves by pilots when entering the aircraft is essential. Mechanical flagging devices must be used.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Do not eat, drink or smoke during work; wash hands and face thoroughly before doing so. Immediately wash off accidental splashes of the concentrate or spray mixture from skin, clothing and out of eyes.

After work, change clothing and wash entire body thoroughly. Wash contaminated working clothes separately from other laundry before reuse. Do not contaminate food or feed.

NOTE: Do not graze animals on treated green crops within three days of application of PRINCETON Fungicide.

- For fruit and specialty crops as listed, do not graze livestock on treated green crops.

DO NOT allow entry into treated area for 12 hours following application. See the **DIRECTIONS FOR USE** section for crop specific restricted entry intervals.

STORAGE: To prevent contamination, store this product away from food or feed.

DISPOSAL:

DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

CONTAINER DISPOSAL:

For recyclable containers:

DO NOT REUSE THIS CONTAINER FOR ANY PURPOSE. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

ENVIRONMENTAL PRECAUTIONS:

Toxic to aquatic organisms and non-target terrestrial plants. The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body. Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application sites such as hedgerows and woodland.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Pamphlet label

GROUP	3	FUNGICIDE
-------	---	-----------

PRINCETON
FUNGICIDE

COMMERCIAL

Emulsifiable Concentrate for:

Broad-spectrum disease control in Wheat, Barley, Oats, Canola, Corn and Soybeans (grown for seed), Dry Edible Beans and for the suppression of Septoria Leaf Mottle on Canary Seed. Broad-spectrum disease control in Wheat, Barley, Oats, Canola, Corn and Soybeans (grown for seed), Dry Edible Beans and for the suppression of Septoria Leaf Mottle on Canary Seed and Peaches, Nectarines, Plums, Sweet Cherries, Sour Cherries, Apricots, Highbush and Lowbush Blueberries, Saskatoon Berries, Cranberries, Caneberries, Strawberries, Rutabagas, Asparagus, Western Red Cedar, and Kentucky Bluegrass Grown for Seed.

ACTIVE INGREDIENT: Propiconazole 418 g/L

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

REGISTRATION NO.: 33840 PEST CONTROL PRODUCTS ACT

WARNING



POISON

EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

Net Contents: 1-1050 litres

SHARDA Cropchem Limited
2nd Floor, Prime Business Park
Dashrathlal Joshi Road
Vile Parle (West)
Mumbai - 400056, India

Canadian Agent:
SHARDA Cropchem Limited
63 Kingsview Blvd
Etobicoke, Ontario, CA
M9R 1V1
1-844-810-5720
1-416-840-5639

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take the container label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Apply symptomatic therapy.

PRECAUTIONS:

KEEP OUT OF THE REACH OF CHILDREN. Do not get in eyes, on skin or on clothing. Avoid breathing spray mist or vapours. Harmful or fatal if swallowed, inhaled or absorbed through the skin. Causes eye and skin irritation. This product may cause skin-sensitization reactions in certain individuals.

Wear coveralls over a long-sleeved shirt and long pants, socks, chemical-resistant footwear and chemical-resistant gloves during mixing, loading, application, clean-up and repair activities. Wear protective goggles or faceshield when handling the concentrated product. The wearing of neoprene gloves by pilots when entering the aircraft is essential. Mechanical flagging devices must be used.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Do not eat, drink or smoke during work; wash hands and face thoroughly before doing so. Immediately wash off accidental splashes of the concentrate or spray mixture from skin, clothing and out of eyes.

After work, change clothing and wash entire body thoroughly. Wash contaminated working clothes separately from other laundry before reuse. Do not contaminate food or feed.

DO NOT allow entry into treated area for 12 hours following application. See the **DIRECTIONS FOR USE** section for crop specific restricted entry intervals.

NOTE: Do not graze animals on treated green crops within three days of application of PRINCETON FUNGICIDE.

- For fruit and specialty crops as listed, do not graze livestock on treated green crops.

STORAGE: To prevent contamination, store this product away from food or feed.

DISPOSAL:

DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

CONTAINER DISPOSAL:

For recyclable containers:

DO NOT REUSE THIS CONTAINER FOR ANY PURPOSE. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

ENVIRONMENTAL PRECAUTIONS:

Toxic to aquatic organisms and non-target terrestrial plants. The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body. Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application sites such as hedgerows and woodland.

DIRECTIONS FOR USE:

DO NOT apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, ditches and wetlands), estuaries or marine habitats.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. DO NOT apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 m/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotor span.

GENERAL INFORMATION:

Introduction: PRINCETON FUNGICIDE is a broad-spectrum systemic fungicide for control of a wide range of diseases on certain crops. PRINCETON FUNGICIDE will protect the crop from yield and quality losses due to disease. PRINCETON FUNGICIDE may be used in conjunction with higher seeding rates, higher fertilizer inputs, plant growth regulators and other fungicides as required. (See “NOTE” under PRECAUTIONS)

DO NOT use in greenhouses.

Crop Information: PRINCETON FUNGICIDE may be applied to:

Winter Wheat
Spring Wheat (including Hard Red, Durum, Canada Prairie, Soft White)
Spring Barley
Oats
Canola
Corn (including Field, Seed and Sweet)

Soybeans (grown for seed)
 Canary Seed
 Dry Edible Beans (including kidney, navy and white bean)

Diseases Controlled:

Winter Wheat and Spring Wheat (including Hard Red, Durum, Canada Prairie, Soft White)	Septoria Leaf Spot; Septoria Glume Blotch; Powdery Mildew; Leaf and Stem Rust; Tan Spot; Stripe Rust
Spring Barley	Net Blotch; Spot Blotch; Scald; Powdery Mildew; Septoria Leaf Spot; Leaf Rust; Stem Rust
Oats	Septoria Leaf Blotch; Crown Rust
Canola	Blackleg
Corn	Rusts, Northern Corn Leaf Blight, Southern Corn Leaf Blight, Helminthosporium Leaf Spot, Eye Spot, Grey Leaf Spot
Soybeans (grown for seed)	Frogeye Leaf Spot (<i>Cercospora spp.</i>), Aerial Web Blight (<i>Rhizoctonia solani</i>)
Canary Seed	Septoria Leaf Mottle
Dry Edible Beans (including kidney, navy and white beans)	Rust

Factors Affecting PRINCETON FUNGICIDE Performance: PRINCETON FUNGICIDE should be applied as a preventative disease control measure. Established diseases are more difficult to control and may have already reduced crop vigour.

If rainfall occurs within one hour of application, reapplication is necessary.

AERIAL APPLICATION:

DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. High humidity and low temperatures (10 – 20°C) allow for a better deposition of spray droplets.

Apply only by fixed-wing or rotary aircraft equipment that has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

SPRAYER AND APPLICATION INFORMATION:

The performance of this product depends on correct application. Follow the guidelines given below for optimal application of PRINCETON FUNGICIDE.

Sprayer Information:

	GROUND APPLICATION	AERIAL APPLICATION
Spray Volume	Minimum 200 L of water per hectare	40 - 50 L of water per hectare
Spray Pressure	200 - 300 kPa	100 - 200 kPa
Nozzle Type	110° Flat Fan (XR11004, 4110-20)	Flat Fan 6510-6515 or Hollow Cone (D8-45)
Droplet Size	Medium Spray (300 - 400 microns VMD)	Medium Spray (350 - 400 microns VMD)
Ground Speed	10 km/h	
Nozzle Angle	90° (straight down)	90° (straight down)
Boom Height	40-50 cm above the crop canopy	2-3 m above the crop canopy

Ground Application:

Mixing and Spraying Instructions:

Spray equipment should be thoroughly flushed with clean water before mixing PRINCETON FUNGICIDE.

Fill spray tank 1/2 full with clean water. Engage gentle agitation.

Add the required amount of PRINCETON FUNGICIDE and agitate thoroughly.

Continue filling the tank with water until the tank is 9/10 full and, if applicable, add the required amount of tank mix partner.

Complete filling the spray tank with water, maintaining agitation during mixing and spraying operations.

Use nozzle screens no finer than 50 mesh. Keep by-pass line on or near the bottom of the tank to minimize foaming.

Aerial Application:

Mixing and Spraying Instructions:

Spray equipment should be thoroughly flushed with clean water before mixing PRINCETON FUNGICIDE.

Fill premix tank 1/2 full with clean water. Engage gentle agitation.

Add the required amount of PRINCETON FUNGICIDE and agitate thoroughly.

Continue filling the tank with water until the tank is 9/10 full and, if applicable, add the required amount of tank mix partner.

Complete filling the premix tank with water.

Maintain gentle agitation during mixing.

Transfer the premix contents into the aircraft spray tank.

Maintain sufficient agitation during the mixing and spraying operation to ensure PRINCETON FUNGICIDE remains in suspension.

Use nozzle screens no finer than 50 mesh. Keep by-pass line on or near the bottom of the tank to minimize foaming.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for a specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

AERIAL APPLICATION USE PRECAUTIONS:

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas.

Coarse sprays are less likely to drift; therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty.

AERIAL APPLICATION OPERATOR PRECAUTIONS:

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

PRODUCT SPECIFIC PRECAUTIONS:

Read and understand the entire label before opening this product. If you have questions call the manufacturer at 1-844-810-5720 or obtain technical advice from the distributor or your provincial agricultural representative.

Application of this specific product must meet and/or conform to the following:

Apply the recommended rate in a minimum spray volume of 40 litres per hectare.

DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application.

High humidity and low temperatures (10 - 20 ° C) allow for a better deposition of spray droplets.

Fertilizer:

Mixing and Spraying Instructions:

If desired, small amounts of nitrogen may be applied with PRINCETON FUNGICIDE. The appropriate amount of urea can be dissolved in water and added to the spray tank before adding PRINCETON FUNGICIDE. The rate of actual nitrogen must not exceed 10 kg/ha.

CAUTION: Excessive nitrogen concentrations may injure the crop.

NOTE: DO NOT add nitrogen when tank-mixing PRINCETON FUNGICIDE with an herbicide.

Buffer zones:

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats

(such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method Of Application	Crop		Buffer Zones (metres) Required for the Protection of:				Terrestrial habitat
			Freshwater Habitats of Depths:		Estuarine/Marine Habitats of Depths:		
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer*	rutabagas, cranberries, asparagus, Kentucky bluegrass, Western cedar		1	0	1	1	1
Airblast	Cherries	Early Growth Stage	5	0	10	3	10
		Late Growth Stage	2	0	4	2	4
	Blueberries, apricots, nectarines, peaches, plums, Saskatoon berries, blackberries, loganberrie, raspberries, other berries	Early Growth Stage	4	0	5	2	5
		Late Growth Stage	2	0	3	1	3
Aerial	Blueberries, Kentucky bluegrass (seed prod.)	Fixed Wing	1	0	3	1	20
		Rotary Wing	1	0	1	1	20

For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

WHEAT, BARLEY AND OATS SPRAY SCHEDULE INSTRUCTIONS FOR USE:

Apply PRINCETON FUNGICIDE at the very early stages of disease. This could occur anytime during tillering or stem elongation. Typically, an application from the beginning of stem elongation up to flag leaf emergence is required.

PRINCETON FUNGICIDE lasts about three weeks in the plant. If conditions favourable to disease continue after this length of time, another application will be necessary to maintain control. The second spray is usually applied at the time of head emergence. In most cases, this second application is essential to maintain control of the Septoria disease complex.

CROP/DISEASE	RATE/HA	EARLY APPLICATION	LATER APPLICATION	MAXIMUM NUMBER OF APPLICATIONS PER SEASON
WHEAT Septoria Leaf Spot and Tan Spot SPRING BARLEY * Net Blotch	150-300 mL	At Growth Stage (G.S). 12-23 (as early as the two leaf stage). For early season disease suppression, use the lower rate for suppression under normal field conditions. Use the higher rate for control if there is a history of high disease pressures in the field and/or field conditions favour disease development.	At the first sign of disease (G.S. 29-37) or before head is half emerged (G.S. 49-55). Apply only the high rate on any application from G.S. 29-55.	2 For net blotch on barley, in areas of high disease pressure, do not apply more than one (1) application of PRINCETON Fungicide before alternating to a labelled fungicide with a different mode of action.
WHEAT Septoria Leaf Spot and Glume Blotch, Powdery Mildew, Leaf and Stem Rust, Tan Spot, Stripe Rust SPRING BARLEY Net Blotch, Spot Blotch, Scald, Powdery Mildew Septoria Leaf Spot, Leaf and Stem Rust OATS Septoria Leaf Blotch, Crown Rust	300 mL	At the first sign of disease, usually at the beginning of stem elongation (G.S. 29-37)	Before head is half emerged (G.S. 49-55)	2 For powdery mildew, net blotch and scald: in areas of high disease pressure, do not apply more than one (1) application of PRINCETON Fungicide before alternating to a labelled fungicide with a different mode of action.
LAST APPLICATION MUST BE MADE AT LEAST 45 DAYS BEFORE HARVEST.				

HERBICIDE TANK-MIXING - WHEAT & BARLEY:

PRINCETON FUNGICIDE can be tank-mixed with ONLY ONE of these herbicides:

2,4-D Amine

MCPA Amine

Badge Emulsifiable Selective Weedkiller Bromotril 240 EC

HORIZON 240 EC

When applied as a tank-mix combination, read and observe all label directions, including rates, personal protective equipment, restrictions and precautions for each product used in the tank-mix. Always use in accordance with the most restrictive label restrictions and precautions.

Tank-mixing Precautions:

Do not tank-mix PRINCETON FUNGICIDE with herbicides for application onto Oats.

Weeds and crops must be at the correct stage of growth as specified in both the PRINCETON FUNGICIDE label and the tank mix partner label.

2,4-D and MCPA formulations may be applied either by ground application or aerial application; tank-mixtures of PRINCETON FUNGICIDE and Badge Emulsifiable Selective Weedkiller or Bromotril 240 EC can only be applied by ground application.

Consult the label of the herbicide partner for a list of weeds controlled, directions for use and precautions.

Compatibility should always be confirmed by premixing small proportional quantities of water, PRINCETON FUNGICIDE, and the tank-mix partner in advance.

NOTE: Do not graze animals on treated green crops within three days of application of PRINCETON FUNGICIDE. Do not feed straw from crops treated with herbicide tank mixes to livestock.

HERBICIDE TANK-MIXING – WHEAT

FOR USE ONLY IN THE PRAIRIE AND PEACE RIVER, OKANAGAN AND CRESTON FLATS REGIONS OF BRITISH COLUMBIA.

PRINCETON FUNGICIDE can be tank-mixed with HORIZON 240 EC Herbicide for disease and grassy weed control.

When applied as a tank-mix combination, read and observe all label directions, including rates, personal protective equipment, restrictions and precautions for each product used in the tank-mix. Always use in accordance with the most restrictive label restrictions and precautions.

Tank-mixing Precautions:

Do not apply by air.

Consult the label of HORIZON 240 EC Herbicide for a list of weeds controlled, directions for use and precautions.

Apply prior to emergence of the 4th tiller (herbicide timing).

CANOLA SPRAY SCHEDULE

INSTRUCTIONS FOR USE:

PRINCETON FUNGICIDE will control blackleg and enhance yield potential during the early stages of canola growth. The disease may reappear later in the season, but with minimal effect on yield. DO NOT APPLY BY AIR.

DISEASE	RATE/HA	REMARKS/TIMING
Blackleg	300 mL	Apply during the rosette stage; between 2 nd true leaf and bolting.
		Seedling Rosette Bud (Bolted) Stage 1 Stage 2 Stage 3
LAST APPLICATION MUST BE MADE AT LEAST 60 DAYS BEFORE HARVEST.		

TANK-MIXING – SEED CORN, FIELD CORN and SWEET CORN

PRINCETON FUNGICIDE can be tank-mixed with the following partner:

Silencer 120 EC Insecticide

A restricted entry interval of 1 day is required for workers hand-harvesting and detasseling treated corn.

When applied as a tank-mix combination, read and observe all label directions, including rates, personal protective equipment, restrictions and precautions for each product used in the tank-mix. Always use in accordance with the most restrictive label restrictions and precautions.

Tank-mixing Precautions:

The tank mix of PRINCETON FUNGICIDE + Silencer 120 EC Insecticide can be applied by air and ground. Use 40 L of water per hectare when applying by air.

Insects and crops must be at the correct stage as specified on the PRINCETON FUNGICIDE as well as the Silencer 120 EC label. Follow the directions for use and precautions on all labels.

A Pre harvest interval (PHI) of 14 days must be respected when using these tank-mixes on field and sweet corn.

Compatibility should always be confirmed by premixing small proportional quantities of water, PRINCETON FUNGICIDE, and the tank-mix partner in advance.

SEED CORN, FIELD CORN AND SWEET CORN SPRAY SCHEDULE

DISEASE	RATE/HA PRINCETON Fungicide alone	RATE/HA Tank Mix partner	REMARKS
Rusts	300 mL	+ 83 mL Silencer 120 EC	Apply 300 mL of PRINCETON FUNGICIDE per hectare when rust pustules first appear. Under severe disease pressure, make a second application 14 days after. Seed corn only , under severe disease pressure, make a third application 14 days after.
Northern Corn Leaf Blight Southern Corn Leaf Blight Helminthosporium Leaf Spot	150-300 mL	+ 83 mL Silencer 120 EC	Apply 150 - 300 mL of PRINCETON FUNGICIDE per hectare when disease first appears. Use the 150 mL rate if disease pressure is low..
Eye Spot Grey Leaf Spot	300 mL	+ 83 mL Silencer 120 EC	Apply 300 mL of PRINCETON FUNGICIDE per hectare when disease first appears.

SOYBEANS GROWN FOR SEED SPRAY SCHEDULE

DISEASE	RATE/HA	REMARKS
Frogeye Leaf Spot (<i>Cercospora spp.</i>) Aerial Web Blight (<i>Rhizoctonia solani</i>)	300-455 mL	Apply 300 - 455 mL of PRINCETON FUNGICIDE per hectare, using ground application equipment only, when disease first appears. Under severe disease pressure, make a second application 14 days after the first application.
Do not harvest soybeans within 50 days of the last application. Harvested soybean seed should not be used for human food or animal feed. For use only on soybeans grown for seed.		

CANARY SEED SPRAY SCHEDULE

DISEASE	RATE/HA	REMARKS
Septoria leaf mottle	300 mL	For the suppression of Septoria leaf mottle; make one application at emergence of flag leaf; ground application only; apply in 200 L water/ha.

DRY EDIBLE BEANS SPRAY SCHEDULE

DISEASE	RATE/HA	REMARKS
Rust	300 mL	Apply 300 mL of PRINCETON FUNGICIDE in minimum of 200 L of water per hectare by ground application or in 40 to 50 L of water per hectare by aerial application at the first detection of disease in the field and a second application 14 to 21 days later.
APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF PRINCETON FUNGICIDE TO DRY EDIBLE BEANS. DO NOT APPLY WITHIN 28 DAYS OF HARVEST.		

DO NOT use in greenhouses.

Diseases Controlled: Cranberries	Cottonball (<i>Monilinia oxycocci</i>)
Kentucky Bluegrass Grown for Seed	Powdery Mildew
Lowbush Blueberries	Monilinia Blight (Mummyberry)
Highbush Blueberries	Mummyberry (<i>Monilinia vaccinii – corymbosi</i>)
Peaches, Nectarines, Plums, Apricots	Brown Rot Blossom Blight, Fruit Brown Rot
Sweet and Sour Cherries	Brown Rot Blossom Blight, Fruit Brown Rot, Cherry Leaf Spot (<i>Blumeriella jaapii</i>)
Plums and Sour Cherries	Black Knot (<i>Apiosporina morbosa</i>) [suppression only]
Rutabagas	Powdery Mildew
Asparagus	Rust (<i>Puccinia asparagi</i>)
Saskatoon Berry	Entomosporium Leaf and Berry Spot, Saskatoon Juniper Rust
Western Red Cedar	Keithia Foliar Blight

Factors Affecting PRINCETON FUNGICIDE Performance: PRINCETON FUNGICIDE should be applied as a preventative disease control measure. Established diseases are more difficult to control and may have already reduced crop vigour.

If rainfall occurs within one hour of application, reapplication is necessary.

DIRECTIONS FOR USE:

DO NOT apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, ditches and wetlands), estuaries or marine habitats.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

NOTE: Do not graze livestock on treated crop.

PEACHES, NECTARINES, PLUMS, AND APRICOTS - SPRAY SCHEDULE		
DISEASE	RATE/HA	REMARKS
Brown Rot Blossom Blight	300 mL	Apply 300 mL of PRINCETON FUNGICIDE in a minimum of 500 L of water per hectare by ground. Make 1st application at early bloom with a 2nd application at 50% - 75% bloom. If disease conditions persist, make a 3rd application at petal fall.
Fruit Brown Rot	300 mL	Apply no more than 2 applications in the 3 weeks prior to harvest. Apply 300 mL of product in a minimum of 500 L of water per hectare by ground.
Suppression of Black Knot (Plums only)	300 mL	Apply 300 mL of PRINCETON FUNGICIDE in a minimum of 500 L of water per hectare by ground. Make 1st application at early bloom with a 2nd application at 50% - 75% bloom. If disease conditions persist, make a 3rd application at petal fall.
DO NOT APPLY WITHIN 3 DAYS OF HARVEST. DO NOT REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION. IF REQUIRED, INDIVIDUALS MAY REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION FOR SHORT TERM TASKS WHICH DO NOT REQUIRE HAND LABOUR IF AT LEAST 4 HOURS HAVE PASSED SINCE APPLICATION AND PROVIDED LONG PANTS, LONG- SLEEVED SHIRTS, AND CHEMICAL-RESISTANT GLOVES ARE WORN.		

SWEET AND SOUR CHERRIES - SPRAY SCHEDULE		
DISEASE	RATE/HA	REMARKS
Brown Rot Blossom Blight	300 mL	Apply 300 mL of PRINCETON FUNGICIDE in a minimum of 500 L of water per hectare by ground. Make 1st application at early bloom with a 2nd application at 50% - 75% bloom. If disease conditions persist, make a 3rd application at petal fall.
Fruit Brown Rot	300 mL	Apply no more than 2 applications in the 3 weeks prior to harvest. Apply 300 mL of product in a minimum of 500 L of water per hectare by ground.
Cherry Leaf Spot (<i>Blumeriella jaapii</i>)	300 mL	Apply a maximum of 3 applications per season for control of Cherry Leaf Spot. Make the 1st application at petal fall. In the 3 weeks prior to harvest make a 2nd and 3rd application at a 7- 10 day interval. Do not apply within 3 days of harvest. Apply 300 mL of product in a minimum of 500 L of water per hectare by ground.
Suppression of Black Knot (sour cherries only)	300 mL	Apply 300 mL of PRINCETON FUNGICIDE in a minimum of 500 L of water per hectare by ground. Make 1st application at early bloom with a 2nd application at 50% - 75% bloom. If disease conditions persist, make a 3rd application at petal fall.
It is recommended that no more than 2 consecutive applications of PRINCETON FUNGICIDE be made before switching to another fungicide with a different mode of action according to disease management practices.		

DO NOT APPLY WITHIN 3 DAYS OF HARVEST. DO NOT REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION. IF REQUIRED, INDIVIDUALS MAY REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION FOR SHORT TERM TASKS WHICH DO NOT REQUIRE HAND LABOUR IF AT LEAST 4 HOURS HAVE PASSED SINCE APPLICATION AND PROVIDED LONG PANTS, LONG- SLEEVED SHIRTS, AND CHEMICAL-RESISTANT GLOVES ARE WORN.

HIGHBUSH BLUEBERRY - SPRAY SCHEDULE

DISEASE	RATE/HA	REMARKS
Mummyberry (<i>Monilinia vaccinii – corymbosi</i>)	300 mL	Apply 1 st application at or near flower bud swelling; make a 2nd application at leaf bud swelling. Apply by ground only, making no more than two applications per year. In BC only, apply by ground, a 3 rd application at pink bloom and a 4 th application 7 to 10 days later at early bloom, making no more than 4 applications per year. Use a minimum of 200 L of water per hectare.

LAST APPLICATION MUST BE MADE AT LEAST 60 DAYS BEFORE HARVEST. A restricted entry interval of 5 days is required for workers hand pruning highbush blueberries.

LOWBUSH BLUEBERRY - SPRAY SCHEDULE

DISEASE	RATE/HA	REMARKS
Monilinia Blight	300 mL	Apply 1 st application when flower bud scales first appear and make a 2 nd application 10 days later. Use ground application or aerial application equipment, making no more than two applications per year. Use a minimum of 200 L of water per hectare if applying by ground equipment; use 40 – 50 L of water per hectare if applying by air.

LAST APPLICATION MUST BE MADE AT LEAST 60 DAYS BEFORE HARVEST.

SASKATOON BERRY - SPRAY SCHEDULE

DISEASE	RATE/HA	REMARKS
Entomosporium Leaf and Berry Spot, Saskatoon Juniper Rust	300 mL	As a foliar spray, apply up to three applications per season. The 1st application to occur at “white tip”, the 2nd application at “petal fall”, and the 3rd application at “green fruit”. Apply 300 mL in a minimum of 200 L of water per hectare by ground, applying to runoff.

LAST APPLICATION MUST BE MADE AT LEAST 38 DAYS BEFORE HARVEST.

CRANBERRY - SPRAY SCHEDULE

DISEASE	RATE/HA	REMARKS
Cottonball (<i>Monilinia oxycocci</i>)	300 mL	Apply the 1st application at leaf bud break. Make a 2nd application 10 – 14 days later, a 3rd application at early bloom and a 4th application 10 – 14 days after the 3rd application. Make no more than four applications per year. Apply product by ground.

LAST APPLICATION MUST BE MADE AT LEAST 45 DAYS BEFORE HARVEST.

RUTABAGAS - SPRAY SCHEDULE		
DISEASE	RATE/HA	REMARKS
Powdery Mildew	240 mL	Make two applications per season with the 1st application at 50 days after planting and the 2nd application 20 days later. Apply to vegetative foliage. Apply 240 mL in a minimum 200 L of water per hectare by ground.
LAST APPLICATION MUST BE MADE AT LEAST 21 DAYS BEFORE HARVEST.		

ASPARAGUS - SPRAY SCHEDULE		
DISEASE	RATE/HA	REMARKS
Rust (<i>Puccinia asparagi</i>)	150 mL	Apply PRINCETON FUNGICIDE to asparagus ferns in Ontario and Quebec only. Once harvest is complete, make the 1st application of PRINCETON FUNGICIDE as soon as fern growth begins, followed by applications at 14 to 21 day intervals. For new, non-harvested plantings, apply PRINCETON FUNGICIDE when first sign of rust is visible, followed by applications at 14 to 21 day intervals. Apply by ground only, making no more than three applications per year. Use a minimum of 370 L of water per hectare.
LAST APPLICATION MUST BE MADE AT LEAST 8 MONTHS BEFORE HARVEST.		

WESTERN RED CEDAR - SPRAY SCHEDULE		
DISEASE	RATE/HA	REMARKS
Keithia Foliar Blight	300 mL	Apply using ground application equipment every four weeks. Make a maximum of 6 applications per year. Apply 300 mL of PRINCETON FUNGICIDE in a volume of 1000 L of water per hectare by ground.

KENTUCKY BLUEGRASS FOR SEED PRODUCTION - SPRAY SCHEDULE		
DISEASE	RATE/HA	REMARKS
Powdery Mildew	300 mL	Apply as a foliar spray. Make no more than 2 applications per crop year with the 1st application at pre-heading and the 2nd at 50% - 100% heading. Apply in 200 - 300 L/ha of water by ground or 40 - 50 L/ha of water by air.

CANEERRIES (CROP GROUP 13A; CULTIVARS AND/OR VARIETIES OF RED AND BLACK RASPBERRY, LOGAN BERRY AND BLACKBERRY) - SPRAY SCHEDULE		
DISEASE	RATE/HA	REMARKS
Yellow rust	300 mL	Apply 300 mL of PRINCETON FUNGICIDE in a minimum of 500 L of water per hectare by ground application at first detection of disease in the field and a second application 14 days later.

It is recommended that no more than 2 consecutive applications of PRINCETON FUNGICIDE be made before switching to another fungicide with a different mode of action according to disease management practices.

APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF PRINCETON FUNGICIDE TO CANEBERRIES. DO NOT APPLY WITHIN 30 DAYS OF HARVEST. DO NOT REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION. IF REQUIRED, INDIVIDUALS MAY REENTER TREATED AREAS WITHIN 3 DAYS OF APPLICATION FOR SHORT TERM TASKS WHICH DO NOT REQUIRE HAND LABOUR IF AT LEAST 4 HOURS HAS PASSED SINCE APPLICATION AND PROVIDED LONG PANTS, LONG SLEEVED SHIRTS, AND CHEMICAL RESISTANT GLOVES ARE WORN.

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Sharda Cropchem Limited under the User Requested Minor Use Label Expansion program. For these uses, Sharda Cropchem Limited has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

STRAWBERRIES – SPRAY SCHEDULE

DISEASE	RATE/HA	REMA
Leaf Spot (<i>Mycosphaerella fragariae</i>)	300 mL	Apply 300 mL of PRINCETON FUNGICIDE per hectare by ground in enough water to ensure thorough coverage. Make 1st application when disease levels are no more than 5%. Apply PRINCETON FUNGICIDE at 10 day intervals

It is recommended that no more than 2 consecutive applications of PRINCETON FUNGICIDE be made before switching to another fungicide with a different mode of action according to disease management practices.

APPLY A MAXIMUM OF 4 APPLICATIONS PER SEASON OF PRINCETON FUNGICIDE TO STRAWBERRIES. DO NOT APPLY WITHIN 1 DAY OF HARVEST.

RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, PRINCETON FUNGICIDE contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to PRINCETON FUNGICIDE and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of PRINCETON FUNGICIDE or other Group 3 fungicides with different groups that control the same pathogens.
- Avoid application of more than the maximum number listed in the label and consecutive sprays of PRINCETON FUNGICIDE or other fungicides in the same group in a season.
- Use tank mixtures with fungicide from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an Integrated Disease Management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor treated fungal populations for sign of resistance development. Notify Sharda Cropchem Limited if reduced sensitivity of the pathogen to PRINCETON FUNGICIDE is suspected.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available. Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information or to report suspected resistance, contact Sharda Cropchem Limited at 1-844-810-5720.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

HORIZON is a registered trademark of a Syngenta group company.