

2024-5612  
2024-11-15

GROUP	2	HERBICIDE
-------	---	-----------

Dakota™

Selective Herbicide

Water Dispersible Granular Herbicide in water-soluble bags

COMMERCIAL (AGRICULTURAL)

FOR SALE FOR USE IN EASTERN CANADA, THE PRAIRIE PROVINCES AND INTERIOR  
OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY

ACTIVE INGREDIENT: Imazamox ..... 70%

REGISTRATION NO 34698

PEST CONTROL PRODUCTS ACT

NET CONTENTS: 470 g (4 x 117.5 g water-soluble bags)

Warning, contains the allergen milk  
Caution - Skin Irritant

READ THE LABEL AND THIS BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN  
DO NOT SELL INDIVIDUAL WATER-SOLUBLE BAGS SEPARATELY

Albaugh LLC  
1525 NE 36<sup>th</sup> Street,  
Ankeny, Iowa  
United States  
1-800-247-8013

**PRECAUTIONS****KEEP OUT OF REACH OF CHILDREN.**

DO NOT APPLY BY AIR

May irritate the skin. Avoid contact with skin

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

**FIRST AID**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION**

Treat symptomatically.

For medical or treatment information from exposure to this product, call 1-888-347-6732 (7 days/week, 24-hr).

For 24-hour chemical spill, leak, fire, exposure, or accident response information, call CHEMTREC toll free at 1-800-424-9300.

**STORAGE**

Store this product away from food or feed.

Store above 5°C.

Store in a cool, dry place away from children, animals, or fertilizers.

Keep unused water-soluble bags in resealed original container.

Keep packages dry at all times.

Keep packages out of rain.

Keep from freezing.

## DISPOSAL

1. When all the water-soluble bags are used, make the empty outer package unsuitable for further use.
2. Dispose of the outer package in accordance with provincial requirements.
3. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

## ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

## DIRECTIONS FOR USE:

**For use only on soybeans in Eastern Canada only and field peas, imazamox tolerant canola, imazamox tolerant lentils, imazamox tolerant sunflowers, and imazamox tolerant canola quality *Brassica juncea* in the Prairie Provinces and Peace River Area of British Columbia only.**

### DIRECTIONS FOR USE (Eastern Canada Only)

Dakota is a selective herbicide that can be applied as an early post-emergence treatment (cotyledon to 4-leaf stage) in soybeans.

DO NOT graze the treated soybeans or cut for hay within 20 days of application.

Pre-harvest Interval: After 60 days, soybeans can be harvested.

### DIRECTIONS FOR USE (Prairie Provinces and Peace River Area of British Columbia only)

Dakota is a selective herbicide that can be applied as an early post-emergence treatment in field peas, imazamox tolerant canola, imazamox tolerant lentils, imazamox tolerant sunflowers and imazamox tolerant canola quality *Brassica juncea*.

When Dakota is applied early post-emergence, absorption may occur through both the roots and foliage. Susceptible weeds stop growing and eventually die.

DO NOT apply more than once per year.

DO NOT graze the treated field peas, imazamox tolerant lentils, imazamox tolerant canola or imazamox tolerant canola quality *Brassica juncea* or cut for hay within 20 days of application.

DO NOT graze treated sunflower plants or cut for straw; sufficient data are not available to support such use.

Pre-harvest Interval: After 60 days, field peas, dry edible beans (*Phaseolus vulgaris*), imazamox tolerant lentils, imazamox tolerant canola or imazamox tolerant canola quality *Brassica juncea* can be harvested.

### Field Sprayer Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply by air.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

### Spray Buffer Zones

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label,
- low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands).

Method of Application	Crop	Spray Buffer zones (metres) Required for the Protection of Terrestrial Habitat
Field sprayer	All crops	1

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

### **BROADLEAF WEED AND GRASS CONTROL (Prairie Provinces and Peace River Area of British Columbia only)**

Dakota at 21 g/ha (15 g a.e./ha) will control only:	
Lamb's-quarters*	Volunteer spring wheat
Stinkweed	(non-imazamox tolerant wheat)
Wild mustard	
Dakota at 29 g/ha (20 g a.e./ha) will control:	
Barnyard grass	Volunteer barley
Cleavers*	Volunteer canary seed
Cow cockle	Volunteer canola (non-imazamox tolerant canola only)
Green foxtail	Volunteer durum wheat
Green smartweed	Volunteer spring wheat (non-imazamox tolerant wheat)
Japanese brome grass*	Volunteer tame oats
Kochia*	Wild buckwheat*
Lamb's-quarters	Wild mustard
Persian dandelion	Wild oats
Redroot pigweed	Yellow foxtail
Round-leaved mallow*	
Russian thistle	
Shepherd's purse	
Stinkweed	

\* = suppression only

Apply in 100 L/ha of water with MERGE® Adjuvant at a rate of 0.5% v/v (e.g. 5 L of MERGE Adjuvant per 1000 L spray solution).

#### **Imazamox Tolerant Canola**

Application should be made from the 2 to 6 true leaf stage of the canola crop and after the weeds have emerged. To control broadleaf weeds, apply Dakota from the cotyledon to the 4-leaf stage. To control grasses, apply Dakota from the 1 to 4-true leaf stage until early tillering. Initial transient crop yellowing may be observed after application, but this is outgrown and should not affect yield.

#### **Imazamox Tolerant Canola Quality *Brassica juncea***

Application should be from the 2 to 6 true leaf stage of the crop and after the weeds have emerged. To control broadleaf weeds, apply Dakota from the cotyledon to the 4-leaf stage. To control grasses, apply Dakota from the 1 to 4 true leaf stage until early tillering. Initial transient crop yellowing may be observed after application, but this is outgrown and should not affect yield.

#### **Imazamox Tolerant Sunflowers**

Application should be made from the 2 to 8 leaf stage of the sunflower crop and after the weeds have emerged. To control broadleaf weeds, apply Dakota from the cotyledon up to the 4-leaf stage. To control grasses, apply Dakota from the 1 to 4-true leaf stage up until early tillering. Initial transient crop yellowing may be observed after application, but this is outgrown and should not affect yield.

Apply in 100 L/ha of water with MERGE® Adjuvant at a rate of 0.5% v/v (e.g. 5 L of MERGE Adjuvant per 1000 L spray solution).

### Imazamox Tolerant Lentils

Application should be made from the 2 to 6 leaf stage of the lentil crop and after the weeds have emerged. To control broadleaf weeds, apply Dakota from the cotyledon up to the 4-leaf stage. To control grasses, apply Dakota from the 1 to 4-true leaf stage up until early tillering.

### For Field Peas

Application should be made from the 1 to 6 true leaf stage and after weeds have emerged. Apply when broadleaf weeds are from the cotyledon to 4-leaf stage and when grassy weeds are at the 1 to 4 true leaf or early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.

### Tank-Mix Combination

#### Dakota + Basagran Forte Herbicide in Field Peas

To provide broad-spectrum weed control, Dakota may be tank-mixed with Basagran Forte Herbicide in field peas.

<b>Timing</b>	Early post-emergence	
<b>Rate</b>	29 g/ha Dakota + 894 mL/ha Basagran Forte Herbicide + 2 L/ha nitrogen source* (UAN 28%)	
<b>Water Volume</b>	100 L/ha	
<b>Weeds Controlled</b>	This tank-mix combination will provide enhanced weed control vs. Dakota alone, as well as control of weeds listed on Dakota label. Refer to the Broadleaf Weed and Grass Control Section of this label.	
	<b>Control</b>	<b>Suppression</b>
	Volunteer Imazamox Tolerant Canola	Wild buckwheat
	Wild mustard (Group 2 resistant) Lentils (including Imazamox Tolerant Lentils)	Cleavers (including Group 2 resistant biotypes)
<b>Remark</b>	Application should be made from the 3 - 6 node stage of field peas and after weeds have emerged. Apply when broadleaf weeds are from the cotyledon to 4-leaf stage and when grassy weeds are at the 1 - 4 true leaf or early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield. For tank mixing, see mixing instructions.	

\*Nitrogen source: A reduction in grass control can be observed without the addition of nitrogen source (UAN 28%).

**Dakota + Basagran Forte Herbicide in Dry Edible Beans (all types of the species *Phaseolus vulgaris*\*)**

To provide broad-spectrum weed control and to reduce the potential for crop injury Dakota must be tank-mixed with Basagran Forte Herbicide in dry edible beans.

**NOTE TO USER**

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Albaugh LLC under the User Requested Minor Use Label Expansion program. For these uses, Albaugh LLC has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

Dry common bean varieties may vary in their tolerance to herbicides, including to Dakota + Basagran Forte Herbicide tank mix. In particular, white (navy) beans are more susceptible to herbicide injury which can result in delayed maturity. Since not all dry common bean varieties have been tested for tolerance to Dakota + Basagran Forte Herbicide tank mix, first use of this tank mix should be limited to a small area of each variety to confirm tolerance prior to adoption as a general field practice.

<b>Timing</b>	Early post-emergence	
<b>Rate</b>	0.029 kg/ha Dakota + 1.25 L/ha Basagran Forte Herbicide + 2 L/ha nitrogen source** (UAN 28%)	
<b>Water Volume</b>	100 L/ha	
<b>Weeds Controlled</b>	This tank-mix combination will provide enhanced weed control vs. Dakota alone, as well as control of weeds listed on Dakota label. Refer to the Broadleaf Weed and Grass Control Section of this label.	
	<b>Control</b>	<b>Suppression</b>
	Group 2 resistant wild mustard	Prostrate pigweed
	Stork's bill	Spiny annual sow thistle
	Volunteer canola [including biotypes with the Imazamox Tolerant trait)	Wild buckwheat
	**Nitrogen source: A reduction in grass control can be observed without the addition of nitrogen source (UAN 28%).	
<b>Remark</b>	<p>Dakota is only registered for use on dry beans when tank-mixed with Basagran Forte Herbicide + UAN.            Application should be made after 1st trifoliolate leaf has fully expanded up to the 2nd trifoliolate leaf stage of the dry bean and after weeds have emerged. Apply when broadleaf weeds are from the cotyledon to 4-leaf stage and when grassy weeds are at the 1 - 4 true leaf or early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.            For tank mixing, see mixing instructions.</p> <p>Do not harvest the treated crop within 60 days of application.</p>	

<b>*Warning</b>	Only the following dry bean types have been tested and demonstrate acceptable tolerance: Pinto, pink, red Mexican, cranberry, black, great northern and navy. Since not all types of dry beans of the species <i>Phaseolus vulgaris</i> have been tested, first application should be limited to a small area or consult your seed supplier or Albaugh LLC Representative for more information.
-----------------	---

**Dakota + Lontrel SG Herbicide in Imazamox Tolerant Canola (e.g., Canola varieties with the Imazamox Tolerant trait)**

**Dakota may be tank-mixed with Lontrel SG herbicide in Imazamox Tolerant Canola (e.g., Canola varieties with the Imazamox Tolerant trait)**

<b>Timing</b>	Early post-emergence		
<b>Rate</b>	29 g/ha Dakota + 100 g/ha Lontrel SG Herbicide		
<b>Water Volume</b>	100 L/ha		
<b>Surfactant/Adjuvant</b>	Add MERGE Adjuvant – 0.5% v/v		
<b>Weeds Controlled</b>	This tank-mix combination will provide control of weeds as listed on the Dakota label. Refer to Broadleaf weed and Grass control Section. Improved weed control with the tank mix vs. Dakota alone.		
	<b>Control</b>		<b>Top Growth Control</b>
	Wild buckwheat		Canada thistle
	Annual sow thistle		Perennial sow thistle
<b>Remark</b>	<p>Application should be made from the 2 - 6 leaf stage of the canola crop and after the weeds have emerged.</p> <p>To control broadleaf weeds, apply Dakota from the cotyledon to the 4-leaf stage. To control grasses, apply Dakota from the 1 - 4 true leaf stage until early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.</p> <p>For tank mixing, see mixing instructions.</p> <p>Refer to Tank mix rotational crop restrictions for details.</p>		

**Tank Mixing**

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Albaugh LLC at 1-800-247-8013 for information before applying any tank mix that is not specifically recommended on this label.



**Tank Mix Rotational Crop Restrictions****Dakota + Lontrel SG Herbicide in imazamox-tolerant Canola**

Fields previously treated with tank mix can be seeded to wheat, barley, oats and rye (not underseeded to forage legumes, clover or alfalfa), canola, corn, field peas, flax, forage grasses or can be summer-fallowed. Seed only those crops listed above in the year following treatment.

Do not seed to field peas for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field peas grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application, delay seeding field peas an additional 12 months (total 22 months following application).

Contact your local Albaugh LLC representative or retailer for more information before seeding field peas following drought conditions in the previous year.

**Tank Mix Pre-harvest / Grazing Interval**

Do not harvest the treated crop within 60 days of application.

**BROADLEAF WEED and GRASS CONTROL (Eastern Canada Only - Soybeans)**

Application must be made after the weeds have emerged. To control broadleaf weeds, apply Dakota from the cotyledon to 4 leaf stage. To control grasses, apply Dakota to green and yellow foxtail when in the 1-4 true leaf stage and to barnyard grass and fall panicum when in the 3-6 leaf stage.

Dakota at 36 g product/ha will control:

Common ragweed\*  
 Lady's thumb  
 Lamb's-quarters\*  
 Redroot pigweed  
 Velvetleaf  
 Wild mustard  
 Cocklebur\*  
 Eastern black nightshade  
 Green foxtail  
 Yellow foxtail  
 Barnyard grass  
 Fall panicum

\*Suppression only

Apply in 200 L/ha of water with a NON-IONIC SURFACTANT (AGRAL®, AGSURF® or SURF® 92) at the rate of 0.25% percent v/v and a fertilizer solution, 28-0-0 or 32-0-0, at a rate of 2 L/ha.

**MIXING INSTRUCTIONS**

1. Use 100 L/ha of water for Western Canada (the Prairie Provinces and Peace River Region of British Columbia) uses and 200 L/ha of water for Eastern Canada uses.
2. Use a 50-mesh (or coarser) filter screen.
3. Fill the spray tank with three-quarters of the required amount of clean water, start agitation and continue agitation throughout the entire mixing and spraying procedure.
4. Add the required amount of Dakota soluble bag(s) directly into the sprayer through the tank opening. For Dakota + Basagran Forte Herbicide tank mix, add Basagran Forte

Herbicide followed by recommended amount of nitrogen source (UAN 28% or ammonium sulphate).

5. For Dakota + Lontrel SG Herbicide tank mix, add recommended Lontrel SG Herbicide.
6. Agitate for at least ten minutes to dissolve the herbicide. After the herbicide is dissolved, continue agitation.
7. If required, add the required amount of MERGE Adjuvant. If excess foaming occurs, a silicone anti-foaming agent may be added (e.g. HALT®). **DO NOT** use MERGE Adjuvant for Dakota + Basagran Forte Herbicide tank mix.
8. Complete filling the tank to the desired level with water. If agitation is stopped for more than 5 minutes, re-suspend spray solution by full agitation prior to commencing spraying again.
9. Between loads of Dakota, check in-line and nozzle screens and rinse clean if necessary.
10. Upon completion of spraying Dakota, thoroughly flush tank, boom, hoses and in-line and nozzle screens with clean water to avoid possible injury to other crops.

## **FOLLOW CROPPING**

### **Prairie Provinces and Peace River Region of British Columbia**

Winter wheat can be planted 3 months after treatment (3 MAT) as a rotational crop.

Initial crop injury to canola may be observed. Avoid spray overlap as yield reduction may result. The following crops may be grown safely the year following an application:

Canary seed  
 Field corn  
 Field peas  
 Imazamox tolerant Canola  
 Canola  
 Lentils  
 Spring wheat  
 Durum wheat  
 Spring barley  
 Sunflower  
 Tame oats  
 Flax  
 Chickpeas  
 Imazamox tolerant Sunflowers

The following crop may be grown safely two years following an application:

Mustard (condiment type only)

There are insufficient data for other follow crops. Conduct a field bioassay (a test strip grown to maturity) the year before growing any crop other than those listed above.

### **Eastern Canada**

The following crops may be grown safely the year following an application:

Soybeans

Imazethapyr-tolerant canola  
 Field corn  
 Spring barley  
 Oats

Winter wheat (Winter wheat should not be planted within 100 days of the last DAKOTA Herbicide application).

Processing tomatoes (transplanted)

There are insufficient data for other follow crops. Conduct a field bioassay (a test strip grown to maturity) the year before growing any crop other than those listed above.

In cases of crop failure, only soybeans may be re-planted in the year of application.

### **RESISTANCE MANAGEMENT RECOMMENDATIONS**

For resistance management, Dakota is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to Dakota and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Dakota or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Albaugh LLC at 1-800-247-8013.

MERGE is a registered trade-mark of BASF Canada Inc.

®All other products listed are registered trademarks of their respective companies.

Outer Foil Pouch Label

GROUP	2	HERBICIDE
-------	---	-----------

Dakota™

Selective Herbicide

Water Dispersible Granular Herbicide in water-soluble bags

COMMERCIAL (AGRICULTURAL)

FOR SALE FOR USE IN EASTERN CANADA, THE PRAIRIE PROVINCES AND INTERIOR  
OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY

ACTIVE INGREDIENT: Imazamox ..... 70%

REGISTRATION NO. 34698      PEST CONTROL PRODUCTS ACT

NET CONTENTS: 470 g (4 x 117.5 g water-soluble bags)

Warning, contains the allergen milk

READ THE LABEL AND THIS BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN  
DO NOT SELL INDIVIDUAL WATER-SOLUBLE BAGS SEPARATELY

Albaugh LLC  
1525 NE 36<sup>th</sup> Street,  
Ankeny, Iowa  
United States  
1-800-247-8013

**PRECAUTIONS****KEEP OUT OF REACH OF CHILDREN.**

DO NOT APPLY BY AIR.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

**ENVIRONMENTAL PRECAUTIONS**

TOXIC to non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**FIRST AID**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

## **TOXICOLOGICAL INFORMATION**

Treat symptomatically.

For medical or treatment information from exposure to this product, call 1-888-347-6732 (7 days/week, 24-hr).

For 24-hour chemical spill, leak, fire, exposure, or accident response information, call CHEMTREC toll free at 1-800-424-9300.

## **STORAGE**

Store this product away from food or feed.

Store above 5°C.

Store in a cool, dry place away from children, animals, or fertilizers.

Keep unused water-soluble bags in resealed original container.

Keep packages dry at all times.

Keep packages out of rain.

Keep from freezing.

## **DISPOSAL**

1. When all the water-soluble bags are used, make the empty outer package unsuitable for further use.
2. Dispose of the outer package in accordance with provincial requirements.
3. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

WSB Label

DAKOTA™  
HERBICIDE

70% Water Dispersible Granular Herbicide

COMMERCIAL (AGRICULTURAL)

NET CONTENTS: 117.5 g

ACTIVE INGREDIENT: Imazamox ... 70%

Warning, contains the allergen milk

KEEP OUT OF REACH OF CHILDREN