

2023-03-14  
2023-0128  
\*Container Label\*



## VIKING™ Bromoxynil Herbicide

GROUP	6	HERBICIDE
-------	---	-----------

EMULSIFIABLE CONCENTRATE

COMMERCIAL (AGRICULTURAL)

For use in WHEAT (spring, durum & winter), BARLEY, OATS, FLAX (including low linolenic acid varieties), CORN, FALL RYE, CANARY SEED (for seed production), TRITICALE, GARLIC, CARROTS, SEEDLING ALFALFA, ESTABLISHED ALFALFA (for seed production only), and SEEDLING GRASSES, FORAGE and GRAIN SORGHUM, FORAGE MILLET, GRAIN PEARL MILLET as a pre-seed/pre-plant application prior to planting CANOLA and ZERO TILL

**REGISTRATION NO.** 34746 PEST CONTROL PRODUCTS ACT

**ACTIVE INGREDIENT:** BROMOXYNIL (present as an ester) .....280 g/L

**NET CONTENTS:** 1 L – 1020 L

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING**

**WARNING**



**POISON**

**In case of emergency call the 24 hour Emergency Response Number 1-800-424-9300 or 1-703-527-3887**

**Viking Crop Production Partners Inc.**

Box 18033  
RPO Shawnessy  
Calgary, AB. T2Y 0K3  
1-587-743-1029

Viking (TM) is a trademark of Viking Crop Production Partners Inc.

## PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes and clothing. Wash concentrate from skin or eyes immediately. Avoid breathing spray mist. After use wash hands and other exposed skin. Wear a chemical-resistant apron during mixing/loading activities.

Do not use in residential areas, which are defined as sites where bystanders may be present during or after spraying, including homes, schools, parks, playgrounds, playing fields, and public buildings.

Apply only to agricultural crops when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools, and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

**Ground applications:** Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, clean-up and repair. Wear a long-sleeved shirt and long pants, chemical resistant gloves, and socks plus shoes during applications. Gloves are not required during application within a closed cab.

**Aerial applications:** the field crew and the mixer/loaders: Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, clean-up and repair.

**Aerial applicators:** Wear a long-sleeved shirt, long pants, socks and shoes during application. Gloves are not required during application within a closed cockpit. Do not allow the pilot to mix chemicals to be loaded onto the aircraft.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry intervals (REIs) specified in the following table:

CROP	POST-APPLICATION ACTIVITY	REI
Garlic	Irrigation, handset	2 days
	All other activities	24 hours
Sweet Corn	Irrigation, handset	5 days
	Harvest, hand	20 days
	All other activities	24 hours

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours for all other crops.

**PRE-HARVEST GRAZING INTERVALS:** (wheat, barley, oats and seedling alfalfa) - Do not use treated crops for grazing of livestock or green feed until 30 days after application of **VIKING Bromoxynil Herbicide** unless otherwise stated on the label. Do not cut treated crops for forage until 30 days after application of **VIKING Bromoxynil Herbicide**.

**CAUTION:** Do not graze other treated crops or cut for feed unless specified above; sufficient data are not available to support such use.

## ENVIRONMENTAL PRECAUTIONS:

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of ponds, streams, rivers and other water sources. Avoid contamination of aquatic systems during application. DO NOT contaminate these systems through direct application, disposal of waste or cleaning equipment.

**FIRST AID:**

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**If swallowed:**

Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**If in eyes:**

Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**If on skin or clothing:**

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:**

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**TOXICOLOGICAL INFORMATION:**

This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat symptomatically.

**STORAGE:**

Store this product away from food or feed. Store the container tightly closed, away from seeds, fertilizer, plants and foodstuffs. This **VIKING Bromoxynil Herbicide** formulation will solidify at temperatures below - 20°C but will become useable again at temperatures above 0°C. Insecticides and fungicides should be segregated from herbicides so as to prevent the possibility of cross-contamination. SHAKE WELL BEFORE USING.

**RECYCLABLE CONTAINER DISPOSAL:**

Do not reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

**RETURNABLE CONTAINER DISPOSAL:**

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**REFILLABLE CONTAINER DISPOSAL:**

For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

**DISPOSAL OF UNUSED, UNWANTED PRODUCT:**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

**NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

\*Booklet Label\*



## VIKING™ Bromoxynil Herbicide

GROUP	6	HERBICIDE
-------	---	-----------

EMULSIFIABLE CONCENTRATE

COMMERCIAL (AGRICULTURAL)

For use in WHEAT (spring, durum & winter), BARLEY, OATS, FLAX (including low linolenic acid varieties), CORN, FALL RYE, CANARY SEED (for seed production), TRITICALE, GARLIC, CARROTS, SEEDLING ALFALFA, ESTABLISHED ALFALFA (for seed production only), and SEEDLING GRASSES, FORAGE and GRAIN SORGHUM, FORAGE MILLET, GRAIN PEARL MILLET as a pre-seed/pre-plant application prior to planting CANOLA and ZERO TILL

**REGISTRATION NO.** 34746 PEST CONTROL PRODUCTS ACT

**ACTIVE INGREDIENT:** BROMOXYNIL (present as an ester) .....280 g/L

**NET CONTENTS:** 1 L – 1020 L

**READ THE LABEL AND BOOKLET BEFORE USING**

**WARNING**



**POISON**

**In case of emergency call the 24 hour Emergency Response Number 1-800-424-9300 or 1-703-527-3887**

**Viking Crop Production Partners Inc.**

Box 18033  
RPO Shawnessy  
Calgary, AB. T2Y 0K3  
1-587-743-1029

Viking (TM) is a trademark of Viking Crop Production Partners Inc.

# Contents

PRECAUTIONS:	2
ENVIRONMENTAL PRECAUTIONS:	1
FIRST AID:	2
TOXICOLOGICAL INFORMATION:	2
STORAGE:	2
RECYCLABLE CONTAINER DISPOSAL:	2
DIRECTIONS FOR USE:	3
FIELD SPRAYER APPLICATION:	3
AERIAL APPLICATION (wheat and barley only):	3
Aerial Use Precautions:	3
Aerial Application Operator Precautions:	4
Product Specific Precautions:	4
BUFFER ZONES:	4
WHEAT (SPRING, HARD AND WINTER) - no sowing of legumes	5
<b>VIKING Bromoxynil Herbicide</b> used alone	5
<b>VIKING Bromoxynil Herbicide</b> + MCPA	6
<b>VIKING Bromoxynil Herbicide</b> + 2,4-D	6
<b>VIKING Bromoxynil Herbicide</b> + ACHIEVE™ LIQUID	6
WHEAT (Spring & Durum – not underseeded to legumes)	8
<b>VIKING Bromoxynil Herbicide</b> + HORIZON® 240 EC Herbicide Tank-Mix (1)	Error! Bookmark not defined.
<b>VIKING Bromoxynil Herbicide</b> + EVEREST® 70 WDG	8
DIRECTIONS FOR USE: WINTER WHEAT (EASTERN CANADA ONLY)	8
<b>VIKING Bromoxynil Herbicide</b> + PEAK® 75WG	8
BARLEY (not underseeded to legumes)	9
<b>VIKING Bromoxynil Herbicide</b> used alone	9
<b>VIKING Bromoxynil Herbicide</b> +MCPA	9
<b>VIKING Bromoxynil Herbicide</b> + 2,4-D	10
<b>VIKING Bromoxynil Herbicide</b> + ACHIEVE LIQUID	10
OATS (not underseeded to legumes)	11
<b>VIKING Bromoxynil Herbicide</b> used alone	11
<b>VIKING Bromoxynil Herbicide</b> + MCPA	11
CORN (Field & Sweet)	12
<b>VIKING Bromoxynil Herbicide</b> used alone	12
<b>VIKING Bromoxynil Herbicide</b> + ATRAZINE	12
<b>VIKING Bromoxynil Herbicide</b> + LOW-RATE ATRAZINE	13
CORN (Field, Silage and Sweet)	13
<b>VIKING Bromoxynil Herbicide</b> +LOW-RATE ATRAZINE	13
FIELD CORN (DO NOT USE ON SWEET CORN)	14
<b>VIKING Bromoxynil Herbicide</b> + BANVEL® or BANVEL II	14
FIELD CORN (DO NOT USE ON SWEET CORN) EASTERN CANADA	14
<b>VIKING Bromoxynil Herbicide</b> + OPTION™ 2.25 OD + AATREX® 480	14
FIELD CORN ONLY – EASTERN CANADA	Error! Bookmark not defined.
<b>VIKING Bromoxynil Herbicide</b> + ULTIM® 75 DF + non-ionic surfactant	Error! Bookmark not defined.
<b>VIKING Bromoxynil Herbicide</b> + Atrazine + ULTIM 75 DF + non-ionic surfactant	Error! Bookmark not defined.
DIRECTIONS FOR USE: (FIELD CORN ONLY)	15
<b>VIKING Bromoxynil Herbicide</b> + Accent™ 75 DF + non-ionic surfactant	15
FALL RYE (not underseeded to legumes)	15
<b>VIKING Bromoxynil Herbicide</b> used alone	15
<b>VIKING Bromoxynil Herbicide</b> + MCPA	16
FLAX (including low linolenic acid varieties)	16
<b>VIKING Bromoxynil Herbicide</b> used alone	16
<b>VIKING Bromoxynil Herbicide</b> + MCPA or MCPA-K	17
CANARY SEED (for seed production)	18

<b>VIKING Bromoxynil Herbicide</b> used alone .....	18
<b>VIKING Bromoxynil Herbicide</b> + MCPA.....	18
TRITICALE .....	19
<b>VIKING Bromoxynil Herbicide</b> used alone .....	19
GARLIC .....	19
<b>VIKING Bromoxynil Herbicide</b> used alone .....	19
CARROT.....	20
<b>VIKING Bromoxynil Herbicide</b> Herbicide used alone .....	20
PROVINCES OF ALBERTA, SASKATCHEWAN, MANITOBA ONLY SEEDLING ALFALFA.....	20
<b>VIKING Bromoxynil Herbicide</b> used alone .....	20
PROVINCES OF ALBERTA, SASKATCHEWAN, MANITOBA ONLY ESTABLISHED ALFALFA.....	21
<b>VIKING Bromoxynil Herbicide</b> used alone .....	21
PRE-SEED/PRE-PLANT APPLICATION PRIOR TO SEEDING CANOLA .....	21
<b>VIKING Bromoxynil Herbicide</b> + glyphosate (present as potassium or isopropylamine salt, dimethylamine, or diammonium salt).....	21
MINIMUM OR ZERO TILL .....	21
<b>VIKING Bromoxynil Herbicide</b> + glyphosate (present as potassium or isopropylamine salt, dimethylamine, or diammonium salt).....	22
SEEDLING GRASSES (not underseeded to legumes) .....	23
<b>VIKING Bromoxynil Herbicide</b> used alone .....	23
<b>VIKING Bromoxynil Herbicide</b> + MCPA.....	23
FORAGE SORGHUM AND FORAGE MILLET.....	24
<b>VIKING Bromoxynil Herbicide</b> used alone .....	24
DIRECTIONS FOR USE: GRAIN SORGHUM and GRAIN PEARL MILLET.....	24
<b>VIKING Bromoxynil Herbicide</b> used alone .....	24
MIXING TABLE.....	25
RESISTANCE-MANAGEMENT RECOMMENDATIONS:.....	27

## PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes and clothing. Wash concentrate from skin or eyes immediately. Avoid breathing spray mist. After use wash hands and other exposed skin. Wear a chemical-resistant apron during mixing/loading activities.

Do not use in residential areas, which are defined as sites where bystanders may be present during or after spraying, including homes, schools, parks, playgrounds, playing fields, and public buildings.

Apply only to agricultural crops when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools, and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

**Ground applications:** Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, clean-up and repair. Wear a long-sleeved shirt and long pants, chemical resistant gloves, and socks plus shoes during applications. Gloves are not required during application within a closed cab.

**Aerial applications:** the field crew and the mixer/loaders: Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, clean-up and repair.

**Aerial applicators:** Wear a long-sleeved shirt, long pants, socks and shoes during application. Gloves are not required during application within a closed cockpit. Do not allow the pilot to mix chemicals to be loaded onto the aircraft.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry intervals (REIs) specified in the following table:

CROP	POST-APPLICATION ACTIVITY	REI
Garlic	Irrigation, handset	2 days
	All other activities	24 hours
Sweet Corn	Irrigation, handset	5 days
	Harvest, hand	20 days
	All other activities	24 hours

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours for all other crops.

**PRE-HARVEST GRAZING INTERVALS:** (wheat, barley, oats and seedling alfalfa) - Do not use treated crops for grazing of livestock or green feed until 30 days after application of **VIKING Bromoxynil Herbicide** unless otherwise stated on the label. Do not cut treated crops for forage until 30 days after application of **VIKING Bromoxynil Herbicide**.

**CAUTION:** Do not graze other treated crops or cut for feed unless specified above; sufficient data are not available to support such use.

## ENVIRONMENTAL PRECAUTIONS:

**TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.**

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.



This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of ponds, streams, rivers and other water sources. Avoid contamination of aquatic systems during application. DO NOT contaminate these systems through direct application, disposal of waste or cleaning equipment.

**FIRST AID:**

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**If swallowed:**

Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**If in eyes:**

Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**If on skin or clothing:**

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:**

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**TOXICOLOGICAL INFORMATION:**

This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat symptomatically.

**STORAGE:**

Store this product away from food or feed. Store the container tightly closed away from seeds, fertilizer, plants and foodstuffs. This **VIKING Bromoxynil Herbicide** formulation will solidify at temperatures below - 20°C but will become useable again at temperatures above 0°C. Insecticides and fungicides should be segregated from herbicides so as to prevent the possibility of cross-contamination. SHAKE WELL BEFORE USING.

**RECYCLABLE CONTAINER DISPOSAL:**

Do not reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

**RETURNABLE CONTAINER DISPOSAL:**

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**REFILLABLE CONTAINER DISPOSAL:**

For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

**DISPOSAL OF UNUSED, UNWANTED PRODUCT:**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of

spills.

**NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

**DIRECTIONS FOR USE:**

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

**TIMING:** For best results spray when weeds are in the seedling stage. Apply in good growing conditions. Application **must** be made before the crop shields the weeds.

**SPRAYER:** Wash the tank and clean all filters, screens and tips. Select nozzle tips to apply the recommended volume of water per hectare. Flood jet type tips are not recommended. Adjust boom height to ensure uniform coverage of weeds.

**MIXING:** Half fill the tank with clean water. Add the required amount of **VIKING Bromoxynil Herbicide** and agitate thoroughly. Fill the tank and agitate again before use.

**FIELD SPRAYER APPLICATION:**

- DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty.
- DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification.
- Boom height must be 60 cm or less above the crop or ground.
- Spray at 5 - 10 km/h.
- Ensure that all tips are in good condition and spraying the same volume.

**AERIAL APPLICATION (wheat and barley only):**

**Aerial application:** DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. To reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

**Use Precautions:**

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

**Operator Precautions:**

Wear a long-sleeved shirt, long pants, socks and shoes during application. Gloves are not required during application within a closed cockpit. Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield, long-sleeved shirt, long pants, socks and shoes during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

**Product Specific Precautions:**

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-587-743-1029 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

**Volume: VIKING Bromoxynil Herbicide** plus either MCPA or 2,4-D must be applied in volume rates of not *less than* 20 L/ha. For best results when there is a heavy crop canopy, or when the majority of the weeds are cow cockle, green or pale smartweed, hemp nettle, redroot pigweed or Canada thistle, a volume rate of 40 L/ha is recommended.

**SPRAY BUFFER ZONES:**

Use of the following spray methods or equipment DO NOT require a spray buffer zone: hand-held or backpack sprayer, inter-row hooded sprayer, spot treatment, soil drench and soil incorporation.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop		Spray Buffer Zones (metres) Required for the Protection of:				
			Freshwater Habitat of Depths:		Estuarine / Marine Habitats of Depths:		Terrestrial Habitat
			Less than 1m	Greater than 1m	Less than 1 m	Greater than 1 m	
Field Sprayer*	All crops		1	1	1	1	1
Aerial	Barley and wheat	Fixed wing	20	5	1	1	55
		Rotary wing	20	3	1	1	45

\* For field sprayer application, spray buffer zones can be reduced with the use of drift-reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled spray buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled spray buffer zone can be reduced by 30%.

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

### Tank Mixtures

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Viking Crop Production Partners Inc. at 1-587-743-1029 for information before applying any tank mix that is not specifically recommended on this label.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on all product labels.

### WHEAT (SPRING, HARD AND WINTER) - no sowing of legumes

Tank mix	VIKING Bromoxynil Herbicide used alone
<b>Spray Volume + pressure</b>	Apply in 100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	<p><b>Spring and durum wheat</b> may be treated from the 2-leaf until the early flag leaf stage.</p> <p><b>Winter wheat</b> may be treated from the 2 to 4 leaf stage in the fall or from the time growth begins to the early flag leaf stage in the spring.</p>
<b>Application rate</b>	<p>1.0 - 1.2 L/ha (The 8 L container treats 6.6-8 ha).</p> <p>If weeds are beyond the leaf stages indicated or under adverse growing conditions, use of the higher recommended rate will improve control.</p>
<b>WEEDS CONTROLLED</b>	<p><b><u>Seedlings up to 4-leaf stage</u></b>            Green smartweed, Pale smartweed, Lady's-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Pigweed*, Velvetleaf***, Bluebur and American nightshade.</p> <p><b><u>Seedlings up to 8-leaf stage</u></b>            Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb's-quarters.</p> <p>* In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better results.            ** Spray before plants are 5 cm high.            *** Spray before plants are 8 cm high.</p>

<b>Tank mix</b>	<b>VIKING Bromoxynil Herbicide + MCPA</b>
<b>Spray Volume + pressure</b>	Apply in 50 - 100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	<b>Spring wheat</b> may be treated from the 2-leaf until the early flag leaf stage. <b>Winter wheat</b> may be treated from the 2 to 4 leaf stage in the fall or from the time growth begins to the early flag leaf stage in the spring
<b>Application rate</b>	<b>Spring wheat - VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank-mixed with 275-550 g active ingredient MCPA per hectare where indicated. <b>Winter wheat - VIKING Bromoxynil Herbicide</b> at 1.0 L/ha can be tank-mixed with 275 g active ingredient MCPA per hectare. Refer to table at end of booklet for correct volume of MCPA (The 8 L container of <b>VIKING Bromoxynil Herbicide</b> treats 8 ha) <b>Note:</b> MCPA-K is preferred for hemp nettle although other formulations may be used. Add MCPA to the spray tank first, agitate, then add <b>VIKING Bromoxynil Herbicide</b> .
<b>WEEDS CONTROLLED</b>	As listed for <b>VIKING Bromoxynil Herbicide</b> used alone plus these additional weeds <b>Seedlings up to 4-leaf stage</b> Redroot pigweed, Flixweed, Shepherd's purse, Scentsless chamomile*, Volunteer sunflower, Volunteer rapeseed/canola**, Hemp nettle***, Night flowering catchfly, Canada thistle****, Perennial sow thistle****, Ball mustard. <b>Seedlings up to 8-leaf stage</b> Stinkweed, Common ragweed, Wild mustard, Wormseed mustard.  *Spring annuals only. Will not control overwintered weeds. **For high infestations tank-mix 550 g active ingredient MCPA per ha. ***Tank-mix 550 g active ingredient MCPA per ha. Plants beyond the 4-leaf stage are not likely to be controlled. Plants emerging after application, which is often the case on peat-type soils, will not be controlled. ****Top growth control.
<b>Tank mix</b>	<b>VIKING Bromoxynil Herbicide + 2,4-D</b>
<b>Spray Volume + pressure</b>	Apply in 100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	<b>Spring and winter wheat</b> may be treated from the 4-leaf until the early flag leaf stage. Application before the 4-leaf stage may injure the crop.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank-mixed with 275-420 g active ingredient 2,4-D per ha. Refer to the table at end of booklet for correct volume of 2,4-D. (The 8 L container of <b>VIKING Bromoxynil Herbicide</b> treats 8 ha). <b>Note:</b> 2,4-D ester is preferred although other formulations may be used. Add 2,4-D to the spray tank first, agitate and then add <b>VIKING Bromoxynil Herbicide</b> . Refer to the 2,4-D label for precautions and limitations.
<b>WEEDS CONTROLLED</b>	As listed for <b>VIKING Bromoxynil Herbicide</b> used alone plus these additional weeds <b>Seedlings up to 4-leaf stage</b> Redroot pigweed, Night-flowering catchfly, Flixweed*, Shepherd's purse, Volunteer sunflower, Ball mustard <b>Seedlings up to 8-leaf stage</b> Stinkweed*, Wild mustard  *The higher rate of 2,4-D is recommended for larger overwintered weeds (fall rosettes).

<b>Tank mix</b>	<b>VIKING Bromoxynil Herbicide + ACHIEVE LIQUID</b>
<b>Spray Volume</b>	Apply in 50-100 L water per hectare at a pressure of 275 kPa

<b>+ pressure</b>	
<b>Timing</b>	<b>Durum wheat, Spring wheat and Winter wheat</b> may be treated from the 2-leaf until the early flag leaf stage of growth.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank mixed with <b>ACHIEVE LIQUID</b> at 0.5 L/ha. When using this tank-mixture, add Turbocharge to the spray tank at a concentration of 0.5% v/v (i.e. 0.5 L of Turbocharge per 100 L of spray solution).  <b>Note: VIKING Bromoxynil Herbicide</b> and <b>ACHIEVE LIQUID</b> tank-mixtures can be used in all varieties of Spring Wheat (including Canada Western red spring, amber durum, soft white spring, extra strong and Canada prairie spring), and red winter wheat.
<b>WEEDS CONTROLLED</b>	As listed for <b>VIKING Bromoxynil Herbicide</b> used alone plus wild oats, volunteer oats, green foxtail (wild millet), yellow foxtail, barnyard grass and Persian darnel.

**WHEAT (Spring – not underseeded to legumes)**

**For use in Manitoba, Saskatchewan, Alberta plus the Interior of British Columbia**

<b>Tank mix</b>	<b>VIKING Bromoxynil Herbicide + EVEREST 70 WDG</b>
<b>Spray Volume + pressure</b>	Apply in 50 – 100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	<b>Spring wheat</b> may be treated from the 1- leaf to 4-leaves on main stem, plus 2 tillers.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank mixed with EVEREST 70 WDG at 43 g/ha plus AGRAL® 90 or AG-SURF® adjuvant at a concentration of 0.25% v/v.  <b>Note:</b> <b>VIKING Bromoxynil Herbicide</b> and EVEREST 70 WDG tank mix can be used in all varieties of Spring wheat (including hard red spring, Canada prairie spring, soft white spring, and extra strong (utility) wheat).
<b>WEEDS CONTROLLED (Spring wheat only)</b>	As listed for <b>VIKING Bromoxynil Herbicide</b> used alone plus wild oats (including ACCase and triallate resistant biotypes) and green foxtail (including ACCase and dinitroaniline resistant biotypes).

**DIRECTIONS FOR USE: WINTER WHEAT (EASTERN CANADA ONLY)**

**VIKING Bromoxynil Herbicide + PEAK 75WG**

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Viking Crop Production Partners Inc. under the User Requested Minor Use Label Expansion program. For these uses, Viking Crop Production Partners Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

<b>Tank mix</b>	<b>VIKING Bromoxynil Herbicide + PEAK 75WG</b>
<b>Spray Volume+ pressure</b>	Apply in a minimum of 150 L water/ha at a pressure of 275 kPa.
<b>Timing</b>	Apply PEAK 75WG Herbicide as a broadcast spray to wheat up to and including BBCH Growth Stage 32 (beginning of stem elongation), by ground application only. Make one application per year. Refer to the PEAK 75WG label for further limitations and precautions.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 0.5 L/ha tank-mixed with PEAK 75WG at 13.3 g/ha plus a non-ionic surfactant at 0.2% v/v.  <b>Note:</b> PEAK 75WG Herbicide must be tank mixed with a non- ionic surfactant (AGRAL 90, Agsurf, or Citowett® Plus). <b>Do not use Crop Oil Concentrate for use on winter wheat.</b>
<b>Weeds controlled</b>	<b>Wild carrot in the 2 to 20 leaf stage</b>

**BARLEY (not underseeded to legumes)**

Tank mix	<b>VIKING Bromoxynil Herbicide used alone</b>
<b>Spray Volume + pressure</b>	Apply in 100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	The crop may be treated from the 2-leaf until the early flag leaf stage.
<b>Application rate</b>	1.0 - 1.2 L/ha (The 8 L container of <b>VIKING Bromoxynil Herbicide</b> treats 6.6-8 ha). If weeds are beyond the leaf stages indicated or under adverse growing conditions, use of the higher recommended rate will improve control.
<b>WEEDS CONTROLLED</b>	<p><b><u>Seedlings up to 4-leaf stage</u></b> Green smartweed, Pale smartweed, Lady's-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Pigweed*, Velvetleaf***, Bluebur, American nightshade.</p> <p><b><u>Seedlings up to 8-leaf stage</u></b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb's-quarters.</p> <p><i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better results.</i> <i>**Spray before plants are 5 cm high.</i> <i>***Spray before plants are 8 cm high.</i></p>

Tank mix	<b>VIKING Bromoxynil Herbicide + MCPA</b>
<b>Spray Volume + pressure</b>	Apply in 50 - 100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	The crop may be treated from the 2-leaf until the early flag leaf stage.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank-mixed with 275-550 g active ingredient MCPA per hectare where indicated. Refer to the table at end of booklet for correct volume of MCPA. (The 8 L container of <b>VIKING Bromoxynil Herbicide</b> treats 8 ha). <b>Note:</b> MCPA-K is preferred for hemp nettle although other formulations may be used. Add MCPA to the spray tank first, agitate, then add <b>VIKING Bromoxynil Herbicide</b> .
<b>WEEDS CONTROLLED</b>	<p>As listed for <b>VIKING Bromoxynil Herbicide</b> used alone <b>plus</b> these additional weeds</p> <p><b><u>Seedlings up to 4-leaf stage</u></b> Redroot pigweed, Flixweed, Shepherd's purse, Scentless chamomile*, Volunteer sunflower, Volunteer rapeseed/canola**, Hemp nettle***, Night flowering catchfly, Canada thistle****, Perennial sow thistle****, Ball mustard.</p> <p><b><u>Seedlings up to 8-leaf stage</u></b> Stinkweed, Common ragweed, Wild mustard, Wormseed mustard.</p> <p><i>*Spring annuals only. Will not control overwintered weeds.</i> <i>**For high infestations tank-mix 550 g active ingredient MCPA per ha.</i> <i>***Tank-mix 550 g active ingredient MCPA per ha. Plants beyond the 4-leaf stage are not likely to be controlled. Plants emerging after application, which is often the case on peat-type soils, will not be controlled.</i> <i>****Top growth control.</i></p>



Tank mix	VIKING Bromoxynil Herbicide + 2,4-D
<b>Spray Volume + pressure</b>	Apply in 100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	The crop may be treated from the 4-leaf until the early flag leaf stage. Application before the 4leaf stage may result in injury to the crop.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank-mixed with 275-420 g active ingredient 2,4-D per ha. Refer to table at end of booklet for correct volume of 2,4-D. (The 8 L container of <b>VIKING Bromoxynil Herbicide</b> treats 8 ha). <b>Note:</b> 2,4-D ester is preferred although other formulations may be used. Add 2,4-D to the spray tank first, agitate and then add <b>VIKING Bromoxynil Herbicide</b> . Refer to the 2,4-D label for precautions and limitations.
<b>WEEDS CONTROLLED</b>	As listed for <b>VIKING Bromoxynil Herbicide</b> used alone <b>plus</b> these additional weeds  <b>Seedlings up to 4-leaf stage</b> Redroot pigweed, Night-flowering catchfly, Flixweed*, Shepherd's purse, Volunteer sunflower, Ball mustard  <b>Seedlings up to 8-leaf stage</b> Stinkweed*, Wild mustard * The higher rate of 2,4-D is recommended for larger over wintered weeds (fall rosettes).

Tank mix	VIKING Bromoxynil Herbicide + ACHIEVE LIQUID
<b>Spray Volume + pressure</b>	Apply in 50-100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	Barley may be treated from the 2-leaf until the early flag leaf stage of growth.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank mixed with ACHIEVE LIQUID at 0.5 L/ha. When using this tank-mixture, add Turbocharge to the spray tank at a concentration of 0.5% v/v (i.e. 0.5 L of Turbocharge per 100 L of spray solution). <b>Note:</b> <b>VIKING Bromoxynil Herbicide</b> and ACHIEVE LIQUID tank-mixtures can be used in all 2 or 6 row varieties of barley (malting and feed varieties).
<b>WEEDS CONTROLLED</b>	As listed for <b>VIKING Bromoxynil Herbicide</b> used alone plus wild oats, volunteer oats, green foxtail (wild millet), yellow foxtail, barnyard grass and Persian darnel.

**OATS (not underseeded to legumes)**

Tank mix	<b>VIKING Bromoxynil Herbicide used alone</b>
<b>Spray Volume + pressure</b>	Apply in 100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	The crop may be treated from the 2-leaf until the early flag leaf stage.
<b>Application rate</b>	1.0 - 1.2 L/ha (The 8 L container treats 6.6-8 ha). If weeds are beyond the leaf stages indicated or under adverse growing conditions, use of the higher recommended rate will improve control.
<b>WEEDS CONTROLLED</b>	<p><b>Seedlings up to 4-leaf stage</b> Green smartweed, Pale smartweed, Lady's-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Pigweed*, Velvetleaf***, Bluebur, American nightshade.</p> <p><b>Seedlings up to 8-leaf stage</b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb's-quarters.</p> <p><i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better results.</i>  <i>**Spray before plants are 5 cm high.</i>  <i>***Spray before plants are 8 cm high.</i></p>

Tank mix	<b>VIKING Bromoxynil Herbicide + MCPA</b>
<b>Spray Volume + pressure</b>	Apply in 50 - 100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	The crop may be treated from the 2-leaf until the early flag leaf stage.
<b>Application rate</b>	<p><b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank-mixed with 275-550 g active ingredient MCPA per hectare where indicated. Refer to the table at end of booklet for correct volume of MCPA. (The 8 L container of <b>VIKING Bromoxynil Herbicide</b> treats 8 ha).  <b>Note:</b> MCPA-K is preferred for hemp nettle although other formulations may be used..            Add MCPA to the spray tank first, agitate, then add <b>VIKING Bromoxynil Herbicide</b>.</p>
<b>WEEDS CONTROLLED</b>	<p>As listed for <b>VIKING Bromoxynil Herbicide</b> used alone <b>plus</b> these additional weeds.</p> <p><b>Seedlings up to 4-leaf stage</b> Redroot pigweed, Flixweed, Shepherd's purse, Scentless chamomile*, Volunteer sunflower, Volunteer rapeseed/canola**, Hemp nettle***, Night flowering catchfly, Canada thistle****, Perennial sow thistle****, Ball mustard.</p> <p><b>Seedlings up to 8-leaf stage</b> Stinkweed, Common ragweed, Wild mustard, Wormseed mustard.</p> <p><i>*Spring annuals only. Will not control overwintered weeds.</i>  <i>**For high infestations tank-mix 550 g active ingredient MCPA per ha.</i>  <i>***Tank-mix 550 g active ingredient MCPA per ha. Plants beyond the 4-leaf stage are not likely to be controlled. Plants emerging after application, which is often the case on peat-type soils, will not be controlled.</i>  <i>****Top growth control.</i></p>

**CORN (Field & Sweet)**

Tank mix	<b>VIKING Bromoxynil Herbicide used alone</b>
<b>Spray Volume + pressure</b>	Apply in 200 - 300 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	Corn may be treated with a broadcast post-emergence application at the recommended rate from the 4-leaf stage onward. To ensure adequate coverage of weeds, drop pipes should be used when corn is beyond the 8-leaf stage or for a second application for later germinating weeds such as cocklebur and velvetleaf. Minimum re-treatment interval for the second application is 21 days. The pre-harvest interval (PHI) is 20 days.
<b>Application rate</b>	1.0 - 1.2 L/ha (The 8 L container treats 6.6-8 ha). Use of the higher recommended rate will improve control when heavy infestations of weeds are present.  <b>VIKING Bromoxynil Herbicide</b> is a contact herbicide thus good coverage of the weeds is essential.  <b>Note:</b> Temporary crop injury in the form of leaf scorching may occur in adverse growing conditions (especially if applied during or after periods of cool and wet, or hot and humid weather conditions). <b>DO NOT ADD OIL OR SURFACTANT.</b>
<b>WEEDS CONTROLLED</b>	<b>Seedlings up to 4-leaf stage</b> Green smartweed, Pale smartweed, Lady's-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Pigweed*, Velvetleaf***, Bluebur, American nightshade.  <b>Seedlings up to 8-leaf stage</b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb's-quarters. <i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better results.</i> <i>**Spray before plants are 5 cm high.</i> <i>***Spray before plants are 8 cm high. Atrazine tank-mix will improve control of larger plants.</i>

Tank mix	<b>VIKING Bromoxynil Herbicide + ATRAZINE</b>
<b>Spray Volume + pressure</b>	Apply in 200 - 300 L water per hectare at a pressure of 275 kPa.
<b>Timing</b>	Corn should be treated from the 4 to 8 leaf stage. Refer to the Atrazine label for limitations on grassy weeds.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank-mixed with Atrazine at 1.1 to 1.5 kg active ingredient per hectare. Add Atrazine to the spray tank first, agitate well, and then add <b>VIKING Bromoxynil Herbicide</b> . (The 8 L container of <b>VIKING Bromoxynil Herbicide</b> treats 8 ha).  <b>Note:</b> Temporary crop injury in the form of leaf scorching may occur in adverse growing conditions (especially if applied during or after periods of cool and wet, or hot and humid weather conditions). <b>DO NOT ADD OIL OR SURFACTANT, OR USE ATRAZINE FORMULATIONS CONTAINING OIL.</b>
<b>WEEDS CONTROLLED</b>	As listed for <b>VIKING Bromoxynil Herbicide</b> used alone plus weeds as indicated on the Atrazine label and, when Atrazine is applied at 1.5 kg active ingredient per hectare, bur cucumber seedlings in the 1 to 4-leaf stage.

Tank mix	<b>VIKING Bromoxynil Herbicide + LOW-RATE ATRAZINE</b>
<b>Spray Volume + pressure</b>	Apply in 200 - 300 L water per hectare at a pressure of 275 kPa.
<b>Timing</b>	Corn may be treated with a broadcast post-emergence application at the recommended rate from the 4-leaf stage onward. To ensure adequate coverage of weeds, drop pipes should be used when corn is beyond the 8-leaf stage or for a second application for later germinating weeds such as cocklebur and velvetleaf. Minimum re-treatment interval for the second application is 21 days. The pre-harvest interval (PHI) is 20 days.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 - 1.2 L/ha tank mixed with Atrazine at 0.5 kg active ingredient per hectare. Add Atrazine to the spray tank first, agitate well and then add <b>VIKING Bromoxynil Herbicide</b> . (The 8L container of <b>VIKING Bromoxynil Herbicide</b> treats 8 ha).  <b>Note:</b> Temporary crop injury in the form of leaf scorching may occur in adverse growing conditions (especially if applied during or after periods of cool and wet, or hot and humid weather conditions). DO NOT ADD OIL OR SURFACTANT, OR USE ATRAZINE FORMULATIONS CONTAINING OIL
<b>WEEDS CONTROLLED</b>	ted for <b>VIKING Bromoxynil Herbicide</b> used alone plus: <b>Seedlings up to the 6-leaf stage:</b> Redroot pigweed (triazine susceptible), velvetleaf* <b>Seedlings up to the 8-leaf stage:</b> Common ragweed  *Spray before weeds reach 10 cm in height.

#### **CORN (Field, Silage and Sweet)**

#### **DIRECTIONS FOR USE: VIKING Bromoxynil Herbicide + AATREX LIQUID 480**

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Viking Crop Production Partners Inc. under the User Requested Minor Use Label Expansion program. For these uses, Viking Crop Production Partners Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

Tank mix	<b>VIKING Bromoxynil Herbicide + AATREX LIQUID 480</b>
<b>Spray Volume + pressure</b>	Apply in 200 L water/ha at a pressure of 275 kPa.
<b>Timing</b>	Apply when the crop is at the 4- to 8-leaf stage, by ground application only. Make one application per year. Refer to the AATREX LIQUID 480 label for further limitations and precautions. Do not use in the province of British Columbia.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank-mixed with AATREX LIQUID 480 at 3.1 L/ha.  <b>Note:</b> Temporary crop injury in the form of leaf scorching may occur in adverse growing conditions (especially if applied during or after periods of cool and wet, or hot and humid weather conditions). DO NOT ADD OIL OR SURFACTANT, OR USE ATRAZINE FORMULATIONS CONTAINING OIL.
<b>WEEDS CONTROLLED</b>	<b>Seedlings at the unifoliolate to fourth trifoliolate leaf stage</b> Volunteer adzuki beans ( <i>Vigna angularis</i> ) <b>Seedlings up to the 10-leaf stage</b> Giant ragweed*

* <i>Suppression only</i>
---------------------------

**FIELD CORN (DO NOT USE ON SWEET CORN)**

Tank mix	<b>VIKING Bromoxynil Herbicide + BANVEL OR BANVEL II</b>
<b>Spray Volume + pressure</b>	Apply in 200 - 300 L water per hectare at a pressure of 275 kPa.
<b>Timing</b>	Field corn should be treated from the 4 - 6 leaf stage as an overall broadcast treatment. Drop pipes should be used when applying to corn over the 6 leaf stage and up to 50 cm; direct drop pipe nozzles on the weeds beneath the corn leaves. Apply no later than 2 weeks prior to tassel emergence and do not apply to corn over 50 cm in height.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank- mixed with BANVEL, or BANVEL II at 290 mL/ha. Add <b>VIKING Bromoxynil Herbicide</b> to the spray tank first, agitate well and then add BANVEL or BANVEL II. <b>Note:</b> Temporary crop injury in the form of leaf scorching may occur in adverse growing conditions (especially if applied during or after periods of cool and wet, or hot and humid weather conditions). <b>DO NOT ADD OIL OR SURFACTANT.</b>
<b>WEEDS CONTROLLED</b>	As listed for <b>VIKING Bromoxynil Herbicide</b> used alone plus: <b><u>Seedlings up to the 6-leaf stage</u></b> Redroot pigweed and Common ragweed including triazine resistant strains.

**FIELD CORN (DO NOT USE ON SWEET CORN) EASTERN CANADA**

Tank mix	<b>VIKING Bromoxynil Herbicide + OPTION 2.25 OD + AATREX 480</b>
<b>Spray Volume + pressure</b>	Apply in a minimum of 150 L water per hectare at a pressure of 175 – 275 kPa.
<b>Timing</b>	Field corn should be treated from the 1 to 8 leaf stage, or 5-6 visible collars (the leaf is counted once the next leaf is visible in the whorl). Refer to the OPTION 2.25 OD and AATREX 480 labels for further limitations and precautions.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 0.5 L/ha tank- mixed with OPTION 2.25 OD at 1.56 L/ha plus AATREX 480 at 1.04 L/ha. Add OPTION 2.25 OD to the spray tank first and ensure the product is thoroughly mixed with water. Add correct amount of dry component next, if appropriate, followed by the correct amount of the liquid component(s).
<b>WEEDS CONTROLLED</b>	Barnyard grass, large crabgrass, fall panicum, bristly foxtail, green foxtail, yellow foxtail, proso millet, witchgrass, common chickweed, redroot pigweed, velvetleaf, wild mustard, wormseed mustard, eastern black nightshade, lamb's quarters, common ragweed, wild buckwheat. Refer to the OPTION 2.25 OD label for appropriate application timing relative to weed growth stages.

**DIRECTIONS FOR USE: (FIELD CORN ONLY)****VIKING Bromoxynil Herbicide + Accent 75 DF + non-ionic surfactant****PRAIRIE PROVINCES**

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Viking Crop Production Partners Inc. under the User Requested Minor Use Label Expansion program. For these uses, Viking Crop Production Partners Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

Tank mix	VIKING Bromoxynil Herbicide + Accent 75 DF + non-ionic surfactant
<b>Spray Volume + pressure</b>	Apply in a minimum of 100 L water per hectare at a pressure of 275 kPa by ground only. <b>DO NOT apply using aerial application equipment.</b>
<b>Timing</b>	Apply as a single post emergent spray. Apply post-emergence to corn when at the 4 to 8 leaf stage (2-6 visible collars). Do not apply prior to the 4-leaf (2 visible collars) or after the 8leaf (6 visible collars) stage of corn. Observe a PHI of 30 days. Refer to the Accent 75 DF label for further limitations and precautions.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank-mixed with Accent 75 DF at 33.4 grams/ha and a recommended non-ionic surfactant, either Citowett Plus, Agral 90 or Agsurf at 2 L per 1000 L spray solution (0.2% v/v).
<b>WEEDS CONTROLLED</b>	As listed for <b>VIKING Bromoxynil Herbicide</b> used alone <b>plus</b> these additional weeds:  <b><u>Seedlings in the 1 to 6-leaf stage (up to early tillering, 2 – two leaf tillers):</u></b> Yellow foxtail*, barnyard grass, fall panicum, green foxtail, old witchgrass  <b><u>Plants in the 3 to 6-leaf stage (10-20 cm in height – leaf extended):</u></b> Quack grass  * <i>Suppression only</i>

**FALL RYE (not underseeded to legumes)**

Tank mix	VIKING Bromoxynil Herbicide used alone
<b>Spray Volume + pressure</b>	Apply in 100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	Fall rye may be treated from the time growth begins to the early flag leaf stage in the spring.
<b>Application rate</b>	1.0 - 1.2 L/ha (The 8 L container treats 6.6-8 ha). If weeds are beyond the leaf stages indicated or under adverse growing conditions, use of the higher recommended rate will improve control.
<b>WEEDS CONTROLLED</b>	<b><u>Seedlings up to 4-leaf stage</u></b> Green smartweed, Pale smartweed, Lady's-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Pigweed*, Velvetleaf***, Bluebur, American nightshade.  <b><u>Seedlings up to 8-leaf stage</u></b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb's-quarters.  <i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better results. **Spray before plants are 5 cm high. ***Spray before plants are 8 cm high.</i>

Tank mix	<b>VIKING Bromoxynil Herbicide + MCPA</b>
<b>Spray Volume + pressure</b>	Apply in 50 - 100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	Fall rye may be treated from the time growth begins to the early flag leaf stage in the spring.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank- mixed with 275 g active ingredient MCPA per hectare where indicated. Refer to the mixing tables at end of book for correct volume of MCPA. (The 8 L container of <b>VIKING Bromoxynil Herbicide</b> treats 8 ha).  <b>Note:</b> MCPA ester is preferred although other formulations may be used. Add MCPA to the spray tank first, agitate, then add <b>VIKING Bromoxynil Herbicide</b> .
<b>WEEDS CONTROLLED</b>	As listed for <b>VIKING Bromoxynil Herbicide</b> used alone <b>plus</b> these additional weeds  <b>Seedlings up to 4-leaf stage</b> Redroot pigweed, Flixweed, Shepherd's purse, Scentless chamomile*, Volunteer sunflower, Volunteer rapeseed/canola**, Night flowering catchfly, Canada thistle**, Perennial sow thistle**, Ball mustard.  <b>Seedlings up to 8-leaf stage</b> Stinkweed, Common ragweed, Wild mustard, Wormseed mustard.  <i>*Spring annuals only. Will not control overwintered weeds.</i> <i>**Top growth control only.</i>

#### FLAX (including low linolenic acid varieties)

Tank mix	<b>VIKING Bromoxynil Herbicide used alone</b>
<b>Spray Volume + pressure</b>	Apply in 100 L water per hectare at a pressure of 275 kPa.
<b>Timing</b>	Flax may be treated from the time it is 5 cm high up to the early flower bud stage but for best results apply <b>VIKING Bromoxynil Herbicide</b> when flax is 5-10 cm high. <b>Note:</b> Spraying in the evening may reduce risk of flax injury. <b>Warning:</b> Do not spray unthrifty crops or when plants are under stress. Do not spray in periods of hot, humid weather. <b>DO NOT APPLY BY AIR.</b> Observe a minimum interval to harvest of 60 days after application of <b>VIKING Bromoxynil Herbicide</b> . Flax is less tolerant of this product than are the cereal crops. Some leaf burn and retarded growth may delay maturity 2-3 days. Avoid Using <b>VIKING Bromoxynil Herbicide</b> when flax is under stress or during hot (over 29°C) humid weather as severe crop injury may occur.
<b>Application rate</b>	1.0 L/ha
<b>WEEDS CONTROLLED</b>	<b>Seedlings up to 4-leaf stage</b> Green smartweed, Pale smartweed, Lady's-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, triazine-resistant Pigweed*, Velvetleaf***, Bluebur, American nightshade.  <b>Seedlings up to 8-leaf stage</b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb's-quarters.  <i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled.</i> <i>**Spray before plants are 5 cm high.</i> <i>***Spray before plants are 8 cm high.</i>

Tank mix	VIKING Bromoxynil Herbicide + MCPA or MCPA-K
<b>Spray Volume + pressure</b>	Apply in 100 L water per hectare at a pressure of 275 kPa.
<b>Timing</b>	<p>Flax may be treated from the time it is 5 cm high up to the early flower bud stage but for best results apply <b>VIKING Bromoxynil Herbicide</b> when flax is 5-10 cm high.</p> <p><b>Note:</b> Spraying in the evening may reduce risk of flax injury.</p> <p><b>Warning:</b> Do not spray unthrifty crops or when plants are under stress. Do not spray in periods of hot, humid weather. <b>DO NOT APPLY BY AIR.</b></p> <p>Observe a minimum interval to harvest of 60 days after application of <b>VIKING Bromoxynil Herbicide</b>.</p> <p>Flax is less tolerant of this product than are the cereal crops. Some leaf burn and retarded growth may delay maturity 2-3 days. Avoid Using <b>VIKING Bromoxynil Herbicide</b> when flax is under stress or during hot (over 29°C) humid weather as severe crop injury may occur.</p>
<b>Application rate</b>	<p><b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank mixed with MCPA or MCPA-K at 275 g a.i./ha.</p> <p>The amine, ester or K-salt formulations of MCPA can be used in tank mixtures.</p>
<b>WEEDS CONTROLLED</b>	<p>As listed for <b>VIKING Bromoxynil Herbicide</b> used alone <b>plus</b> these additional weeds</p> <p><b>Seedlings up to 4-leaf stage</b> Redroot pigweed, Flixweed, Shepherd's purse, Scentless chamomile*, Volunteer sunflower, Volunteer rapeseed/canola**, Hemp nettle***, Night flowering catchfly, Canada thistle****, Perennial sow thistle****, Ball mustard.</p> <p><b>Seedlings up to 8-leaf stage</b> Stinkweed, Common ragweed, Wild mustard, Wormseed mustard.</p> <p><i>*Spring annuals only. Will not control overwintered weeds.</i></p> <p><i>**For high infestations tank-mix 550 g active ingredient MCPA per ha.</i></p> <p><i>***Tank-mix 550 g active ingredient MCPA per ha. Plants beyond the 4-leaf stage are not likely to be controlled. Plants emerging after application, which is often the case on peat-type soils, will not be controlled.</i></p> <p><i>****Top growth control only.</i></p>



**CANARY SEED (for seed production)**

Tank mix	<b>VIKING Bromoxynil Herbicide used alone</b>
<b>Spray Volume + pressure</b>	Apply in 100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	The crop may be treated from the 3 to 5 leaf stage.
<b>Application rate</b>	1.0 L/ha (The 8 L container treats 8 ha).
<b>WEEDS CONTROLLED</b>	<p><b><u>Seedlings up to 4-leaf stage</u></b> Green smartweed, Pale smartweed, Lady's-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Pigweed*, Velvetleaf***, Bluebur, American nightshade.</p> <p><b><u>Seedlings up to 8-leaf stage</u></b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb's-quarters.</p> <p><i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled.</i> <i>**Spray before plants are 5 cm high.</i> <i>***Spray before plants are 8 cm high.</i></p>

Tank mix	<b>VIKING Bromoxynil Herbicide + MCPA</b>
<b>Spray Volume + pressure</b>	Apply in 50 - 100 L water per hectare at a pressure of 275 kPa
<b>Timing</b>	The crop may be treated from the 3 to 5 leaf stage.
<b>Application rate</b>	<p><b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank- mixed with 275 g active ingredient MCPA per hectare. Refer to the table at end of book for correct volume of MCPA. (The 8 L container of <b>VIKING Bromoxynil Herbicide</b> treats 8 ha). <b>Note:</b> MCPA ester is preferred although other formulations may be used. Add MCPA to the spray tank first, agitate, then add <b>VIKING Bromoxynil Herbicide</b>.</p>
<b>WEEDS CONTROLLED</b>	<p>As listed for <b>VIKING Bromoxynil Herbicide</b> used alone <b>plus</b> these additional weeds</p> <p><b><u>Seedlings up to 4-leaf stage</u></b> Redroot pigweed, Flixweed, Shepherd's purse, Scentless chamomile*, Volunteer sunflower, Volunteer rapeseed/canola**, Night flowering catchfly, Canada thistle**, Perennial sow thistle**, Ball mustard.</p> <p><b><u>Seedlings up to 8-leaf stage</u></b> Stinkweed, Common ragweed, Wild mustard, Wormseed mustard.</p> <p><i>*Spring annuals only. Will not control overwintered weeds.</i> <i>**Top growth control only.</i></p>

## TRITICALE

Tank mix	VIKING Bromoxynil Herbicide used alone
Spray Volume + pressure	Apply in 100 L water/ha at a pressure of 275 kPa.
Timing	The crop may be treated from the 2-leaf until the early flag leaf stage.
Application rate	1.0 - 1.2 L/ha (The 8L container treats 6.6-8 ha). If weeds are beyond the leaf stage indicated or under adverse growing conditions, use of the higher recommended rate will improve control.
WEEDS CONTROLLED	<p><b><u>Seedlings up to 4-leaf stage</u></b> Green smartweed, Pale smartweed, Lady's-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Pigweed*, Velvetleaf***, Bluebur, American nightshade.</p> <p><b><u>Seedlings up to 8-leaf stage</u></b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb's-quarters.</p> <p><i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better results.</i> **Spray before plants are 5 cm high. ***Spray before plants are 8 cm high</p>

## GARLIC

Tank mix	VIKING Bromoxynil Herbicide used alone
Spray Volume + pressure	Apply in 200 - 300 L water per hectare at a pressure of 275 kPa.
Timing	Apply early post emergent to weeds by ground application only. One application per year. Observe a PHI of 58 days.
Application rate	1.0 L/ha
WEEDS CONTROLLED	<p><b><u>Seedlings up to 4-leaf stage</u></b> Green smartweed, Pale smartweed, Lady's-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Pigweed*, Velvetleaf***, Bluebur, American nightshade.</p> <p><b><u>Seedlings up to 8-leaf stage</u></b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb's-quarters.</p> <p><i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better results.</i> **Spray before plants are 5 cm high. ***Spray before plants are 8 cm high.</p>

**CARROT**

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Viking Crop Production Partners Inc. under the User Requested Minor Use Label Expansion program. For these uses, Viking Crop Production Partners Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

Tank mix	VIKING Bromoxynil Herbicide used alone
<b>Spray Volume + pressure</b>	Apply in 200 - 300 L water per hectare at a pressure of 275 kPa.
<b>Timing</b>	Apply only to carrots grown on muck soils containing greater than 10% organic matter. Apply 3-4 days prior to crop emergence by ground application only. One application per year. Observe a PHI of 45 days. Do not irrigate carrots that have received a preemergence application of <b>VIKING Bromoxynil Herbicide</b> for 2 days following application or 3 days of crop emergence.
<b>Application rate</b>	1.0 - 1.2 L/ha <b>VIKING Bromoxynil Herbicide</b> is a contact herbicide thus good coverage of the weeds is essential.
<b>WEEDS CONTROLLED</b>	<p><b>Seedlings up to 4-leaf stage</b> Green smartweed, Pale smartweed, Lady's-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Pigweed*, Velvetleaf***, Bluebur, American nightshade.</p> <p><b>Seedlings up to 8-leaf stage</b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb's-quarters.</p> <p><i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better results.</i> **Spray before plants are 5 cm high. ***Spray before plants are 8 cm high.</p>

**PROVINCES OF ALBERTA, SASKATCHEWAN, MANITOBA ONLY SEEDLING ALFALFA**

Tank mix	VIKING Bromoxynil Herbicide used alone
<b>Spray Volume + pressure</b>	Apply in 100 L water/ha at a pressure of 275 kPa – ground application only.
<b>Timing</b>	Seedling alfalfa may be treated at the 2 to 6 trifoliolate leaf stage.
<b>Application rate</b>	1.0 L/ha (The 8L container treats 8 ha). <b>Note:</b> Use of this product may cause temporary leaf scorch or foliar burn in adverse growing conditions, especially if applied during or after periods of cool and wet, or hot and humid weather conditions. Do not apply if crop is under stress. Do not apply to alfalfa seedlings in the unifoliolate or 1st trifoliolate leaf stage. Avoid overlapping as severe crop injury may occur.
<b>WEEDS CONTROLLED</b>	<p><b>Seedlings up to 4-leaf stage:</b> Green smartweed, Pale smartweed, Lady's-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Velvetleaf***, Bluebur, American nightshade.</p> <p><b>Seedlings up to 8-leaf stage:</b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb's-quarters.</p> <p><i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better results.</i> **Spray before plants are 5 cm high. ***Spray before plants are 8 cm high.</p>

**PROVINCES OF ALBERTA, SASKATCHEWAN, MANITOBA ONLY ESTABLISHED ALFALFA  
(for seed production only)**

Tank mix	<b>VIKING Bromoxynil Herbicide used alone</b>
<b>Spray Volume + pressure</b>	Apply in 100 L water/ha at a pressure of 275 kPa – ground application only.
<b>Timing</b>	Established alfalfa may be treated until alfalfa is 25 cm tall. Maximum of 2 applications per year. Minimum re-treatment interval for the second application is 21 days
<b>Application rate</b>	1.0 - 1.2 L/ha (The 8L container treats 6.6-8 ha). If weeds are beyond the leaf stage indicated or under adverse growing conditions, use of the higher recommended rate will improve control.  <b>Note:</b> Use of this product may cause temporary leaf scorch or foliar burn in adverse growing conditions, especially if applied during or after periods of cool and wet, or hot and humid weather conditions. Crop will recover and yield will not be affected.
<b>WEEDS CONTROLLED</b>	<b>Seedlings up to 4-leaf stage</b> Green smartweed, Pale smartweed, Lady’s-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Velvetleaf***, Bluebur, American nightshade.  <b>Seedlings up to 8-leaf stage</b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb’s-quarters.  <i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better results.</i> <i>**Spray before plants are 5 cm high.</i> <i>***Spray before plants are 8 cm high.</i>

**PRE-SEED/PRE-PLANT APPLICATION PRIOR TO SEEDING CANOLA**

Tank mix	<b>VIKING Bromoxynil Herbicide + glyphosate (present as potassium or isopropylamine salt, dimethylamine, or diammonium salt)</b>
<b>Spray Volume + pressure</b>	Apply in 100 L of <b>clean</b> water per hectare at a pressure of 275 kPa.
<b>Timing</b>	Application can be made up to one day prior to seeding. <b>DO NOT</b> apply after seeding or crop emergence.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 – 1.25 L/ha tank-mixed with glyphosate at 448 – 1782 g a.e./ha. Under adverse growing conditions or heavy weed populations use of the higher recommended rate of <b>VIKING Bromoxynil Herbicide</b> will improve control.  <b>VIKING Bromoxynil Herbicide</b> should be added to the spray tank first, then add glyphosate.
<b>WEEDS CONTROLLED</b>	Annual and perennial weeds listed on the glyphosate label, volunteer canola* seedlings at the 1-4 leaf stage  <i>* Including herbicide resistant biotypes.</i>

**MINIMUM OR ZERO TILL**

<b>Tank mix</b>	<b>VIKING Bromoxynil Herbicide + glyphosate (present as potassium or isopropylamine salt, dimethylamine, or diammonium salt)</b>
<b>Spray Volume + pressure</b>	Apply in 100 L of <b>clean</b> water per hectare at a pressure of 275 kPa
<b>Timing</b>	This recommendation applies only to minimum or zero till situations <b>before</b> emergence of the crop. Application can be made just prior to or immediately after seeding. Under no circumstances should application be made after crop emergence. Weeds in the seedling stage are most susceptible. This treatment should only be used for minimum or zero-tillage cropping systems in wheat, barley and oats.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 – 1.25 L/ha tank-mixed with glyphosate at 396 g a.e./ha. Under adverse growing conditions or when high populations of volunteer canola are present, use of the higher recommended rate of <b>VIKING Bromoxynil Herbicide</b> will improve control. <b>VIKING Bromoxynil Herbicide</b> should be added to the spray tank first, then add glyphosate.
<b>WEEDS CONTROLLED</b>	<b><u>Seedlings up to 4-leaf stage</u></b> Green smartweed, Pale smartweed, Lady's-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Pigweed*, Velvetleaf***, Bluebur, American nightshade, Volunteer canola****. <b><u>Seedlings up to 8-leaf stage</u></b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb's-quarters. <b><u>Grassy weeds up to 15 cm</u></b> Wild oats, Volunteer cereals, Green foxtail  <i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled.</i> <i>**Spray before plants are 5 cm high,</i> <i>***Spray before plants are 8 cm high.</i> <i>****Including herbicide tolerant types.</i>

**SEEDLING GRASSES (not underseeded to legumes)**

<b>Tank mix</b>	<b>VIKING Bromoxynil Herbicide used alone</b>
<b>Spray Volume + pressure</b>	Apply in 150 L water per hectare at a pressure of 275 kPa.
<b>Timing</b>	Seedling grasses may be treated from the 2 to 4 leaf stage.
<b>Application rate</b>	1.0 - 1.2 L/ha (The 8 L container treats 6.6-8 ha). If weeds are beyond the leaf stage indicated or under adverse growing conditions, use of the higher recommended rate will improve control. <b>Note:</b> Grasses grown for seed production in the year of establishment only. <b>Crops:</b> Brome grass, Crested wheat grass, Intermediate wheat grass, Slender wheat grass, Tall wheat grass, Russian wild rye, Timothy, Orchard grass, creeping red fescue, Meadow fescue, Reed canary grass.
<b>WEEDS CONTROLLED</b>	<b>Seedlings up to 4-leaf stage</b> Green smartweed, Pale smartweed, Lady's-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Pigweed*, Velvetleaf***, Bluebur, American nightshade. <b>Seedlings up to 8-leaf stage</b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb's-quarters.  <i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better results.</i> <i>**Spray before plants are 5 cm high.</i> <i>***Spray before plants are 8 cm high.</i>

<b>Tank mix</b>	<b>VIKING Bromoxynil Herbicide + MCPA</b>
<b>Spray Volume + pressure</b>	Apply in 150 L water per hectare at a pressure of 275 kPa.
<b>Timing</b>	Seedling grasses may be treated from the 2 to 4 leaf stage.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha tank- mixed with 275 g active ingredient MCPA per hectare. Refer to the table at end of book for correct volume of MCPA. (The 8 L container of <b>VIKING Bromoxynil Herbicide</b> treats 8 ha). <b>Note:</b> MCPA ester is preferred although other formulations may be used. Add MCPA to the spray tank first, agitate, then add <b>VIKING Bromoxynil Herbicide</b> .
<b>WEEDS CONTROLLED</b>	As listed for <b>VIKING Bromoxynil Herbicide</b> used alone <b>plus</b> these additional weeds <b>Seedlings up to 4-leaf stage</b> Redroot pigweed, Flixweed, Shepherd's purse, Scentless chamomile*, Volunteer rapeseed/canola**, Volunteer sunflower, Night flowering catchfly, Canada thistle**, Perennial sow thistle**, Ball mustard. <b>Seedlings up to 8-leaf stage</b> Stinkweed, Common ragweed, Wild mustard, Wormseed mustard.  <i>*Spring annuals only. Will not control overwintered weeds.</i> <i>**Top growth control only.</i>

**FORAGE SORGHUM AND FORAGE MILLET**

<b>Tank mix</b>	<b>VIKING Bromoxynil Herbicide used alone</b>
<b>Spray Volume + pressure</b>	Apply in 200 – 300 L water per hectare at a pressure of 275 kPa.
<b>Timing</b>	Apply when the crop is at or beyond the 4 leaf stage and less than 20 cm in height by ground application only. One application per year. Observe a PHI of 30 days.
<b>Application rate</b>	1.0 L/ha
<b>WEEDS CONTROLLED</b>	<p><b><u>Seedlings up to 4-leaf stage</u></b> Green smartweed, Pale smartweed, Lady’s-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Pigweed*, Velvetleaf***, Bluebur, American nightshade.</p> <p><b><u>Seedlings up to 8-leaf stage</u></b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb’s-quarters.</p> <p><i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better results.</i>  <i>**Spray before plants are 5 cm high.</i>  <i>***Spray before plants are 8 cm high.</i></p>

**DIRECTIONS FOR USE: GRAIN SORGHUM and GRAIN PEARL MILLET**  
**VIKING Bromoxynil Herbicide**

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Viking Crop Production Partners Inc. under the User Requested Minor Use Label Expansion program. For these uses, Viking Crop Production Partners Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

<b>Tank mix</b>	<b>VIKING Bromoxynil Herbicide used alone</b>
<b>Spray Volume + pressure</b>	Apply in 200-300 L water/ha at a pressure of 275 kPa.
<b>Timing</b>	Apply when the crop is at or beyond the 4 leaf stage and less than 20 cm in height by ground application only. One application per year. Observe a PHI of 100 days.
<b>Application rate</b>	<b>VIKING Bromoxynil Herbicide</b> at 1.0 L/ha.
<b>WEEDS CONTROLLED</b>	<p><b><u>Seedlings up to 4-leaf stage</u></b> Green smartweed, Pale smartweed, Lady’s-thumb, Wild mustard*, Cow cockle*, Russian thistle**, Stinkweed*, Cocklebur, Common ragweed, Pigweed*, Velvetleaf***, Bluebur, American nightshade</p> <p><b><u>Seedlings up to 8-leaf stage</u></b> Wild buckwheat, Tartary buckwheat, Common buckwheat, Common groundsel, Lamb’s-quarters.</p> <p><i>*In normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better results.</i>  <i>**Spray before plants are 5 cm high.</i>  <i>***Spray before plants are 8 cm high.</i></p>

**MIXING TABLE**

For calculating the amount of product to add to the spray tank when applying 100 L spray solution per hectare.

		VIKING BROMOXYNIL HERBICIDE (280 g/L)		MCPA AMINE 500 or MCPA ESTER 500		MCPA-K 400		MCPA SODIUM SALT 300	
<b>Rate active per hectare</b>		280 g	340g	275g	550g	275g	550g	275g	550g
<b>Rate product per hectare</b>		1.0L	1.2L	550 mL	1.1L	690 mL	1.38 L	920 mL	1.83 L
<b>Size of Spray tank</b>		<b>Litres of Chemical to add to spray tank (approx.)</b>							
<b>Gallons</b>	<b>Liters</b>								
500	2273	22.7	27.3	12.5	25.0	15.7	31.4	20.9	41.6
400	1818	18.2	21.8	10.0	20.0	12.5	25.0	16.7	33.3
350	1591	16.0	19.1	8.75	17.5	11.0	22.0	14.6	29.2
300	1363	13.6	16.4	7.5	15.0	9.4	18.8	12.5	25.0
250	1136	11.4	13.6	6.25	12.5	7.8	15.7	10.5	20.8
200	909	9.1	10.9	5.0	10.0	6.25	12.5	8.4	16.6
100	455	4.5	5.5	2.5	5.0	3.2	6.3	4.2	8.4
50	227	2.3	2.7	1.25	2.5	1.6	3.2	2.1	4.2
22	100	1.0	1.2	550 mL	1.1	690 mL	1.38	920 mL	1.8
10	45	450 mL	540 mL	250 mL	500 mL	310 mL	620 mL	420 mL	820 mL

		VIKING BROMOXYNIL HERBICIDE (280 g/L)		2,4-D AMINE 500		2,4-D 600 LV ESTER		2,4-D 700 LV ESTER	
<b>Rate active per hectare</b>		280 g	340g	275g		275g		275g	
<b>Rate product per hectare</b>		1.0L	1.2 L	550 mL		458 mL		393 mL	
<b>Size of Spray Tank</b>		<b>Litres of chemical to add to the spray tank (approx.)</b>							
<b>Gallons</b>	<b>Liters</b>								
500	2273	22.7	27.3	12.5		10.4		8.9	
400	1818	18.2	21.8	10.0		8.3		7.1	
350	1591	16.0	19.1	8.75		7.3		6.25	
300	1363	13.6	16.4	7.5		6.25		5.4	
250	1136	11.4	13.6	6.25		5.2		4.5	
200	909	9.1	10.9	5.0		4.2		3.6	
100	455	4.5	5.5	2.5		2.1		1.8	
50	227	2.3	2.7	1.25		1.0		890 mL	
22	100	1.0	1.2	550 mL		458 mL		390 mL	
10	45	450 mL	540 mL	250 mL		210 mL		180 mL	





## RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, **VIKING Bromoxynil Herbicide** is a Group 6 herbicide. Any weed population may contain or develop plants naturally resistant to **VIKING Bromoxynil Herbicide** and other Group 6 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed. To delay herbicide resistance:

- Where possible rotate the use of **VIKING Bromoxynil Herbicide** or other Group 6 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your Viking Crop Production Partners Inc representative, or call Viking Crop Production Partners Inc. at 1-587-743-1029.

Viking (TM) is a trademark of Viking Crop Production Partners Inc.  
All other products mentioned are trademarks of their respective holders.