

86.03.24

DC-2000

DISINFECTANT - SANITIZER - DEODORIZER

Disinfectant - Sanitizer - Deodorizer for Hospitals,
Institutional, Industrial, School, Dairy,
and other Farm and Home use.

GUARANTEE:

ALKYL (C₁₂, 40%, C₁₄ 50%, C₁₆ 10%)
5% Dimethyl Benzyl Ammonium Chloride

READ THE LABEL BEFORE USING

CAUTION CORROSIVE

DIN 67617g

REGISTRATION NO. 19287
PEST CONTROL PRODUCTS ACT

Net Contents: 4 litres

Manufactured by: CHEMAC INDUSTRIES INC, #1-368 Industrial
Avenue, Kelowna, B.C. V1Y 7E8

Sanitizing of Food Processing Equipment and other Hard Surfaces
in Food Contact Locations For sanitizing food processing
equipment, dairy equipment, food utensils, dishes, silverware,
glasses, sink tops, countertops, refrigerated storage and display
equipment and other surfaces. Wash and rinse all articles
thoroughly, then apply a solution of 4 mL DC-2000 in 1 litre of
water. (1 part in 250 parts of water) Apply to sink tops,
countertops, refrigerated storage, display equipment and other
stationary hard surfaces by cloth or brush. Dishes, silverware,
glasses, cooking utensils and other similar size food processing
equipment can be sanitized by immersion in a 4 mL per litre
dilution of DC-2000. No potable water rinse is required. At
4 mL/litre DC-2000 has a hard water tolerance of 500 ppm of
hardness calculated as CaCO₃ when evaluated by the AOAC
Germicidal and Detergent Sanitizer Method against Escherichia
coli and Staphylococcus aureus.

The udders, flanks, and teats of dairy cows can be sanitized by
washing with a solution of 4 mL DC-2000 in 1 litre of warm
water. Use a fresh towel for each cow. Avoid contamination of
sanitizing solution by dirt and soil. Do not dip used towel back
into sanitizing solution. When solution becomes visibly dirty,
discard and provide fresh solution.

PRECAUTIONS: DANGER Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID INSTRUCTIONS In case of contact, immediately flush eyes or skin with plenty of water for at least 15 min. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

TOXICOLOGICAL INFORMATION Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

STORAGE AND DISPOSAL Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Triple rinse (or equivalent) and dispose.

Disinfection in Hospitals, Nursing Homes, and other Health Care Institutions For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard surfaces. Add 16 - 17 mL DC-2000 to 1 litre water. (1 part to 60 parts) Apply to previously cleaned hard surface with mop or cloth. At this use-level, DC-2000 is effective against *Pseudomonas aeruginosa*.

Disinfectant in Institutions (i.e. Educational, Penal, Military), Industry Schools and Homes For disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard surfaces. Add 8 mL DC-2000 to 1 litre of water. (1 part in 125 parts). Apply to previously cleaned hard surface with mop or cloth. At 8 mL per 1 litre use-level, DC-2000 is effective against *Staphylococcus aureus*, and *Salmonella choleraesuis*

Disinfection of Barber Tools Barber tools such as combs, brushes, razors, and scissors can be disinfected by immersing in a 8 mL per 1 litre solution of DC-2000.

Disinfectant of Poultry Equipment, Animal Quarters and Kennels. Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Poultry

brooders, watering founts, feeding equipment and other animal quarters can be disinfected after thoroughly cleaning by applying a solution of 16 -17 mL DC-2000 per 1 litre of water with a mop, cloth or brush. Small utensils should be immersed in this solution. After disinfection, ventilate buildings, coops, and other closed spaces. All treated equipment that will contact feed or drinking water must be rinsed with potable water.

NOTICE TO USER This control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))