

03-MAY-2004 Notification received of label changes. See below for these changes.

GROUP **1 2** HERBICIDE

**READ LABEL AND DIRECTIONS FOR USE
BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

**NET CONTENTS:
CHAMPION FM - 8.1 LITRES (1.8 Gal)
PLUS - 81 GRAMS (2.9 OZ)**

**CHAMPION(TM) PLUS
POST-EMERGENT HERBICIDE
TANK MIX**

**FOR USE IN BARLEY
FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND
PEACE RIVER REGION OF BRITISH COLUMBIA ONLY**

COMMERCIAL

WARNING - EYE IRRITANT



THIS TANK MIX USES:

**CHAMPION FM Reg. No. 23524
PEST CONTROL PRODUCTS ACT**

GUARANTEE:

Fenoxaprop-p-ethyl: 45 g/L (7.2 oz/gal)

MCPA (present as 2-ethylhexyl ester): 210 g/L (33.7 oz/gal)

2,4-D (present as 2-ethylhexyl ester): 70 g/L (11.2 oz/gal)

PLUS Reg. No.: 23566

PEST CONTROL PRODUCTS ACT

GUARANTEE: Thifensulfuron methyl 75%

DISTRIBUTED BY:

Notification Change

~~DuPont Canada Inc.~~ E.I. DU PONT CANADA COMPANY

Agricultural Products

Box 2300, Streetsville,
Mississauga, ON L5M 2J4
Toll Free: 1-800-667-3925

Champion FM is manufactured
and registered by:
Aventis CropScience Canada Co.
295 Henderson Drive
Regina, SK S4N 6C2
Toll Free: 1-888-AVENTIS
(1-888-283-6847)

DIRECTIONS FOR USE

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA ONLY

GENERAL INFORMATION

CHAMPION PLUS Tank Mix provides control of certain annual grassy and broadleaf weeds in Barley. ALL THE COMPONENTS IN THE CHAMPION PLUS TANK MIX PACKAGE MUST BE APPLIED TOGETHER ACCORDING TO LABEL RECOMMENDATIONS OR A REDUCTION IN WEED CONTROL AND/OR CROP TOLERANCE WILL OCCUR. This CHAMPION PLUS Tank Mix package contains one 8.1 litre (1.8 gal.) container of CHAMPION FM, and one 81 gram (2.9 oz.) container of PLUS.

CHAMPION PLUS Tank Mix may NOT be used on any crop for which it is NOT registered. This product is registered for use in barley only.

SUSCEPTIBLE WEEDS

Annual Grassy Weeds - Wild Oats, Green and Yellow Foxtail (Wild Millet).

Annual Broadleaf Weeds - Annual Sunflower, Burdock, Chickweed, Cocklebur, Corn Spurry, Cow Cockle, Field Horsetail*, Flixweed**, Green Smartweed, Hemp-nettle, Hoary Cress*, Kochia, Lady's-thumb, Lamb's-quarters, Mustards (except Dog and Green Tansy), Plantain*, Prickly Lettuce, Ragweeds, Redroot Pigweed, Russian Pigweed, Russian Thistle, Shepherd's-purse**, Stinkweed**, Vetch, Volunteer Rapeseed, Wild Buckwheat, and Wild Radish.

*top growth control only

**spring seedlings only

Canada Thistle Suppression - Suppression can be expressed as a reduction in height and/or a retardation and/or a yellowing of the leaves and/or a browning of the leaves. Suppressed plants will usually recover.

TIME OF APPLICATION

Treatment at the 3 to 4 leaf stage of both crop and grassy weeds combines maximum crop tolerance and weed susceptibility. However, Champion Plus may be applied to barley that has a minimum of 2 leaves and up to a maximum of 5 leaves on the main stem. Plants must not have more than 2 tillers.

Application of Champion Plus past the emergence of the second tiller will result in crop damage. The two tillers stage of barley usually occurs within 25 days after seeding.

[BARLEY LEAF STAGE DIAGRAM]

NOTE: CROP DAMAGE WILL OCCUR IF APPLICATION IS MADE EARLIER OR LATER THAN THE RECOMMENDED LEAF STAGES ON THIS LABEL.

NOTE: CHAMPION FM CONTAINS THE HERBICIDES MCPA AND 2,4-D. THESE HERBICIDES HAVE BEEN KNOWN TO CAUSE HEAD DEFORMATION IN CEREALS IF APPLIED AT EARLY STAGES. THESE SYMPTOMS DO NOT USUALLY AFFECT YIELD OR QUALITY OF GRAIN.

ANNUAL GRASSY WEEDS - Apply when the annual grassy weeds have 1 leaf up to a maximum of 5 leaves on the main stem plus 2 tillers.

BROADLEAF WEEDS - Apply according to the following chart:

BROADLEAF WEEDS	TIME OF APPLICATION	BROADLEAF WEEDS	TIME OF APPLICATION
Annual Sunflower		Corn Spurry	
Ball Mustard		Cow Cockle	
Burdock		Green Smartweed	
Cocklebur		Hemp-nettle	
Field Horsetail		Lady's Thumb	
Flixweed		Lamb's-quarters	Weeds less than 10
Hoary Cress		Redroot Pigweed	cm (4 inches) tall or
Kochia		Russian Thistle	across
Mustards*	Weed Seedling (2 to 4	Stinkweed	
Plantain	leaf stage)	Volunteer Rapeseed	
Prickly Lettuce		Wild Mustard	
Ragweeds			
Russian Pigweed			
Shepherd's-purse		Chickweed	Weed 1-6 leaf stage
Vetch			
Wild Radish			Weed 1-3 leaf stage
		Wild Buckwheat	
			Weeds less than 10
			cm (4 inches)
		Canada Thistle	
		(suppression only)	

* except dog and green tansy

Apply to emerged, young actively growing weeds. Weeds that emerge after application will not be controlled.

NOTE: Treatment at the 3 to 4 leaf stage of crops and weeds usually combines maximum crop tolerance and weed susceptibility. Some broadleaf weeds may not be controlled if infestation is

heavy, weeds are in bud or weather is dry and cool. Under stressed conditions and/or heavy crop canopy, earlier application will result in improved grassy weed control.

RATE OF APPLICATION OF CHAMPION PLUS TANK MIX

One container of CHAMPION FM 8.1L (1.8 gal.) combined with one container of PLUS 81g (2.9 oz.) treats 4.0 hectares (10 acres).

BARLEY

Apply CHAMPION FM at 2.0 Litres per hectare (0.178 gal. per acre). Apply PLUS at 20 grams per hectare (0.29 oz. per acre). **DO NOT USE A SURFACTANT WITH THIS TANK MIX.**

CAUTION

PLUS MUST BE TANK-MIXED AT THE RECOMMENDED RATE WITH CHAMPION FM OR CROP DAMAGE AND/OR A REDUCTION IN GRASSY AND BROADLEAF WEED CONTROL WILL RESULT.

CAUTION

DO NOT TREAT BARLEY IF UNDERSEEDED TO FORAGE CROPS.

METHOD OF APPLICATION

Application should be made in a minimum of 100 litres of water per hectare (10 gallons of water per acre) at a pressure of 275 kPa (40 psi) or 310 kPa (45 psi) if using check valves, and at a groundspeed of 6 to 8 km per hour (4 to 5 mph). The use of 80 degree or 110 degree flat fan nozzles is recommended for optimal spray coverage.

DO NOT USE FLOOD JET NOZZLES, CONTROLLED DROPLET APPLICATION EQUIPMENT OR SPRAFOIL EQUIPMENT.

DO NOT APPLY CHAMPION PLUS TANK MIX BY AIR.

UNIFORM, THOROUGH COVERAGE IS IMPORTANT TO OBTAIN GOOD CONTROL.

Application of the spray at a forward angle of 45 degrees will result in better penetration of the canopy and better coverage. Apply when weeds are young and actively growing. Do not apply if rain is expected within 2 hours after spraying.

CAUTION

Apply uniformly and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over application.

CAUTION

A time interval of seven (7) days prior to application, or four (4) days after application of CHAMPION PLUS is required before any other pesticide can be applied.

PRE-HARVEST INTERVAL:

DO NOT APPLY CHAMPION PLUS TANK MIX WITHIN 50 DAYS OF HARVEST.

GRAZING RESTRICTION:

Do not graze the treated crop or cut for hay; sufficient data are not available for fenoxaprop-p-ethyl to support such use.

NOTE: Prior to using CHAMPION PLUS Tank Mix, complete a thorough cleaning of the spray tank, lines, and filter. Spray equipment should be triple rinsed using a strong detergent solution. A scrubbing of the tank is also highly recommended.

MIXING INSTRUCTIONS

1. Assure that the spray tank is thoroughly clean.
2. Fill the tank half full with clean water and start agitation or bypass system.
3. Mix PLUS (container #1) in water by slowly adding the correct amount of PLUS to 20 litres (4.5 gal) of water and add to spray tank. Agitate thoroughly until PLUS is completely in suspension. Put in all containers of #1 if using more than one container.
4. Add the correct amount of CHAMPION FM (container #2) and continue agitation. Put in all containers of #2 if using more than one container.
5. Triple rinse the emptied containers into the spray tank.
6. Add the remaining amount of water while agitation continues. Spray out immediately. Spray mixture should not be left in tank overnight.
7. Thoroughly clean the spray tank after using CHAMPION PLUS Tank Mix and before using any other pesticide.

SPRAYER CLEANUP

Do not drain or flush equipment on or near desirable vegetation. To avoid subsequent injury to other crops, immediately after spraying and prior to spraying other crops, thoroughly remove all traces of CHAMPION PLUS Tank Mix from mixing and spray equipment as follows:

SPRAYER CLEANOUT INSTRUCTIONS WHEN MOVING INTO WHEAT, RYE OR FLAX

When moving into WHEAT, SPRING or FALL RYE, or FLAX immediately following the application of CHAMPION PLUS Tank Mix, clean the sprayer by thoroughly flushing with water/detergent or cleaning agent.

NOTE: BROADLEAF CROPS CAN BE DAMAGED BY CHAMPION PLUS TANK MIX RESIDUES IN THE SPRAY TANK EVEN AFTER A NUMBER OF APPLICATIONS OF A DIFFERENT PRODUCT. IT IS CRITICAL TO THOROUGHLY CLEAN AND REMOVE ALL TRACES OF CHAMPION PLUS TANK MIX FROM THE SPRAY TANK PRIOR TO MOVING INTO A BROADLEAF CROP.

WHEN MOVING INTO BROADLEAF CROPS

IN ALL CASES, prior to spraying a broadleaf crop (such as CANOLA, PEAS, LENTILS, ALFALFA, SUGARBEETS, VEGETABLES, etc.) complete a thorough cleaning of the tank, because the Plus component of the CHAMPION PLUS tank mix can cause crop injury to sensitive crops at very low concentrations. Follow the cleanup instructions below to ensure adequate sprayer cleaning and removal of the CHAMPION PLUS Tank Mix.

CLEANUP INSTRUCTIONS PRIOR TO SPRAYING BROADLEAF CROPS

The cleaning agent referred to may be either household ammonia (minimum 3% ammonia) or an ammonia-based clean agent containing at least 3% ammonia.

1. Remove nozzles and screens and clean separately in a bucket containing water and cleaning agent.
2. Drain tank and flush tank, boom, and hoses with clean water for at least 10 minutes. Inspect tank to ensure all visible residues have been removed. If necessary, repeat this step. **DO NOT CLEAN SPRAYER NEAR WELL OR WATER SOURCE OR NEAR DESIRABLE VEGETATION.**

If the sprayer tank is equipped with rotary washing nozzles, go to step 6.

3. Fill the tank with clean water while adding 1 litre (35 oz.) cleaning agent per 100 litres (22 gal.) water. Flush solution through boom and hoses and then add enough water to fill the tank. Allow to stand for 15 minutes with agitation. Flush booms and hoses again, and drain tank.
4. Repeat step 3.
5. Go to and complete step 9.
6. If the sprayer tank is equipped with rotary nozzles, use the following procedure. Fill tank approximately one quarter full with water, and add 1 litre (35 oz.) of cleaning agent for each 100 litres (22 gal.) of water.
7. Operate the rotary nozzles for at least 15 minutes, ensuring that all inside surfaces of the tank are covered by the water/cleaning agent mixture. After about 5 minutes of agitation, flush the solution through the hoses and boom for about 2 minutes. Drain tank.
8. Repeat steps 6 and 7.
9. Thoroughly rinse the tank, hoses and boom with clean water for at least 5 minutes.

CAUTION

Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour which may cause eye, throat and lung irritation.

CAUTION

Do not allow spray or vapour to drift onto crops which can be damaged such as peas, beans, sugar beets, tomatoes, clovers, alfalfa, canola, sunflowers, potatoes, tame oats, turnips, vegetables, flowers, trees or shrubs.

CAUTION

Do not tank-mix with any other chemical additives, pesticides, or fertilizers unless recommended on this label. Do not use a surfactant with this tank mix.

CAUTION

Application of CHAMPION PLUS Tank Mix must take place within the leaf stages outlined for all weeds on this label. Late application will result in reduced weed control.

CAUTION

During periods of stress, plants are not actively growing. When daytime temperatures before or after application are very hot combined with very dry conditions and low humidity, plants are under stress. Application of CHAMPION PLUS Tank Mix during these periods may result in substantially reduced control.

CAUTION

If application of CHAMPION PLUS Tank Mix is not possible due to uneven weed germination or differential germination of weeds and crop, consult your local weed specialist or the provincial weed control recommendations for the best alternative method of weed control.

CAUTION

Do not apply to crop that is stressed by severe weather conditions, frost, low fertility, drought, water saturated soil, disease or insect damage as crop injury may result.

CAUTION

TOXIC TO FISH. Do not apply within 15 metres (50 feet) of a body of water or wetland area. Do not contaminate water supply, ponds, lakes, rivers, streams and irrigation ditches by direct application, spray drift or when cleaning and rinsing spray equipment or containers.

CAUTION

Do not apply to land where tail water will be used to irrigate crop land. Do not contaminate irrigation water.

CAUTION

Do not exceed a total of 20 grams per hectare (0.29 oz. per acre) per crop year of PLUS (thifensulfuron-methyl). This tank-mix should be applied only once a year.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, CHAMPION FM and PLUS Herbicides of the CHAMPION PLUS Post-Emegent Herbicide Tank Mix are Group 1 and 2 herbicides, respectively. Any weed population may contain or develop plants naturally resistant to the components of CHAMPION PLUS Post-Emegent Herbicide Tank Mix and other Group 1 and 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of CHAMPION PLUS Post-Emegent Herbicide Tank Mix or other Group 1 and 2 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your local DuPont representative, or call DuPont Canada Inc. at 1-800-667-3925 toll free.

STORAGE

Keep away from fire or open flame or other sources of heat. CANNOT be stored below freezing. If stored for 1 year or longer, shake well before using. Store the tightly closed container away from feed, seeds, fertilizer, plants and foodstuffs. Do not use or store in or around the home. Keep in original container during storage.

RECYCLABLE CONTAINER DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

DISPOSAL OF UNUSED SPRAY SOLUTION:

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the field just sprayed, not over crop just sprayed, away from water supplies, ditches, and irrigation ditches.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. WARNING EYE IRRITANT. CAUSES SKIN IRRITATION. DO NOT GET IN EYES, ON SKIN OR CLOTHING.

HARMFUL IF INHALED. DO NOT BREATHE VAPOUR OR SPRAY MIST. WEAR A RESPIRATOR WITH A SUITABLE CARTRIDGE DURING MIXING AND LOADING.

TO MINIMIZE EXPOSURE, THE FOLLOWING PRECAUTIONS SHOULD BE FOLLOWED:

1. Protective Clothing and Equipment

- **Handling the concentrate (mixing and loading):** wear long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes, and protective eye wear (face shield or safety glasses). Rinse gloves before removal.
- Coveralls or a chemical-resistant apron should also be worn when open pouring from containers greater than 5 L.
- **Handling the dilute spray solution (during application or repairing or cleaning equipment):** wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Rinse gloves before removal.

- Gloves are not required during application when applicator is in an enclosed tractor.

2. Operator Use Precautions

- Wear freshly laundered clothing and clean protective equipment daily.
- Rinse gloves before removal.
- Wash hands before eating, drinking, using tobacco or using the toilet.
- If herbicide penetrates clothing remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate.
- After using this product, remove clothing and launder separately and promptly, and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.
- After work, remove all clothing and shower using soap and water.

Avoid spray drift to susceptible plants and USE ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED RATES.

This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. DO NOT contaminate these systems through direct application, disposal of waste or cleaning equipment.

If this product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., contact 1-866-375-4648 or www.cropro.org/.

FIRST AID: IN CASE OF POISONING, CALL A PHYSICIAN OR POISON CONTROL CENTRE IMMEDIATELY. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

INGESTION:

If the tank-mix spray solution or CHAMPION FM are swallowed do NOT induce vomiting but rush patient to nearest hospital taking these Directions For Use or the labelled containers with you. If PLUS is swallowed, administer water freely and induce vomiting. Repeat until vomit fluid is clear. This tank mix contains PETROLEUM DISTILLATES. Contact a doctor or Poison Control Centre.

SKIN CONTACT:

In case of skin contact, remove contaminated clothing and wash skin thoroughly with soap and water. If irritation persists get medical attention.

EYE CONTACT:

If in eyes, flush with plenty of water for 15 minutes and take patient to hospital at once, taking these Directions For Use or the labelled containers with you.

INHALATION:

If inhaled, remove patient from site of exposure.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label, and subject thereto the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

FOR MORE INFORMATION CONTACT:

E.I. DU PONT CANADA COMPANY

~~DU PONT CANADA INC.~~ AGRICULTURAL PRODUCTS

Notification Change

Box 2300, Streetsville
Mississauga, Ontario
L5M 2J4
Toll Free: 1-800-667-3925

CHAMPION PLUS is Distributed by DuPONT CANADA INC.
CHAMPION PLUS is a trademark of DuPONT CANADA INC.

SPRAFOIL is a registered trademark of D&W Industries.

ALLY, REFINE, EXPRESS, MUSTER are registered trademarks of E.I. DuPont de Nemours.

AMBER is a registered trademark of Novartis Crop Protection, Inc.

LASER and TRIUMPH PLUS are trademarks of the Aventis Group.

05-JUL-2002

PLUS DRY FLOWABLE HERBICIDE

GROUP □ 2 HERBICIDE

For Sale For Use on Barley in the Prairie Provinces and Peace River Region of British Columbia.

COMMERCIAL

READ THE LABEL AND ENCLOSED DIRECTIONS FOR USE BEFORE USING

This product must be used in combination with Champion F.M.
For use only on crops registered on the Champion Plus Herbicide Tank Mix Booklet.

NET CONTENTS: 81 grams

REGISTRATION NO. 23566 PEST CONTROL PRODUCTS ACT

GUARANTEE: Thifensulfuron methyl75%

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.
AVOID BREATHING SPRAY MIST.
AVOID CONTACT WITH SKIN, EYES AND CLOTHING.
DO NOT CONTAMINATE ANY BODY OF WATER.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., contact 1-866-375-4648 or www.cropro.org/.

FIRST AID:

EYE CONTACT: Flush eyes immediately with plenty of water for 15 minutes. Contact a physician or a poison control centre IMMEDIATELY

SKIN CONTACT: Wash skin with plenty of soap and water. Remove and wash contaminated clothing before re-use.

INGESTION: Administer water freely and induce vomiting. Repeat until vomit fluid is clear. Contact a physician.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies and transportation emergencies such as spills, leaks or fires, call 1-613-348-3616 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.
DUPONT CANADA INC.
AGRICULTURAL PRODUCTS
BOX 2300 STREETSVILLE, MISSISSAUGA,
ONTARIO L5M 2J4

1-800-667-3925

STORAGE AND DISPOSAL

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, seed, food or feed. Not for use or storage in or around the home. Keep container closed.

DISPOSAL:

1. Triple- or pressure-rinse the container thoroughly. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the empty container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of spill, and for clean up of spills.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, PLUS DRY FLOWABLE HERBICIDE is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to PLUS DRY FLOWABLE HERBICIDE and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the used of PLUS DRY FLOWABLE HERBICIDE or other Group 2 herbicides with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from a different group when such use is permitted
- ◆ Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- ◆ Monitor treated weed populations for resistance development.
- ◆ Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information or to report suspected resistance, contact your local DuPont representative or the DuPont hotline at 1-800-667-3925 for further information.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

PLUS HERBICIDE is manufactured by E.I. du Pont de Nemours and Co. and is registered by DuPont Canada Inc.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))