

2002/2005

CO-OP (R)

WARFARIN RAT KILLER

RODENTICIDE PELLETS

DOMESTIC

500 g

GUARANTEE:

WARFARIN	0.025%
SULFAQUINOXALINE	0.025%

REGISTRATION NO. 11164

PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

Interprovincial Cooperative Limited
P.O. Box 1050,
Saskatoon, Saskatchewan
S7K 3M9
WINNIPEG, TORONTO, MONTREAL, MONCTON

RC-000-0297

DIRECTIONS:

Rats: Place 250 g to 500 g of bait in a dry container in locations frequented by rats. (Tunnels, Feeding areas and Runways.)

Mice: Place 15 - 30 grams of bait in shallow containers where mice are seen or signs of mice are apparent. Place bait stations 3 - 4 metres apart and as many bait stations as practical.

Assure an uninterrupted supply of fresh bait for a period of not less than 12 days or until feeding ceases. Always make water available in shallow containers near bait.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN, PETS, DOMESTIC ANIMALS AND WILDLIFE. If taken in large or repeated doses by humans or

animals, it may reduce the clotting ability of the blood and cause internal bleeding. Do not expose treated bait in areas accessible to children, dogs, domestic animals and wildlife except when enclosed within a tamper-resistant bait box.

- (A) Keep away from feed or food stuffs.
- (B) Do not get in eyes, on skin or on clothing. Wash arms, hands and face with soap and water after handling and before eating or smoking.
- (C) Keep out of lakes, streams or ponds.

FIRST AID:

If in eyes, flush with water for 15 minutes. If on skin, wash with soap and water. If swallowed, give patient one to two glasses of water and induce vomiting by touching back of throat with a finger or blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. Contact a physician or Poison Control Centre immediately.

TOXICOLOGICAL INFORMATION:

Vitamin K1 in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

DISPOSAL:

Do not reuse empty container. Dispose of empty container by placing in household garbage.

RODENT DISPOSAL:

Dispose of dead rodent in garbage or by burying.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

(R) Registered trade-mark of TMC Distributing Ltd. Used under Licence.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))