

90.12.17

CO-OP (R) Warble Killer

R.T.U.
READY TO USE

Contains Trichlorfon

READ THE LABEL BEFORE USING

POUR-ON TREATMENT, READY-TO-USE.

KILLS CATTLE GRUBS AND LICE ON BEEF
AND NON-LACTATING DAIRY ANIMALS

AGRICULTURAL

GUARANTEE: Trichlorfon 8%

Registration No. 15258
Pest Control Products Act

Protect from extreme heat and protect from sunlight.

4 L

INTERPROVINCIAL COOPERATIVE
P.O. Box 1050, Saskatoon, Saskatchewan
Winnipeg, Toronto, Montreal, Moncton

DIRECTIONS:

Co-Op Warble Killer is ready-to-use. No dilution is necessary. Do not apply during December, January or February. Controls cattle grubs and prevents the build-up of lice on all beef and non-lactating dairy cattle.

Apply 30 mL of product per 100 kg of body weight. Pour the required amount uniformly along the back line. Apply as soon as possible in the fall after heel fly activity has ceased. Consult

your local Ag rep. or veterinarian for the optimum time to apply in your area. Can also be applied in the spring, ideally at least six weeks before grubs appear in the back.

Animals heavily or chronically infested with lice should be treated in the fall with an overall spray such as Co-op Malathion. Do not make a separate louse spray within two weeks of warble treatment.

The contents of this package will treat:

Weight of animals (kg)

100	150	200	250	300	350	400	450	500
-----	-----	-----	-----	-----	-----	-----	-----	-----

Number of animals:

133	89	67	53	44	38	33	30	27
-----	----	----	----	----	----	----	----	----

PRECAUTIONS:

KEEP OUT OF THE REACH OF CHILDREN. May be harmful if swallowed or absorbed through the skin. Avoid contact with skin and eyes. Wash body and clothing thoroughly after handling. Do not contaminate food or feed.

1. Do not apply to animals less than three months of age.
2. Do not apply to lactating animals within seven days of freshening. If freshening should occur within seven days of treatment, do not use milk for human consumption for the balance of the seven day interval.
3. Do not apply to cattle within twenty-one days of slaughter.
4. Do not treat animals for ten days before or after shipping, weaning or dehorning or after exposure to contagious or infectious diseases.
5. Do not apply in conjunction with oral drenches, other internal medications or with organic phosphates.
6. Do not apply during December, January or February.

FIRST AID:

If swallowed, induce vomiting. If in eyes, flush with water for fifteen minutes. In either case, obtain prompt medical attention.

TOXICOLOGICAL INFORMATION:

Cholinesterase inhibitor. Atropine is the antidote. 2-PAM may be administered in conjunction with Atropine.

NON-WARRANTY:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising out of the use or handling of this product and accepts the product on that condition.

NOTICE TO USER:

This control product is to be used in accordance with the directions on the label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))