

90.09.18

RHONE-POULENC

MANEB 80-W

NET WEIGHT 10 KG

FUNGICIDE  
WETTABLE POWDER

AGRICULTURAL

GUARANTEE:

MANEB 80%

REGISTRATION NO. 14768  
PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

RHONE-POULENC CANADA INC.  
2000 ARGENTIA ROAD, PLAZA 3, SUITE 400,  
MISSISSAUGA, ONT. L5N 1V9

IN CASE OF SPILLS, POISONING OR FIRE  
TELEPHONE EMERGENCY RESPONSE NUMBER COLLECT  
(416) 634-2359 (24 HOURS A DAY).

COMPATIBILITY: This product is compatible with most commonly used insecticides and fungicides, except Bordeaux and lime.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Hazardous if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin. Avoid inhalation of dust or spray mist. After handling, wash thoroughly with soap and water before eating, drinking or smoking and change to clean clothing. Do not apply or allow to drift to areas occupied by unprotected persons or to streams, lakes or ponds to protect wildlife. Avoid contamination of feed or food, including such crops on which residue is unsafe.

FIRST AID: If swallowed, induce vomiting by touching the back of the tongue with a blunt object; repeat until vomit fluid is clear. Take patient to nearest hospital taking this labelled container with you. In case of skin contact, remove contaminated clothing and wash skin with soap and water. If in eyes, flush with plenty of water for 15 minutes and get medical attention.

STORAGE: Store product in a cool, dry place away from food or feed. Prevent the contents from becoming wet as this will reduce effectiveness and may create flammable vapours. Keep away from fire and sparks.

#### DISPOSAL

1. Empty bag.
2. Follow provincial instructions for any required additional cleaning of the bag prior to its disposal.
3. Make the empty bag unsuitable for further use.
4. Dispose of the bag in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills contact the regional office of Conservation and Protection, Environment Canada.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

DIRECTIONS FOR USE Apply MANEB 80W in sufficient water for thorough coverage of the foliage; the required volume depends on the spray equipment and size of plants.

TOMATO: Anthracnose; Early Blight; Grey Leaf Spot; Grey Mould; Late Blight; Septoria Leaf Spot.

METHOD/RATE: Apply 2.25 kg/ha when the first fruit has formed and repeat at 7-10 day intervals. When weather favours disease or in case of severe disease outbreak, apply 3.25 kg/ha every 5-7 days. For Anthracnose apply not less than 3.25 kg/ha. Do not apply later than 3 days before each picking.

CUCUMBER, MELON, PUMPKIN, SQUASH: Alternaria and Septoria Leaf Spot; Anthracnose; Downy Mildew.

METHOD/RATE: Apply 2.25 - 3.25 kg/ha as soon as plants begin to vine or disease threatens. Repeat at 7-10 day intervals. Use the higher rate when vines cover the ground and if disease is present. Do not apply later than 5 days before each picking.

ONION: Botrytis Leaf Blight; Downy Mildew; Purple Blotch.

METHOD/RATE: Apply 3.25 kg/ha when disease threatens and repeat at 7-10 day intervals. Add 125-250 mL of Super Spred per 1000 litres of spray mixture. Do not apply later than 10 days before harvest.

CABBAGE, ENDIVE, LETTUCE: Downy Mildew; Alternaria Leaf Spot.

METHOD/RATE: Apply 2.25 kg/ha when disease threatens and repeat at 7-10 day intervals. If disease is present and weather favours disease apply every 3-5 days and increase rate for lettuce to 3.25 kg/ha. On cabbage plants beds start spraying 7-10 days after seeding and repeat every 3 days until transplanting. Do not apply later than 7 days before harvest.

CARROT: Alternaria Blight; Cercospora Blight.

METHOD/RATE: Apply 3.25 kg/ha starting in late July and repeat at 7 day intervals. A total of 5 sprays may be needed. Do not apply later than 5 days before harvest.

CELERY: Early Blight; Late Blight.

METHOD/RATE: Apply 2.25 kg/ha when seedlings emerge and repeat at 3-5 day intervals until transplanting. Continue spraying in the field every 7 days. Do not apply later than 14 days before harvest and trim stalks in the field during harvest.

POTATO: Early Blight; Late Blight.

METHOD/RATE: Apply 2.25 kg/ha. For Early Blight start spraying when plants are 15 cm high; for Late Blight start in mid-July or when disease is reported in the area. Repeat every 7-10 days and shorten interval to 5-7 days when weather favours disease. May be applied up to 1 day before harvest.

SUGAR BEETS: Cercospora Leaf Spot.

METHOD/RATE: Apply 2.25 kg/ha when disease is first noticed and repeat at 7-10 day intervals depending on weather. Do not apply later than 14 days before harvest.

PEACH, APRICOT: Coryneum Blight; Brown Rot; Leaf Curl.

METHOD/RATE: In British Columbia, apply 9kg/ha, 2.5 kg/1000 L. For Coryneum Blight, apply at Husk-fall; for Brown Rot, apply at Pink and Full-bloom; for Leaf Curl, apply at Dormant stage.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))