

2001/2006

**WILSON**  
**PROLIN**  
**CONCENTRATE**

25 kg

COMMERCIAL

Kills Rats & Mice

REGISTRATION NO. 8699 PEST CONTROL PRODUCTS ACT

GUARANTEE:

Warfarin.....	0.5%
Sulfaquinoxaline.....	0.5%

For Making Your Own Baits

**READ LABEL BEFORE USE**

**DIRECTIONS FOR USE:** Thoroughly mix 1 part of the contents of this package with 19 parts suitable bait material to make a finished bait. Fresh corn meal, rolled oats, dog food or similar substances that rodents prefer in your locality are suitable bait materials. If the bait tends to be dry or dusty, a small amount of salad oil or cooking oil may be added.

Place approximately 50 grams for house mice (*mus musculus*); 200 g to 400 g for rats; norway or brown rat (*rattus norvegicus*), roof or black rat (*rattus rattus*), in shallow, dry containers, where rodents are accustomed to feed. Bait stations at intervals of approximately 3 m for mice and as many as practical for rats are recommended. Assure an uninterrupted supply of bait until all signs of feeding have stopped. Make water available in shallow containers near bait.

For areas not directly related to food processing: Use only in non-food or non-feed areas where feed, food, packaging and handling equipment are never opened or exposed.

For areas where feed or food is processed, served or stored: In meat and food processing plants (processing areas) use only when plant is not in operation. Remove or cover all food, packaging

material and utensils before placing bait in baiting stations. Remove all baits and dead rodents before re-use of the plant (processing areas include storage and service).

**PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK.**

Cover with bait boxes if necessary. If swallowed by humans, pets or domestic animals, this material can reduce clotting ability of the blood.

**FIRST AID: If swallowed,**

cause vomiting until fluid is clear, call a doctor or Poison Control Centre in case of accident. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**NOTICE TO USER:** This control product is to be used only in accordance with the directions on this label. It is an offence under The Pest Control Products Act to use a control product under unsafe conditions.

**TOXICOLOGY:** Vitamin K1 in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

**STORAGE AND DISPOSAL: DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OF DISPOSAL.**

**STORAGE:** Store only in original container in a dry place inaccessible to children and pets.

**DISPOSAL:**

1. Follow provincial instructions for any required additional cleaning of the container prior to disposal.
2. Make the empty container unsuitable for other uses.
3. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused unwanted product and cleanup of spills contact the provincial regulatory agency or the manufacturer.

**Wilson Laboratories Inc.,  
10 Craig St., Brantford, ON  
N3R 7J1**

**77-0960-001**

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace

the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+) )