

85.12.31

GREEN CROSS

ROSE DUST

SQUEEZE DUSTER

INSECTICIDE/FUNGICIDE

DOMESTIC

READ THE LABEL BEFORE USING

GUARANTEE:

CARBARYL 5%, MALATHION 4%, CAPTAN 5%, SULPHUR 20%

REG. NO.: 10559

P.C.P. ACT

300 g

WARNING: Exposure to captan may produce long-term health Effects. To minimize exposure, follow directions outlined on this label.

GREEN CROSS

DIVISION OF CIBA-GEIGY CANADA LTD.

6860 CENTURY AVENUE

MISSISSAUGA, ONTARIO L5N 2W5

DIRECTIONS: To control aphids, red spider mites, rose chafers, rose leafhoppers and certain chewing insects, dust the upper and lower leaf surfaces lightly and evenly. Application is recommended before early morning dew dries. Dusting at other times is less effective. Repeat at frequent intervals. Helps control powdery mildew and black spot. NOTE: 1. Do not use on Boston ivy or virginia creeper. 2. HIGHLY TOXIC TO BEES. 3. Do not refill more than 2/3 full. 4. Do not apply when temperature is above 27°C.

CAUTIONS: KEEP OUT OF REACH OF CHILDREN. Poisonous if swallowed. Never handle material with bare hands. Use rubber gloves; do not use leather or cloth gloves. Wash hands, face and arms thoroughly after handling this product and before drinking, eating or smoking. Do not breathe dust. When handling or

applying, wear goggles and clothing that completely covers arms and legs to minimize exposure. Change contaminated clothing and wash thoroughly before re-use. If an accident occurs, remove contaminated clothing immediately, and wash contaminated skin thoroughly with soap and water. Wash clothing and gloves before re-use. Always work out of the wind when handling, or applying treatments. Keep unprotected persons out of operating area or vicinity where there may be drift. Avoid contaminating food and feed.

FIRST AID: In case of poisoning, call a physician immediately. IF ON SKIN, remove contaminated clothing and wash skin thoroughly with soap and water. IF IN EYES, flush with water for 5-10 minutes and obtain medical attention. IF SWALLOWED, give the patient one to two glasses of water and cause vomiting by giving one dose of syrup of ipecac. If the patient does not vomit within 20 minutes, give a second dose. If syrup of ipecac is not available, give the patient one to two glasses of water and cause vomiting by inserting a finger down the throat. Repeat with water until vomit fluid is clear. The patient should be lying down with the head below the level of the feet. DO NOT TRY TO CAUSE VOMITING IF THE PATIENT IS UNCONSCIOUS OR IN A CONVULSIVE STATE.

DECONTAMINATION AND DISPOSAL: Discard empty containers with household garbage.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))