

95/2000

DOMESTIC
Free Measuring
Cup Inside

250 mL
500 mL
1 L

C-I-L

AMBUSH(R)

Tree and Garden Insect Killer
FOR OUTDOOR USE

READ THE LABEL AND BOX BEFORE USING

KEEP OUT OF REACH OF CHILDREN

GUARANTEE: permethrin 12.5 g per litre

REGISTRATION NO. 18868 PEST CONTROL PRODUCTS ACT

KILLS OVER 40 DIFFERENT INSECTS

ZENECA Agro
A business ZENECA Corp.
Stoney Creek, Ontario
L8G 3Z1

Made in Canada

A93/D-3

KEEP BOX FOR USE DIRECTIONS

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

Wear protective gloves and long-sleeved clothing when handling product. Avoid inhaling spray or contact with eyes, skin or clothing. Avoid contamination of food, feed, utensils or drinking water. DO NOT SPRAY WITHIN 15 M OF ANY BODY OF WATER. Wash thoroughly after use and before eating, drinking and smoking. This product is very toxic to fish. Remove fish bowls

and aquariums from the spray area. Do not apply when weather conditions favour drift. Toxic to bees - Do not spray if bees are present.

KEEP CHILDREN AND PETS AWAY FROM TREATED AREA DURING APPLICATION AND UNTIL DRY.

DISPOSAL

Do not reuse empty container. Dispose of container with household garbage.

FIRST AID

IF SWALLOWED, induce vomiting by giving patient 1 to 2 glasses of water and inserting finger down throat. IF ON SKIN, wash with soap and water. IF IN EYES, flush with plenty of clean water. If any complications arise, see a doctor promptly.

PRODUCT INFORMATION

C-I-L Garden Insect Killer contains permethrin, a synthetic pyrethroid insecticide of low mammalian toxicity. Kills insects on contact or by ingestion.

DIRECTIONS FOR USE

5 mL = 1 teaspoon
15 mL = 1 tablespoon

Mix with water as directed below and apply with a sprayer to give thorough coverage of plants. Apply when insects or damage are first noticed and repeat as necessary.

ORNAMENTALS

CONIFEROUS TREES AND SHRUBS: Douglas-Fir Tussock Moth, White-Marked Tussock Moth, Spruce Budworm, and Yellow-Headed Spruce Sawfly, Coneworm and Open Feeding Sawflies - Mix 50 mL with 10 L of water and thoroughly apply when first signs of feeding occur. Repeat once if reinfestation occurs.

DECIDUOUS TREES AND SHRUBS: Gypsy Moth, Eastern Tent Caterpillar, Forest Tent Caterpillar, Fall Webworm, Euonymus Webworm and Open Feeding Aphids - Mix 70 mL with 10 L of water and thoroughly apply when insects appear. Repeat once if

reinfestation occurs.

FRUIT TREES

APPLES: Winter Moth, Eastern Tent Caterpillar, Eye-spotted Budmoth, Plum Curculio, White Apple Leafhopper: Apply 100 mL in 10 L of water. Tentiform Leafminer, Green Fruitworm, Apple Maggot, Leafrollers, Tarnish Plant Bug, Mullein Bug, Lesser Appleworm, Codling Moth: Apply 200 mL in 10 L of water.

PEARS: Eastern Canada - Pear Psylla (adults and nymphs) Codling Moth, Green Fruitworm: Apply 200 mL in 10 L of water. Do not apply less than 7 days before harvest.

PEACHES AND NECTARINES: Oriental Fruit Moth, Plum Curculio, Plant Bugs: Apply 200 mL in 10 L of water. Do not apply within 7 days of harvest.

PLUMS: Plum Curculio: Apply 200 mL in 10 L of water. Do not apply within 7 days of harvest.

GRAPES: Grape Berry Moth: Apply 150 mL in 10 L of water. Grape Leafhopper: Apply 75 mL in 10 L of water. Do not apply less than 21 days before harvest.

VEGETABLES

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS: Imported Cabbage Worm, Cabbage Looper, Diamondback Moth Larvae and Crucifera Flea Beetle - Mix 50 mL with 10 L of water. Do not spray cabbage, cauliflower and Brussels sprouts within 3 days of harvest; broccoli within 7 days of harvest.

SWEET CORN: European Corn Borer, Corn Earworm and Sap Beetle - Mix 100 mL with 10 L of water. European Corn Borer - In early plantings, apply when first feeding is seen on the leaves. In later plantings, apply before tassels show and repeat at 5 day intervals until early silk stage.

Corn Earworm - When 25% of ears show silk, start spraying and fully cover ears and silk. Apply 4 sprays allowing 4 days between sprays (3 day intervals in hot weather). Sap Beetles - Apply when pink-spotted beetles appear. Do not treat within 1 day of harvest.

PEPPERS: European Corn Borer - Mix 100 mL with 10 L of water and

apply if corn borer is noticed on plants (usually end of July to early September). Do not apply within 1 day of harvest.

POTATOES AND TOMATOES: Colorado Potato Beetle, Potato Flea Beetle, Potato Leafhopper, Tarnished Plant Bug and White Flies - Mix 75 mL with 10 L of water. Do not apply to tomatoes within 1 day of harvest.

Cutworms - These insects attack and cut off seedlings, transplants and low-lying plants like petunias and pansies. To confirm the cause, finger soil around damaged plant looking for a soft, dark-coloured worm in a curled position. Mix 75 mL with 10 L of water and apply to cover a ground area of 100 m². If soil is organic (muck soil) or very dry, apply 150 mL in 10 L of water per 100 m².

For Variegated cutworms (climbing cutworms) on Potatoes, Tomatoes and Ornamental Plants - Mix 75 mL with 10 L of water. Thoroughly spray the plant and underlying ground at the first sign of damage. Do not apply within one day of harvest.

NOTE: Only cutworms feeding on the soil surface and above will be controlled. Subterranean cutworms will not be controlled unless they come up and move on the treated surface.

AMBUSH is a registered trademark of ZENECA, England.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))