

85.12.31

(R)MORESTAN

2% DUST

MITICIDE, INSECTICIDE, FUNGICIDE

COMMERCIAL

FOR CONTROL OF MITES AND POWDERY MILDEW

GUARANTEE: Chinomethionat 2%

NET WEIGHT _____ KILOGRAMS

U.S. Patent No. 3,091,613
Canadian Patent No. 659,270

REGISTRATION NO. 9498
PEST CONTROL PRODUCTS ACT

CAUTION
KEEP OUT OF REACH OF CHILDREN

(See rear panel for first aid and caution statements)

READ THE LABEL BEFORE USE

CHEMAGRO LIMITED
1355 Aerowood Drive
Mississauga, Ontario,
Canada
L4W 1C2

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

DIRECTIONS FOR USE

MORESTAN Miticide-Fungicide is recommended for residual control of mites and mite eggs on deciduous fruits, non-bearing strawberries, and ornamentals.

It is also effective in controlling powdery mildew on non-bearing apples. For best results, apply a thorough coverage before mites or mite eggs are present in large numbers and before mites cause bronzing of the foliage. Apply during good drying conditions. (See Restrictions.) Consult your local Agricultural Authorities for detailed information on timing of applications for most effective control.

RECOMMENDED APPLICATIONS

CROP	PEST	DOSAGE MORESTAN 2% DUST	REMARKS
BEARING FRUIT Pre-bloom & Post-Harvest Apples Apricots Cherries Peaches Pears Plums Rune Plums	Mites Mite eggs Pear Psylla	32.5 to 55 kg/ha 85 kg/ha	Apply specified dosage per hectare as a pre-bloom treatment (delayed dormant, pre-pink, pink, or calyx.) Do not apply after calyx stage except on apples and pears (see below.) Additional applications may be made to tree foliage after harvest.
NON-BEARING FRUIT Apples	Powdery Mildew	32.5 kg/ha	Apply specified dosage per hectare. Repeat as necessary. For control of powdery mildew begin preventive applications before disease appears and apply at 7 to 14 day intervals.
Apples Apricots Cherries Peaches Pears Plums Prune Plums	Mites Mite eggs	32.5 to 55 kg/ha	Apply specified dosage per hectare. Repeat as necessary. For control of powdery mildew begin preventive applications before disease appears and apply at 7 to 14 day intervals.

Strawberries	Mites (except cyclamen mite) Mite eggs	32.5 to 55 kg/ha	Applied specified dosage per hectare. Repeat as necessary.
ORNAMENTALS Arborvitae Juniper Roses Yew	Mites Mite eggs	32.5 to 55 kg/ha	Apply specified dosage per hectare. Repeat as necessary.

RESTRICTIONS

Under some conditions, injury may occur to foliage of strawberries or deciduous fruits (apples, apricots, cherries, peaches, pears, plums and prune plums.)

Do not apply oils after delayed dormant where MORESTAN 2% Dust is to be used in pink or calyx application.

Do not contaminate streams, lakes or ponds with this material.

Do not allow dust to drift onto other crops, including pastures. Do not graze treated areas.

Do not use on other crops used for food or forage. Use only according to label directions.

Phytotoxicity has occurred on the foliage and flowers of several varieties of roses, especially during periods of high temperatures. Treat roses only after cutting and before new growth appears. Before treating large plantings for the first time, treat a few plants and observe for 4 to 5 days. Under certain conditions, some injury to other ornamentals may occur.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy it by burying with waste or burning (stay away from smoke).

CAUTION

Harmful if swallowed or inhaled. MORESTAN 2% Dust may cause skin irritation and, therefore, contact with skin, eyes, or clothing should be avoided. Avoid breathing of dust. Wash thoroughly with soap and warm water after handling.

Do not contaminate feed or food. Keep out of reach of children.

FIRST AID: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, vomiting should be induced. Administer milk or water freely and induce vomiting by giving one dose (15 mL) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. The patient should be lying down with the head below the foot level and facing down or to one side. Professional medical assistance should be secured immediately. DO NOT INDUCE VOMITING TO AN UNCONSCIOUS PERSON OR TO PERSONS IN A CONVULSIVE STATE. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash immediately with flowing water for at least 15 minutes.

TOXICOLOGICAL INFORMATION: In experimental animals acutely poisoned by MORESTAN, diarrhea and decreased activity were the only symptoms observed. The patient should be treated symptomatically.

NOTICE

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

MORESTAN is a Reg. TM of Bayer AG.

Printed in U.S.A.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))