

96.11.19

SANEX(R)

5% SEVIN DUST

DOMESTIC

INSECTICIDE

Controls insects on potatoes, cucumbers, beans
and other vegetables

300 g

CAUTION

POISON

GUARANTEE:

Carbaryl..... 5.0%

READ THE LABEL BEFORE USING

REGISTRATION NO. 17538
PEST CONTROL PRODUCTS ACT

SANEX AGRO INC.
36 Head Street
Dundas, Ontario
L9H 3H3

DIRECTIONS FOR USE: Apply a thin uniform coating of dust to the upper and lower leaf surfaces when the air is still, preferably in early morning or evening.

POTATOES: To control Colorado Potato Beetles, Flea Beetles and Leafhoppers, apply a light even coating on upper and lower leaf surfaces. For larger areas, apply 225 grams per 100 m² to control Colorado Potato Beetle and 450 grams per 100 m² to control Flea Beetles and Leafhoppers. Repeat treatment at 7-10 day intervals if necessary.

CUCUMBERS & SQUASH: To control Cucumber Beetles, Squash Bugs and Flea Beetles, dust upper and lower leaf surfaces evenly. For larger areas, use 275-450 grams per 100 m². Repeat at 7-10 day intervals if necessary.

BEANS: To control Mexican Bean Beetle, Leafhoppers and Flea Beetles, dust upper and lower leaf surfaces evenly. For larger areas use 225-450 grams per 100 m². Repeat in 10-14 days if necessary.

SWEET CORN: Corn Borer - Make first application when egg masses on leaves are in black-head stage. Apply evenly on upper and lower leaf surfaces. A minimum of 4 applications are required at 4-5 day intervals. For larger areas, apply 450-675 grams per 100 m².

CORN EARWORM: Dust leaves and ears thoroughly and evenly at 2-3 day intervals starting at first silking and continuing until silks start drying. For larger areas apply 450-675 per 100 m². Three or more applications are necessary depending upon the length of silking and the severity of the infestation.

TOMATOES: To control Flea Beetles, Fall Armyworm, Stink Bugs, Tomato Fruit Worm and Tomato Hornworm, apply a light even coating to upper and lower leaf surfaces. For larger areas use 225-325 grams per 100 m². Repeat at 7-10 day intervals if necessary.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. This product is toxic to bees. Avoid using during flowering period of the crop. Do not breathe dust or take internally. Avoid skin contact. Change clothing daily. Avoid contamination of feed, feed troughs and water receptacles. Do not treat within 7 days of harvest except tomatoes and corn which may be treated 1 day before harvest.

FIRST AID:

If swallowed, induce vomiting by touching finger to back of throat. Get medical attention or contact a Poison Control Centre. If in eyes, flush thoroughly with plenty of water for 15 minutes. If on skin, wash with soap and water.

TOXICOLOGICAL INFORMATION:

Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

DISPOSAL:

Do not reuse empty container. Wrap in newspaper and discard with household garbage.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

(R)Registered trademark of Sanex Agro Inc.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))