

----- LABEL -----

Sendran (R)

Collar for Cats of all Sizes

Kills Fleas & Ticks up to 4 Months

For continuous protection replace collar
when effectiveness
diminishes

SOLD THROUGH
VETERINARIANS

CAUTION:
DO NOT ALLOW CHILDREN TO PLAY WITH THIS COLLAR

DOMESTIC

READ THE LABEL BEFORE USING

GUARANTEE: Propoxur.....9.4%

CONTENTS: 1 Collar, Net Weight 8.25 g

Registration No. 17493 P.C.P. Act

Distributed by BAYVET Division
Chemagro Limited
Etobicoke, Ontario, Canada M9W 1G6

HAYER (R)

----- INSERT -----

CODE NO. 2380

SENDRAN (R)

FLEA AND TICK COLLAR
For Cats

DOMESTIC

GUARANTEE:

Propoxur.....9.4%

DIRECTIONS

Remove collar from package and place around cat's neck. Adjust for proper fit and buckle in place. Collar must be worn loosely but securely enough to prevent easy removal and loss. Cut off any excess length of collar, leaving ample length for expansion as the cat grows, and dispose of by discarding in trash. Check and adjust collar periodically to assure a proper fit.

Do not use on kittens under 12 weeks of age.

RESULTS EXPECTED

The flea and tick collar starts killing fleas and ticks when it is placed around the cat's neck. Fleas will be killed and new ones which may temporarily appear on the cat will also be killed for up to 4 months. Non-engorged ticks are tough occasional pests of cats and will be killed within a few days. Collar will kill ticks encountered in first two months of wear. FOR CONTINUOUS PROTECTION, REPLACE COLLAR WHEN EFFECTIVENESS DIMINISHES. Some cats may be sensitive to this collar. When collar is first worn, observe neck area every few days for irritation. Any collar, when fastened too tightly, may cause skin irritation. Remove collar at the first sign of irritation or adverse reaction. This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Do not use on sick or convalescing animals. Do not administer other cholinesterase inhibitors (i.e. oral pesticides) while collar is worn or for a few days after removing collar.

PRECAUTIONS

Do not open inner envelope until ready to use. DO NOT ALLOW CHILDREN TO PLAY WITH UNATTACHED COLLAR. Dust will form on this collar during storage. Do not get dust or collar in mouth, harmful if swallowed. Wash hands thoroughly with soap and water after handling collar.

FIRST AID

In case of poisoning, call a physician immediately. If dust from collar contacts skin, wash thoroughly with soap and water. If dust contacts eyes, flush eyes with water and get medical attention.

TOXICOLOGICAL INFORMATION

Dust in eyes will cause temporary pupillary constriction*. Dust released from collar is a cholinesterase inhibitor. Note to Physician: Atropine and homatropine* are antidotal.

Reg. No. 17493 P.C.P. Act

SENDTRAN (R) is a Reg. TM of the parent Company of
Farbenfabriken Bayer, GmbH, Leverkusen.

MFG FOR:

BAYVET Division
CHEMAGRO Ltd.
Rexdale, Ontario M9W 1G6

HAYER (R)

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))