

96.05.29

SANEX

PROFESSIONAL PRODUCTS
LES PRODUITS PROFESSIONNELS

COMMERCIAL

AN EMULSIFIABLE CONCENTRATE, CONTAINS CHLORPYRIFOS
CONCENTRE EMULSIFIABLE CONTENANT DU CHLORPYRIFOS

FOR USE ONLY BY PEST CONTROL OPERATORS ONLY TO CONTROL NUMEROUS PEST OF
HOUSEHOLDS, TURF, AND SOD FARMS, ORNAMENTAL PLANTS, INDUSTRIAL AND
INSTITUTIONAL LOCATIONS.

STRICTEMENT RESERVE AUX EXTERMINATEURS LICENCIES POUR LA REPRESSION DE
NOMBREUX INSECTES NUISIBLES DANS LES MAISONS, LES PELOUSES, LES GAZONNIERES,
LES PLANTES ORNEMENTALES ET LES ETABLISSEMENTS INDUSTRIELS ET
INSTITUTIONNELS.

READ THE LABEL BEFORE USING
LIRE L'ETIQUETTE AVANT L'EMPLOI

REG.NO./NO.D'ENR. 16740
P.C.P. ACT/L.P.A.

DANGER POISON

GARANTIE/GUARANTEE:
CHLORPYRIFOS 240 g/L

4 L

SANEX AGRO INC.
36 Head St.,
DUNDAS, ONTARIO
L9H 3H3

DIRECTIONS FOR USE

INDOOR PEST CONTROL:

Use FOSBAN 2E insecticide to control cockroaches, carpet beetles, crickets,

ants, firebrats, silverfish, spiders, brown and American dog ticks and fleas in and around residential, institutional and commercial buildings including homes, apartment buildings, hospitals, stores and warehouses. Dilute FOSBAN 2E with water or base oil to make a 0.24% or 0.48% chlorpyrifos spray solution as shown in the following SPRAY DILUTION CHART.

WATER-BASED SPRAY			OIL-BASED SPRAY		
Litres of Spray Mixture desired	mL of FOSBAN 2E to use		Litres of Spray desired	mL of FOSBAN 2E to use	
	0.24% spray	0.48% spray		0.24% spray	0.48% spray
1	10	20	1	8	16
2	20	40	2	16	32
5	50	100	5	40	80
10	100	200	10	80	160
25	250	500	25	210	420

Application within food handling establishments (places other than private residences in which food is held, processed, prepared or served) including but not limited to restaurants, grocery stores, bakeries, bottling plants, canneries and grain mills. IN FOOD PROCESSING, PREPARATION AND SERVING AREAS USE ONLY DURING CLOSE DOWN PERIODS. Use as a spot treatment to control cockroaches (including strains resistant to certain other insecticides), ants, crickets, firebrats, silverfish and spiders in and around both food and non-food areas by application as a coarse, low pressure (20 psi or less) spray to localized areas where the above pests have been seen or are suspected of hiding or entering. Apply only enough spray to thoroughly cover the surfaces treated using special care to avoid unnecessary runoff. Do not introduce the spray into the air or allow the spray to contact food or food contacting surfaces. Areas treated may include dark corners of rooms and closets, along and behind baseboards, beneath and behind stoves, refrigeration units and similar food preparation areas and equipment, floor drains and around plumbing and other utility installations. For ants apply to ant trails and wherever these pests may find entrance. Spot treatment may encompass crack and crevice treatment by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment legs and bases or at points between different elements of construction or between equipment and floors. REPEAT TREATMENT AS NEEDED BUT NOT MORE OFTEN THAN ONCE EVERY 7 DAYS IN RESTAURANTS AND SIMILAR FOOD SERVICE ESTABLISHMENTS OR MORE OFTEN THAN ONCE EVERY 14 DAYS IN OTHER TYPES OF FOOD HANDLING ESTABLISHMENTS. Remove or cover all food, packaging materials and utensils before spraying. After spraying wash all surfaces that may contact food and rinse thoroughly with potable water before re-use for food processing preparation or serving. APPLICATION OF THIS PRODUCT IN FOOD HANDLING ESTABLISHMENTS OTHER THAN AS A SPOT AND/OR CRACK AND CREVICE TREATMENT ARE

NOT PERMITTED.

For control within food and non-food areas of residential buildings including homes and apartment buildings and within industrial, institutional, commercial buildings including hospitals (in non-patient areas), stores, manufacturing plants and warehouses. COCKROACHES, ANTS, CRICKETS, FIREBRATS, SILVERFISH, SPIDERS.

Use a 0.24% solution to control light infestations and a 0.48% solution to quickly reduce heavy infestations. Apply as a spot treatment to localized areas where pests occur or are found, including, dark corners of rooms and closets, cracks and crevices in walls, baseboards, beneath and behind stoves, fridges and cupboards, around plumbing and utility installations. Preferably as a coarse low pressure spray.

CARPET BEETLES

Apply as a 0.48% spray as a spot treatment along baseboards and edges of carpeting, under carpets, rugs or furniture, in closets, on shelves and wherever these insects are seen or suspected.

BROWN AND AMERICAN DOG TICKS & FLEAS

Apply a 0.48% spray as a spot treatment to infested areas such as pet beds and resting quarters; nearby cracks and crevices, along and behind baseboards, window and doorframes, and localized areas of floor and floor covering where these pests may be present. Old pet bedding should be removed and replaced with clean fresh bedding after treatment. DO NOT TREAT PETS WITH THIS PRODUCT. DO NOT APPLY THIS PRODUCT TO FURNITURE.

WARNING:

(PESTS INDOORS) Do not use in poultry houses, other animal buildings or in edible products, areas of food storage or processing plants. Do not allow spray to contact food, feedstuffs and water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow humans or pets to contact treated surfaces until spray has dried. Remove pets and cover fish tanks before spraying. Cover or remove all food before spraying FOSBAN. In hospitals use only in non-patient areas. Do not use in nurseries, nor in any rooms occupied by infants, sick or aged persons.

PESTS IN LAWNS AND RECREATIONAL TURF GRASS:

Use FOSBAN 2E to control ants and chinch bugs. Mix FOSBAN 2E with water following instructions in the chart, apply as a coarse spray uniformly to the lawn using low pressure and suitable application equipment. To obtain best results lawn should be moist at time of application. Spray when pests first appear and repeat as necessary.

DOSAGE & DILUTIONS FOR USE ON TURF

AREA TO BE TREATED	VOLUME FOSBAN 2E	DILUTE WITH WATER TO GIVE
10 m ²	4.0 mL	11 L
100 m ²	42 mL	110 L
500 m ²	210 mL	550 L
1000 m ²	420 mL	1100 L

TO CONTROL WHITE GRUBS, EUROPEAN CHAFER (LARVAE), JAPANESE BEETLE (LARVAE)

Use twice as much FOSBAN 2E as shown in the above Turf Chart, i.e. 8.40 mL diluted in 1,100 litres of water per 1000 square metres of area. For European Chafer(larvae), Japanese Beetle(larvae) and White Grubs(June Beetles), treat when grubs are small and feeding near the soil surface, usually in late July and during August. Consult your local Agricultural Extension Service Specialist for proper time to treat in your area.

NOTE:

To help prevent infestation of buildings by ants, treat a band of soil 2-3 metres wide around and adjacent to the building, also the building foundation to a height of one metre, where pests are active and may find entrance. Use the same dosage and dilution recommendations for control of these pests on turf as shown in the above table. Apply as a coarse spray to thoroughly and uniformly wet the band area.

WARNING

(LAWN PESTS) Keep humans and pets off treated areas until spray has dried. Keep out of lawn and garden fishpools and other bodies of water. Fish may be killed if the waters are contaminated with this product. Do not apply where runoff is likely to occur. Do not apply when weather conditions favour drift from areas treated. Do not mow or collect clippings for 24 hours after treatment.

USE PRECAUTIONS:

WARNING: This product is toxic to fish and crustaceans. Do not contaminate water by cleaning of the equipment or disposal of wastes.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. May be fatal if swallowed. May be absorbed through skin. May be injurious to eyes and skin. Do not take internally. Do not get in eyes, on skin or clothing. Wash thoroughly after handling. Wash contaminated clothing before re-use. Flush contaminated eyes with plenty of

water and get prompt medical attention. Avoid breathing vapours and spray mist. Avoid spraying food crops. Keep away from food, feedstuffs and domestic water supplies. Keep container closed. Keep away from heat and open flame. Handle concentrate in a ventilated area. Do not permit humans or pets to contact treated areas until thoroughly dry. Wash dishes and food handling utensils thoroughly in soap and water if they become contaminated during application of the product.

FIRST AID:

If accidentally swallowed, call a physician or a poison control centre immediately. Do not induce vomiting in an unconscious person. If conscious, induce vomiting to rid stomach of Fosban 2E. Look for secondary signs of pulmonary pneumonia which may be delayed 24 hours or more. Flush contaminated eyes and skin with plenty of water for 15 minutes. Get medical attention for eyes. If on skin, wash thoroughly with soap and water.

TOXICOLOGICAL INFORMATION:

Chlorpyrifos is a Cholinesterase inhibitor. SYMPTOMS OF POISONING: Headache, weakness, tightness in the chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, abdominal cramps. Treat symptomatically. Atropine is antidotal.

NOTICE TO USER:

This product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

