

31-JAN-2006

30-JUN-2006 - Notification of Registrant Name Change. See below for details.

C-I-L[®]
GARDEN DOCTOR⁽¹⁾
INSECTICIDE/FUNGICIDE

Controls insects and diseases on tomatoes, potatoes
and many other vegetables and small fruits.

DOMESTIC

REG. NO. 26445 P.C.P. ACT

GUARANTEE:

Carbaryl 5%

Copper 7%

(from Tri-basic Copper Sulphate)

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

275g (300, 500g)

Spectrum Brands IP Inc.

~~Nu-Gro IP Inc.~~

10 Craig St.,
Brantford, ON
Canada N3R 7J1

Notification Change

DIRECTIONS: Apply a thin uniform coating of dust to both sides of foliage, preferably in early morning or evening when air is still. Start applications when insects or damage caused by insects or disease appear and repeat at intervals of 7 to 10 days or as necessary for control. For potato, apply when plants are 13 to 18 cm high. Repeat at intervals of 5 to 10 days as long as disease threatens. Some leaf injury may occur to carrots and cucumber if applied to tender wet foliage or in the presence of prolonged humidity.

* days interval to harvest

CROP	ITH*	DISEASES	INSECTS
Grape	3	black rot, dead arm, downy mildew	berry moth, leafhoppers
Strawberry	1	leaf spot	spittlebug, leafroller
Bean	3	anthracnose, leaf spot, downy mildew	
Beet, Carrot	14 1	<i>Cercospora</i> leaf spot	
Broccoli, Cabbage, Brussels Sprouts, Cauliflower	3 3 3 3	black leaf spot, grey leaf spot, downy mildew, leaf spot	flea beetles, leafhoppers, armyworms, cabbage loopers, diamondback moth, imported cabbageworm, spittlebug, stink bug
Celery	3	early blight, late blight	
Spinach	14	downy mildew, white rust	
Cucumber Melon Squash	3 3 3	anthracnose, leaf spot, scab	cucumber beetle, climbing cutworm, flea beetle, leafhoppers, squash bugs
Eggplant Pepper Potato	1 1 7	early blight, late blight	Colorado Potato Beetle, corn borer, fall armyworm, tomato horn worm, tomato fruitworm leaf hoppers, flea beetles, stink bugs, plant bugs, climbing cutworm
Tomato	1	anthracnose, early blight, late blight, <i>Septoria</i> leaf spot	
Ornamentals (rose)		black spot, powdery mildew, downy mildew	beetles, leafhoppers, leafrollers, plant bugs, rose aphid, rose chafer, rose slug

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Hazardous if swallowed, inhaled, or absorbed through the skin. Wash thoroughly after use and before eating, drinking or smoking. Do not contaminate feed, food, ponds or streams. Do not use when fruit trees are in bloom, or directly on flower blossoms as it is very poisonous to foraging bees. Store in a cool, dry, ventilated place.

FIRST AID: If on skin, wash thoroughly with soap and water. **If in eyes,** flush with water for 5 to 10 minutes and get medical attention. **If swallowed,** induce vomiting and go to hospital taking labelled container with you. **DO NOT INDUCE VOMITING IF PATIENT IS UNCONSCIOUS OR CONVULSIVE.** Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

DISPOSAL: Dispose of empty container with regular garbage.

For additional product information or in case of emergency, spills or fire, call toll-free 1-800-268-2806. www.cil.nu-gro.com

CIL Oval® ICI Canada Inc. / Lic. Nu-Gro
GARDEN DOCTOR is a trademark of Zeneca Corp.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.