

15-JAN-1998

Later's
Malathion
4% Dust
Insecticide

An INSECTICIDE for use on PETS, ORNAMENTALS and VEGETABLES.

DOMESTIC

READ THE LABEL BEFORE USING.

KEEP OUT OF REACH OF CHILDREN

REG. NO. 14211
P.C.P. ACT

GUARANTEE: Malathion.... 4%

500 g

Notification Change

~~Later Chemicals Ltd.
12080 Horseshoe Way
Richmond, B.C.
V7A 4V5~~

Nu-Gro IP Inc
10 Craig Street
Brantford, ON N3R 7J1

DIRECTIONS FOR USE:

PETS: Limitations - 1) Do not apply to pets under 8 weeks of age.
2) Do not apply to adult animals which are nursing young.

DOGS, CATS: To control Fleas, Lice and Ticks, apply directly to fur so that the dust reaches the skin. Repeat in two weeks if necessary.

ANIMAL QUARTERS (Dogs, Cats): To control Fleas, clear the area to be treated of debris and old bedding. Use 1 kg of dust per 100 m². Provide fresh bedding after treatment.

ORNAMENTAL TREES, SHRUBS, AND FLOWERS: To control Aphids, Bagworms, European pine shoot moths, Lace bugs, Leafhoppers,

Leafminers, Mealybugs, Native scale, Pine needle scale, Spider mites, Tent caterpillars, Thrips, Four-lined plant bugs, and Tarnished plant bugs, apply at first appearance of pest. Obtain full coverage. Repeat as needed.

AFRICAN VIOLET: To control Aphids, Midges, Spider mites, and White flies, dust plants lightly.

VEGETABLES:

ASPARAGUS: To control Asparagus beetle, use at the rate of 330 g per 100 m². Do not apply within 1 day of harvest.

BEANS: To control Aphids, Leafhoppers, Mexican bean beetles, and Spider mites, use at the rate of 330 g per 100 m². Repeat as necessary. Do not apply within 1 day of harvest.

BROCCOLI: To control Aphids, and Cabbage loopers, use at the rate of 330 g per 100 m². Repeat as necessary. Do not apply within 3 days of harvest.

BRUSSEL SPROUTS: To control Aphids, Cabbage Worms and Cabbage loopers, use at the rate of 330 g per 100 m². Do not apply within 7 days of harvest.

CABBAGE, KALE, TURNIPS: To control Aphids, Cabbage loopers, and imported Cabbage worms, use at the rate of 330 g per 100 m². Repeat as necessary. Do not apply within 7 days of harvest.

CARROTS: To control Aphids, use at the rate of 330 g per 100 m². Do not apply within 7 days of harvest.

CAULIFLOWER: To control Aphids, Cabbage loopers, and Cabbage worms, use at the rate of 330 g per 100 m². Do not apply within 7 days of harvest.

CUCUMBER ,SQUASH: To control Aphids, Leafhoppers, and Spider mites, use at the rate of 280 g per 100 m². Do not apply within 3 days of harvest. Do not apply unless the foliage is dry.

FIELD LETTUCE (Leaf, Head): To control Aphids, Cabbage loopers, Mites, and six-spotted Leafhoppers, use at the rate of 330 g per 100 m². Repeat as necessary. Do not apply within 3 days of harvest on field grown head lettuce and within 14 days of harvest on field grown leaf lettuce.

ONIONS (Bulb, Green), GARLIC: To control Thrips use at the rate of 280 g per 100 m². Repeat as needed. Do not apply within 3 days of harvest.

PEAS: To control Leafhoppers, Mites, Pea aphids, Pea moths, and Pea weevils, use at the rate of 280 g per 100 m². Repeat as necessary. Do not apply unless foliage is dry. Do not apply within 3 days of harvest.

POTATOES: To control Aphids, Colorado potato beetles, Leafhoppers, and Mites use at the rate of 250 g per 100 m². Repeat as necessary. Do not apply within 3 days of harvest.

RADISHES: To control Aphids, Flea beetles, and Leafhoppers, use at the rate of 330 g per 100 m². Repeat as necessary. Do not apply within 7 days of harvest.

TOMATOES: To control Aphids, Leafhoppers, and Spider mites, use at the rate of 250 g per 100 m². Repeat as necessary. Do not apply within 3 days of harvest.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN and DOMESTIC ANIMALS. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash exposed skin areas thoroughly with soap and water after working with malathion. Clean contaminated clothing before re-use. Do not treat dairy barns. Wash treated food before eating. Do not apply to eyes, ears or nose of any animal.

ENVIRONMENTAL HAZARDS: Highly toxic to fish, and bees. Do not treat any plants while in bloom. Avoid contaminating any body of water.

FIRST AID: In case of exposure to malathion, move patient from area and remove all contaminated clothing. Wash all exposed skin with soap and water. IF IN EYES flush with water for 15 minutes and get medical attention. IF SWALLOWED, induce vomiting by inserting a finger into mouth and touching the back of the throat. Never give anything by mouth to an unconscious person. In case breathing is weak or irregular, give artificial respiration. Contact a doctor or poison control centre at once.

TOXICOLOGICAL INFORMATION: Symptoms of poisoning; headache, weakness, sweating, giddiness, blurred vision, nausea, abdominal cramps, diarrhea, and discomfort in the chest. Atropine is antidotal. Do not use 2-PAM for malathion poisoning.

STORAGE:

Containers should be kept tightly sealed when not in use. Do not store near food, feed, fertilizer, or seed.

DISPOSAL: Do not re-use empty container. Wrap container and discard in household garbage.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))