AGRICULTURAL

MICROFINE SULPHUR 92

FUNGICIDE

For Apples, Peaches, Plums, Cucumbers and Ornamentals

READ THE LABEL BEFORE USING KEEP OUT OF REACH OF CHILDREN

GUARANTEE: sulphur 92%

REGISTRATION No. 3918 PEST CONTROL PRODUCTS ACT

ZENECA Agro a business of ZENECA Corp. Stoney Creek, Ontario L8G 3Z1

A93/A-1

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

- * Avoid Breathing Dust or Spray Mist.
- * Avoid Contact with Skin.
- * Wear Gloves.
- * Wash Hands Thoroughly After Use.
- * Do not contaminate food or feed.

FIRST AID

If a known exposure occurs or is suspected, immediately initiate the recommended procedures below. If further treatment is required, contact Poison Control Centre, a physician, or the nearest hospital.

If Swallowed: Immediately give several glasses of water and induce

15 kg

vomiting by gagging the victim with a finger or blunt object placed on the back of the victim's tongue. Give fluids until vomitus is clear. If victim is unconscious, do not induce vomiting or give anything by mouth.

If in Eyes: Immediately flush eyes with large amounts of running water f at least 15 minutes. Hold eyelids apart during flushing to ensure rinsir of the entire surface of the eye and lids with water. Get medical attention.

If on Skin: Flush all affected areas with plenty of water for 15 minutes Remove contaminated clothing and shoes. Seek medical attention if skin irritation occurs. Wash clothing before reuse.

If Inhaled: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

EMERGENCY TELEPHONE NUMBER

All hours, 1-800-263-0984 ONLY for health and environmental information.

STORAGE

Store in a cool dry well ventilated area away from foodstuffs and out of reach of children and animals. This product is not affected by freezing.

SPILL CLEANUP

Wear Appropriate protective equipment (gloves, glasses, apron) when attempting to clean up the spill. If the container is leaking, secure le and place the container into a drum or heavy gauge plastic bag. Contact ZENECA Agro (See EMERGENCY NUMBER) for further information.

For spills and leaks; clean up as much of the product as possible with a broom and shovel. If available, use an absorbent such as DUSTBANE to $cl\epsilon$ the spill area. Shovel the spilled product and spend absorbent in sealak waste container.

<u>On hard surfaces</u> - sprinkle spill area with detergent and scrub in a smal quantity of water with a coarse broom. Let stand for 10 minutes then absorb onto an inert material and shovel waste into a waste container.

<u>On soil</u> - remove the top 15 cm of soil in the spill area and replace with fresh soil. Dispose of all waste including scrub brush in accordance wit provincial requirements. For more information on the disposal of waste ϵ the clean up of spills, contact the regional office of Conservation and Protection, Environment Canada.

DISPOSAL

Rinse the empty container thoroughly and add the rinsings to the spray mixture in the tank. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the emp container unsuitable for further use. Dispose of container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office c Conservation and Protection, Environment Canada.

PRODUCT INFORMATION

Chipman Microfine Sulphur 92 is a high quality wettable sulphur for spray or dust application. It is compatible with the commonly used insecticide and fungicides except oil.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to u a control product under unsafe conditions.

DIRECTIONS FOR USE

Sulphur should not be used in combination with, immediately before or immediately after an oil spray. Sulphur sprays should not be applied whe temperature is above 30EC.

CROPS	DISEASES	RATE	REMARKS
Apples	Apple Scab	20 to 27 kg/ha	In pre-pink, blossom and calyx sprays, use 27 kg/ha In cover sprays use 20 kg/ha.
Peaches Sweet Cherries and Plums	Brown Rot Scab Leaf Spot	20 to 27 kg/ha	Use in all sprays where sulphur is recommended.
Sour Cherries	Brown Rot	20 to 27 kg/ha	Use in spray just before blossoms open. For other sprays use ZENECA Ferbam.
Gooseberries	Powdery Mildew	27 kg/ha	Do not use on "sulphur-shy American varieties.
Roses, Chrysanthemums and other Ornamentals	Powdery Mildew	20 kg/ha	Apply on appearance of disease.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arisin from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulator Agency to provide efficient searching for label information. This servic and this information do not replace the official hard-copy label. The PN does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore no liable for any loss resulting, directly or indirectly, from reliance upor this service.

+))