

30-JAN-2003. Notification of Label Changes received. See below for label changes.

16-DEC-1998

((BAG))

MORESTAN (R)
SOLUPAK (TM)

25% WETTABLE POWDER
MITICIDE, FUNGICIDE

COMMERCIAL

In Convenient Water Soluble Packets

NET WEIGHT 2 KG

contains four 500 gram water soluble packets

FOR CONTROL OF MITES AND POWDERY MILDEW

GUARANTEE: Chinomethionat.....25%

REGISTRATION NO. 21,367
PEST CONTROL PRODUCTS ACT

KEEP OUT OF REACH OF CHILDREN

KEEP WATER SOLUBLE PACKETS IN THIS CONTAINER
AND STORE IN A COOL DRY PLACE.

DO NOT ALLOW PACKETS TO BECOME WET
PRIOR TO ADDING TO THE SPRAY TANK.

DANGER - EYE IRRITANT

READ THE LABEL AND PRINTED DIRECTIONS BEFORE USING

(See side panel for caution statements)

FOR COMPLETE DIRECTIONS FOR USE
SEE MANUFACTURER'S PRINTED DIRECTIONS ON BOTTOM OF BAG

WATER SOLUBLE PACKETS MAY NOT BE SOLD SEPARATELY

Notification Change

~~BAYER INC.~~

Bayer CropScience Inc. □
Suite 100 □
3131-114th Ave. S.E. □
Calgary, AB T2Z 3X2

~~CROP PROTECTION~~
~~77 BELFIELD ROAD~~
~~ETOBICOKE, ONTARIO,~~
~~CANADA M9W 1G6~~

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

ENVIRONMENTAL HAZARDS

MORESTAN is toxic to fish. Do not apply where runoff is likely to occur. In cleaning of equipment or disposal of wastes, do not contaminate streams, lakes or ponds with this material.

DANGER

Causes substantial eye injury. Harmful if inhaled. Avoid breathing dust or spray mist. Do not get in eyes or on clothing. WEAR GOGGLES OR FACE SHIELD. Wear latex or Neoprene gloves and rubber apron. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash separately from household laundry before reuse.

Do not contaminate feed or food. KEEP OUT OF REACH OF CHILDREN.

FIRST AID: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. **If swallowed**, vomiting should be induced. Administer water freely and induce vomiting by giving one dose (15 mL) of syrup of ipecac. If vomiting does not occur with a 10 to 20 minutes, administer second dose.

If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. The patient should be lying down with the head below the foot level and facing down or to one side. Professional medical assistance should be secured immediately. DO NOT INDUCE VOMITING TO AN UNCONSCIOUS PERSON OR TO PERSONS IN A CONVULSIVE STATE. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash immediately with flowing water for at least 15 minutes.

TOXICOLOGICAL INFORMATION: In experimental animals acutely poisoned by MORESTAN, diarrhea and decreased activity were the symptoms observed.

the patient should be treated symptomatically.

DISPOSAL

1. Do not reuse the container.
2. Dispose of the empty container in accordance with provincial requirements.
3. For information on the disposal of unused, unwanted product and the cleanup of spills contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto the buyer assumes the risk to persons or property arising from the use or handling of this product on that condition.

Bayer Inc. Crop Protection 77 Belfield Road, Etobicoke, Ontario, Canada
M9W 1G6

MORESTAN is a Reg. TM of Bayer AG, Germany
SOLUPAK is a Reg. TM of Chemagro Limited

((PAMPHLET))

MORESTAN (R)
SOLUPAK (TM)

25% WETTABLE POWDER
MITICIDE, FUNGICIDE

In Convenient Water Soluble Packets

COMMERCIAL

CONTAINS FOUR 500 GRAMS OF WATER SOLUBLE
PACKETS TOTAL NET CONTENTS 2 KG

FOR CONTROL OF MITES AND POWDERY MILDEW

GUARANTEE: Chinomethionat.....25%

REGISTRATION NO. 21,367
PEST CONTROL PRODUCTS ACT

KEEP OUT OF REACH OF CHILDREN

KEEP WATER SOLUBLE PACKETS IN THIS CONTAINER
AND STORE IN A COOL DRY PLACE.

DO NOT ALLOW PACKETS TO BECOME WET
PRIOR TO ADDING TO THE SPRAY TANK.

DANGER - EYE IRRITANT

READ THE LABEL BEFORE USING
(See rear panel for first aid and danger statements)

WATER SOLUBLE PACKETS MAY NOT BE SOLD SEPARATELY

Notification Change

~~BAYER INC.~~
~~CROP PROTECTION~~
~~77 BELFIELD ROAD~~
~~ETOBICOKE, ONTARIO,~~
~~CANADA M9W 1G6~~

Bayer CropScience Inc. □
Suite 100 □
3131-114th Ave. S.E. □
Calgary, AB T2Z 3X2

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

DIRECTIONS FOR USE

MORESTAN miticide-fungicide is recommended for residual control of mites, mite eggs and pear psylla on deciduous fruits, non-bearing strawberries, and ornamentals. MORESTAN SOLUPAK mixes readily with water to form a suspension which is usable in all conventional hand or machine-operated sprayers. For best results, apply a thorough coverage before mites or mite eggs are present in large numbers and before mites cause bronzing of the foliage. Apply during good drying conditions (See Restrictions). Consult your local Agricultural Authorities for detailed information on timing of applications for most effective control.

MIXING: The enclosed packets containing MORESTAN SOLUPAK miticide-fungicide are water soluble. Do not allow packets to become wet prior to adding to the spray tank. Wear latex or Neoprene gloves when handling packets. Reseal carton to protect remaining packets.

To prepare the spray mixture, merely drop the required number of unopened packets, as determined under "Recommended Applications", into the spray tank while filling with water to the desired level. Operate the agitator while mixing. Depending on the water temperature and the degree of agitation, the packets should be completely dissolved within approximately 5 to 10 minutes from the time they were added to the water.

CAUTION: Do not use packets in liquid fertilizer. Do not use packets in a tank-mix with products that contain Boron or release free chlorine. The resultant reaction of water soluble packets and Boron or free chlorine is a plastic which is not soluble in water or solvents such as diesel oils, kerosene, gasoline or alcohol.

DOSAGE: Use specified dosage of MORESTAN SOLUPAK in the amount of water necessary to give complete coverage of foliage. Determine the total amount of wettable powder to be added to the spray tank based on the rates under "Recommended Applications". Add the required number of 500 gram packets to the spray tank while filling with sufficient water to give complete coverage of foliage. The type of equipment used will determine the concentration required; however, do not use these packets for making highly concentrated mixtures such as used for aircraft spraying.

RECOMMENDED APPLICATIONS

CROP	PEST	Dosage MORESTAN SOLUPAK	REMARKS
BEARING FRUIT Pre-Bloom & Post Harvest Apples Apricots Cherries Peaches Pears Prune Plums	Mites Mites eggs	625 g to 1.25 kg/1000 L	Apply specified dosage per 1000 L of water in early season sprays as required. (delayed dormant, pre-pink or pink). Additional applications may be made to tree foliage after harvest.
	Pear psylla	2 kg/1000 L	

NON-BEARING FRUITS Apples	Powdery mildew Mites Mites eggs	625 g to 1.25 kg/1000 L	Apply specified dosage per 1000 L of water as a full coverage spray*. Spray all foliage surfaces including the underside of leaves for complete coverage. Repeat as necessary. For control of powdery mildew, begin preventive applications before disease appears and apply at 7 to 14-day intervals. During heavy infection periods, make more frequent applications.
Apricots Cherries Pears Plums Prune Plums	Mites (except cyclamen mite) Mite eggs	625 g to 1.25 kg/1000 L	
NON-BEARING FRUITS Strawberries	Mites (except cyclamen mite)	1.1 1.75 kg/ha	Apply specified dosage per hectare in sufficient water for complete coverage. Repeat as necessary.
ORNAMENTALS Roses Arborvitae Juniper Yew	Mites Mite eggs	625 g to 1.25 kg/1000 L	Apply specified dosage per 1000 L of water (1g/L). Spray all foliage surfaces including the underside of leaves for complete coverage. Repeat as necessary.

* This concentration is calculated for conventional hydraulic-type sprayers. When lower volumes of spray are applied per hectare as with low-pressure, low-volume, or mist-type equipment, increase the concentration of MORESTAN SOLUPAK in the spray mixture in order to apply amount of MORESTAN SOLUPAK per hectare equivalent to a full coverage spray.

RESTRICTIONS

Do not apply oils after delayed dormant where MORESTAN SOLUPAK is to be used in the pink application.

Do not use spreader-sticker materials with MORESTAN SOLUPAK.

MORESTAN applied within 3 days following a frost or near frost weather conditions and prior to a long wetting period may result in russetting. Cultivars such as MacIntosh, Empire, Spartan and Idared appear to be more susceptible.

Do not use on other crops used for food and forage. Do not allow sprays to drift onto other crops, including pastures. Do not graze treated areas.

Use only according to label directions.

Under certain adverse weather conditions, particularly those that cause slow drying of spray deposits, injury may occur to foliage of strawberries, and deciduous fruit trees (apricots, apples, cherries, peaches, pears, plums, prune plums) that have new and/or tender growth.

Phytotoxicity has occurred on the foliage and flowers of several varieties of roses, especial during periods of high temperatures. Treat roses only after cutting and before new growth appears. Before treating large plantings for the first time, treat a few plants and observe for 4 to 5 days. Under certain conditions, some injury to other ornamentals may occur.

ENVIRONMENTAL HAZARDS:

MORESTAN is toxic to fish. Do not apply where runoff is likely to occur. In cleaning of equipment or disposal of wastes, do not contaminate streams, lakes or ponds with this material.

DANGER

Causes substantial eye injury. Harmful if inhaled. Avoid breathing dust or spray mist. Do not get in eyes or on clothing. WEAR GOGGLES OR FACE SHIELD. Wear latex or Neoprene gloves and rubber apron. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash separately from household laundry before reuse. Do not contaminate feed or food. Keep out of reach of children.

FIRST AID: In case of poisoning call a physician or contact a poison control centre immediately. Have patient lie down and keep quite. If swallowed, vomiting should be induced. Administer water freely and induce vomiting by giving one dose (15 mL) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. The patient should be lying down with the head below the foot level and facing down or to one side. Professional medical assistance should be secured

immediately. DO NOT INDUCE VOMITING TO AN UNCONSCIOUS PERSON OR TO PERSONS IN A CONVULSIVE STATE. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash immediately with flowing water for at least 15 minutes.

TOXICOLOGICAL INFORMATION: In experimental animals acutely poisoned by MORESTAN, diarrhea and decreased activity were the only symptoms observed. The patient should be treated symptomatically.

DISPOSAL:

1. Do not reuse the container.
2. Dispose of the empty container in accordance with provincial requirements.
3. For information on the disposal of unused, unwanted product and the cleanup of spills contact the provincial regulatory agency or the manufacturer.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

MORESTAN is a Reg. TM of Bayer AG, Germany.
SOLUPAK is a Reg. TM of Chemagro Limited

Notification Change

~~BAYER INC.~~
~~CROP PROTECTION~~
~~77 BELFIELD ROAD~~
~~ETOBICOKE, ONTARIO~~
~~M9W 1G6~~

Bayer CropScience Inc. □
Suite 100 □
3131-114th Ave. S.E. □
Calgary, AB T2Z 3X2

190201

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))