

85.12.31

DU PONT
CANADA
BENLATE(R) T

FUNGICIDE

Benomyl - Thiram Seed Dressing
For Dry, Slurry or Drill - Box Treatment of Rape Seed

COMMERCIAL

READ THE LABEL BEFORE USING

GUARANTEE: Benomyl +20%
Thiram20%

REGISTRATION NO. 14,841 PEST CONTROL PRODUCTS ACT

WARNING POISON

PRECAUTION: Keep out of reach of children

MAY IRRITATE EYES, NOSE, THROAT & SKIN

Avoid breathing dust or spray mist.
Avoid contact with skin, eyes, and clothing
In case of contact, flush skin or eyes with plenty of water,
for eyes, get medical attention.

Wash thoroughly after handling and before eating, drinking or
smoking.

Do not consume alcoholic beverages immediately before or within
24 hours after working with this product.

When treating or handling treated seed, work in well ventilated
area and wear a suitable dust mask, goggles and gloves.

Do not contaminate food or feed.

Treated seed may be harmful to birds if ingested. Highly toxic
to fish.

Do not contaminate any body of water with this material.

IMPORTANT - Never allow "Benlate" T to become wet during storage.
This may lead to certain chemical changes which will reduce the
effectiveness of "Benlate" T as a fungicide. Keep container
closed when not in use. Do not re-use container, bury when
empty.

Manufactured by E.I. DU PONT DE NEMOURS & COMPANY (INC.),

Agricultural Products,
Wilmington, Delaware, U.S.A.

Distributed in Canada by:
DU PONT CANADA INC. Agricultural Products,
P.O. Box 2300, Streetsville Post Office,
Mississauga, Ontario
L5M 2J4

"Benlate" Du Pont Reg. Trademark in Canada
+Canadian 1969 Pat. No. 820317

DIRECTIONS

Du Pont "Benlate" T Fungicide should be used only in accordance with recommendations on this label, or in separate Du Pont Agricultural Bulletins.

"Benlate" T is recommended for the treatment of Rape Seed for the control of Black Leg (*Leptosphaeria Maculans* and *Plendodomus Lingam*), seed decay, seedling blight and damping-off.

HOW TO USE: Apply 310 g "Benlate" T to 25 kg of Rape seed by the following methods:

Dry or Slurry Treatment - Apply after cleaning and before bagging.

GUSTAFSON TREATERS - Insure that the treater is properly calibrated to apply the rate selected for application.

FEED OR CEMENT MIXERS - Fill unit not more than 1/2 full of seed. Add the correct amount of "Benlate" T for the weight of seed in the mixer. Rotate slowly until gray colour is imparted to all the seed.

Drill Box Treatment - Place 1/2 the seed in the drill box, do not exceed 1/2 the capacity, and sprinkle 1/2 the total amount of "Benlate" T to be used over the seed surface. Mix by stirring with a stick or paddle until gray colour is imparted to all the seed. Add remaining 1/2 of seed and "Benlate" T and stir as outlined above.

Note: Label all unplanted treated seed as follows:

"Poisonous to man and animals. This seed has been treated with "Benlate" T fungicide for the control of disease. Do

not use as food or feed. Do not sell to oil mills".

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))