

86.07.28

NOXALL

DOMESTIC

PROLIN
RAT BAIT MEAL

KILLS RATS AND MICE

CONTAINS: WARFARIN AND SULFAQUINOXALINE

READ LABEL BEFORE USING

GUARANTEE:
WARFARIN - 0.025%,
SULFAQUINOXALINE - 0.025%

"WISCONSIN ALUMNI RESEARCH FOUNDATION
CANADIAN LETTERS PATENT NO. 460,333"

REG.NO. 03758 P.C.P. ACT

227g

NOXALL PRODUCTS LTD.
1820 W. 3RD AVENUE,
VANCOUVER, B.C., CANADA

DIRECTIONS FOR USE:

FOR RATS: Place two to eight teaspoons of contents of this package in locations frequented by rats, or where rats have been known to feed. Bait must be protected from children, dogs, cats, poultry, livestock, etc. by bait boxes, cages or other suitable means. An uninterrupted supply of bait should be maintained for at least ten days. Baits should be examined regularly and replenished before completely eaten. If bait moulds, decomposes or becomes sour it should be replaced. Where a continuous source of infestation prevails from nearby dumps or fields, permanent bait stations should be used and bait replenished as needed.

Baiting should continue until complete lack of feeding is noted.

FOR MICE: For controlling mice follow same general directions as for rats except that the amount of bait replacements should be tablespoon amounts to be placed at 3 m intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

PRECAUTION: This product contains as its active ingredient an anticoagulant chemical which if taken accidentally by humans, domestic animals or pets, may reduce the clotting ability of the blood and serious hemorrhage may result. Keep out of reach of children and domestic animals. Wash hands after using. Do not contaminate feed or foodstuffs.

FIRST AID: In case baits are accidentally eaten, give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Call a Physician immediately.

TOXICOLOGY: When a human has been known to have accidentally ingested Noxall Rat Bait containing prolin, warfarin and sulfaquinoxaline, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated, as in the case of hemorrhage caused by overdoses of Dicumarol R.

READY MIXED FOR INSTANT USE.

KEEP IN A DRY PLACE.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))