

05-MAY-1997

Net Contents: 600 g

CO-OP (R)  
D-L+C DRILL BOX

Seed Treatment Powder  
Insecticide-Fungicide

CORN, BEANS, SOYBEANS AND PEAS:

For the control of Wireworms and Seed Corn Maggots and to supplement previous fungicidal treatment for seedling blight and seed rot.

DANGER                      POISON

DANGER                      CAUTION

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION: Exposure to captan may produce long term health effects. To minimize exposure, follow directions outlined on this label.

NOTE: This product contains 0.6% Rhodamine B Dye which has been determined to be of toxicological concern.

GUARANTEE: diazinon 15%  
gamma BHC (from Lindane) 25%  
captan 15%

REGISTRATION NO. 11451 PEST CONTROL PRODUCTS ACT

INTERPROVINCIAL COOPERATIVE LIMITED  
P.O. BOX 1050,  
SASKATOON, SASKATCHEWAN  
S7K 3M9

WINNIPEG              MISSISSAUGA              MONTREAL              MONCTON

## PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Never handle material with bare hands. Do not breathe dust. Avoid spilling on skin or clothing.

When treating or handling treated seed, work in a well-ventilated area, and wear a respirator, goggles, and rubber gloves. Do not use leather or cloth gloves. Wear clothing that completely covers arms and legs to minimize exposure. Change contaminated clothing daily and wash thoroughly before reuse. Wash hands, face and arms thoroughly after handling CO-OP D-L+C SEED TREATMENT and before drinking, eating or smoking. Shower after use or at the end of the work day. Do not contaminate food, feed or any body of water. Lindane is toxic to fish, birds and other wildlife. Label unplanted treated seed as follows: "Poisonous to man and animals. Do not use for food or feed purposes. Do not sell to oil mills. This seed has been treated with captan, diazinon and lindane."

## FIRST AID

In case of poisoning, call physician or contact a Poison Control Centre. IF ON SKIN, wash thoroughly with soap and water. IF IN EYES, flush with water for 5-10 minutes and get medical aid. IF SWALLOWED, give the patient one to two glasses of water and induce vomiting by either giving a dose of syrup of ipecac or inserting a finger down the throat. Repeat ipecac dose if unsuccessful within 20 minutes. Continue with water until vomit is clear. Patient should be lying down with the head below the level of feet while vomiting. DO NOT TRY TO CAUSE VOMITING IF THE PATIENT IS UNCONSCIOUS OR IN A CONVULSIVE STATE.

## TOXICOLOGICAL INFORMATION

Diazinon is a cholinesterase inhibitor. Atropine is antidotal. Lindane is an organochlorine insecticide. Barbiturates may be given to control convulsions. Oxygen may be indicated. Keep patient quiet. Avoid use of morphine and adrenaline. Symptoms of Poisoning: nausea, vomiting, hyper-irritability, convulsions and/or coma and other symptoms typical to organochlorine or organophosphate insecticide poisoning.

## DISPOSAL:

1. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.

3. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

#### NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

#### DIRECTIONS FOR USE

USE THIS PRODUCT ONLY ON SEED PREVIOUSLY TREATED WITH CAPTAN OR THIRAM. DO NOT USE ON SEED ALREADY TREATED WITH AN INSECTICIDE (other than methoxychlor or malathion).

CORN, BEANS, SOYBEANS AND PEAS: For the control of Wireworms and Seed Corn Maggots and to supplement previous fungicidal treatment for Seeding Blight and Seed Rot, apply the contents of one 600 g container to 300 kg (12 units) of seed at planting or near planting time (50 g per 25 kg of seed). Use the scoop to measure 50 g. When starting with an empty planter, place enough treated seed in the planter box to cover the bottom. Treat this seed in a separate container that later can be destroyed. Add the remaining seed to the planter box and treat by thoroughly stirring with a paddle or lath so that the seed is coated with the seed treatment.

#### NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

(cap)

Cap measures 50 g to the ridge.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace

the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))