

95/2000

CLEAN CROP®

ATRAZINE 80W

HERBICIDE

COMMERCIAL

READ THE LABEL BEFORE USING

GUARANTEE:

ATRAZINE 76%
Related Triazines 4%

REGISTRATION NO. 14524
PEST CONTROL PRODUCTS ACT

Wettable Powder Herbicide for Selective
Weed Control in corn, triazine tolerant
Canola and lowbush blueberries.

UNITED AGRI PRODUCTS
R.R. #2
Dorchester, Ontario
N0L 1G5

NET: 4 kg

GENERAL INFORMATION: ATRAZINE 80W provides selective control of broadleaf weeds and wild oats in corn and certain broadleaf weeds in triazine tolerant canola and low bush blueberries. This product may be applied before or after weeds emerge.

WEEDS CONTROLLED:

Wormseed Mustard
Wild Mustard
Common Purslane
Ragweed
Smartweed

Wild Buckwheat
Lamb's-quarters
Redroot Pigweed
Volunteer Clover
Wild Oats

Lady's-Thumb

NOTE: Some naturally occurring triazine tolerant biotypes of the above listed broadleaf weeds may not be controlled by atrazine alone. Refer to specific recommendations in the tank mix section of this label or consult your local agricultural authority.

TIMING OF APPLICATION:

Can be applied:

Pre-plant incorporated (PPI)
Pre-emergence
Post-emergence

in either conventional tillage or minimum tillage programs. If weeds escape a ppi or pre-emergence application, an additional post-emergence application may be necessary. Do not make more than one reapplication per year and only in the spring.

CAUTION: Do not graze or cut for fodder, treated immature crops. (In the case of corn, immature means before ear emergence) there are not sufficient data available to support such use.

LIQUID FERTILIZERS (CORN):

For pre-plant incorporated and pre-emergence application, nitrogen solutions or complete Liquid Fertilizers may replace the water as a carrier.

CAUTION: Do not apply after the corn has emerged as there is danger of Liquid Fertilizer causing crop injury. If there is any doubt about compatibility, check by mixing a small amount of ATRAZINE 80W with the Liquid Fertilizer.

DRY BULK GRANULAR FERTILIZERS:

Many dry bulk granular fertilizers may be impregnated with ATRAZINE 80W or ATRAZINE 80W plus DUAL® as a pre-plant incorporated application to control weeds in corn. Prepare the herbicide/fertilizer mixtures by using any commercially used dry bulk fertilizer blender. Spray the herbicide(s) uniformly onto the fertilizer. If the herbicide/fertilizer mixture is too wet, the use of a highly absorptive powder such as Microcel E, diatomaceous earth or finely powdered clay is recommended. Generally less than 2% by weight of absorptive powder will be needed. For best results, apply ATRAZINE 80W or ATRAZINE 80W plus DUAL/fertilizer mixtures uniformly to the soil with properly calibrated equipment immediately after blending and incorporate

according to label directions.

NOTE: Apply a minimum of 225 kg per hectare of dry granular fertilizer impregnated with recommended rates of herbicide. Do not impregnate ATRAZINE 80W or ATRAZINE 80W plus DUAL onto nitrate fertilizers, super phosphates or limestone.

ATRAZINE 80W In Oil-Water Emulsions (Post-Emergent Treatment):

The post-emergent activity of ATRAZINE 80W is increased when applied in an oil-water emulsion. This combination produces satisfactory control of seedling broadleaf weeds up to 10 cm high. Always apply ATRAZINE 80W before the crop reaches 30 cm in height in order to favour degradation during the growing season. The combination of ATRAZINE 80W in oil-water emulsions has been particularly useful following periods of hot, dry weather, when weeds are more difficult to kill.

NOTE: Some injury to corn can occur with ATRAZINE 80W applied in oil-water emulsions, especially during cold, backward weather. There is no risk of corn injury when ATRAZINE 80W is applied in water alone. Do not add hormone-type weed killers such as 2,4-D, 2-4-DB, MCPA, MCPB and dicamba, or other herbicides, to combinations of oil-water emulsions and ATRAZINE 80W. The risk of damage to other vegetation from spray drift is increased with the use of ATRAZINE 80W in an oil-water emulsion.

MIXING AND SPRAYING INSTRUCTIONS:

This product should not be mixed/loaded, or used within 30 metres of all wells, lakes, streams, ponds or sink holes in order to avoid the potential of well or surface water contamination. Where possible, use a nurse tank to fill your spray tank.

Mixing steps:

1. Spray equipment should be thoroughly flushed with clean water before mixing ATRAZINE 80W.
2. Fill spray tank 1/2 full with carrier. Engage gentle agitation.
3. Pour in the required amount of ATRAZINE 80W or wettable granule, if applicable.
4. Fill tank 3/4 full and add flowable liquid products if applicable.
5. Fill tank 9/10 full and add emulsifiable concentrate, liquid solutions or emulsifiable oils, as required. If adding emulsifiable oils, avoid excessive agitation to prevent

RESTRICTIONS.

** Rate of *XA OIL CONCENTRATE or equivalent depends on water volume used.

Use 300 L water/ha at higher rate

1. DO NOT APPLY TO SWEET CORN.
2. DO NOT USE SUSPENSION FORMULATIONS OF BLADEX (CYANAZINE) POST-EMERGENCE DUE TO RISK OF CROP INJURY.

CAUTION: Do not mix oils, oil concentrates or surfactants or hormone-type herbicides with ATRAZINE 80W/BLADEX.

* ALWAYS APPLY ATRAZINE 80W BEFORE THE CORN REACHES 30 CM IN HEIGHT.

* SOME INJURY TO CORN CAN OCCUR WITH ATRAZINE 80W APPLIED IN OIL-WATER EMULSIONS UNDER CROP STRESS CONDITIONS.

* USE ONLY CROP OILS OR OIL CONCENTRATES LABELLED FOR USE WITH ATRAZINE 80W.

* OILS CONTAMINATED WITH WATER OR OTHER MATERIALS CAN CAUSE COMPATIBILITY PROBLEMS, CROP INJURY, OR BOTH.

WEED CONTROL IN TRIAZINE TOLERANT CANOLA - EASTERN CANADA ONLY:

NOTE: Use ATRAZINE 80W on Triazine Tolerant Canola: pedigree seed only.

WEEDS CONTROLLED:

Non-triazine resistant varieties of:

Wild mustard

Lamb's-Quarters

APPLICATION TIMING:

ATRAZINE 80W may be applied as a PRE-EMERGENCE application or as a POST-EMERGENCE application in an oil-water emulsion when the Triazine Tolerant Canola is at the 2-leaf stage (3 cm in height).

APPLICATION RATES:

Pre-Emergent:

SOIL TYPE

RATES PER HECTARE

Light and Sandy 1.4 kg
Loam and Clay 1.9 kg

Post-Emergent:

Apply 0.9 kg of ATRAZINE 80W plus 17 litres of an emulsifiable crop oil (CORN OIL) in 150 - 300 L of water per hectare.

CAUTION: WHERE PLANTING OF TRIAZINE TOLERANT CANOLA IS DELAYED DUE TO INCLEMENT WEATHER CONDITIONS, OR, IF THIS CROP IS GROWN IN LOW HEAT UNIT AREAS, CHECK FOR RESIDUE CARRYOVER PRIOR TO ROTATING TO SUSCEPTIBLE CROPS.

WEED CONTROL, IN LOWBUSH BLUEBERRIES

WEEDS CONTROLLED:

Browntop	Canada goldenrod*
Poverty grass	Narrowleaf goldenrod*
Sheep fescue	Solidago bicolor*
Hawkweed	Wild strawberry

*suppression only

APPLICATION RATES: Apply 5 kg ATRAZINE 80W per hectare.

TIMING OF APPLICATION:

Apply to native stands in the spring after burn and before new blueberry growth has emerged. CAUTION: Do not apply ATRAZINE 80W to steep terrain where excellent weed control will result in soil erosion. Do not apply ATRAZINE 80W to coarse textured soils low in organic matter.

WARNING:

Injury may occur if land treated with ATRAZINE 80W is planted to any crop other than corn or triazine tolerant canola in the same season. One year after application there is virtually no hazard to most rotational crops when annual weed label rates are applied. Sugar beets and tobacco are sensitive to ATRAZINE 80W and should not be planted on land treated with ATRAZINE 80W the previous year. When extended periods of dry weather occur during the year of treatment, there can be some injury to succeeding crops such as white beans, onions, peas, tomatoes and turnips. Injury is most likely to occur when the seedling crop is subjected to periods of stress e.g., during periods of abnormally hot, dry weather. Under conditions of heavy rainfall sufficient

to cause erosion, soil containing ATRAZINE 80W may wash to lower areas of land and injury to existing or subsequent crops other than corn may result.

THOROUGH TILLAGE INCLUDING PLOUGHING SHOULD PRECEDE THE PLANTING OF CROPS OTHER THAN CORN. PLOUGH THE LAND FLAT AS OPPOSED TO SETTING THE FURROW ON EDGE. UNEVEN APPLICATION OF ATRAZINE 80W OR APPLICATION IN EXCESS OF RECOMMENDED RATES WILL NOT INJURE CORN BUT MAY RESULT IN THE INJURY OF SUCCEEDING CROPS OTHER THAN CORN.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with eyes or prolonged contact with skin. Avoid inhalation of vapours or spray mist. When applying the product wear a coverall over a long-sleeved shirt and pants and chemical resistant boots. Additionally, wear chemically resistant gloves, when servicing the sprayer and during cleanup. Persons involved in mixing/loading operations should wear the previously mentioned clothing, chemical resistant gloves and a face shield. Do not eat, drink or smoke when using product. Wear freshly laundered clothes daily. Wash hands and face before eating, drinking, smoking or using the toilet. Store and wash all protective clothing separately from household laundry. Wash in detergent and hot water before reuse. This product should not be mixed/loading within 30 meters of any wells, lakes, streams, ponds or sink holes in order to avoid the potential of well or surface water contamination. Do not use this product within 10 meters of the above mentioned water sources. Avoid contamination of food and feed, domestic or irrigation water supplies, lakes, streams and ponds. Do not apply this product through any type of irrigation system. Use only on labelled crops. Do not apply at rates greater than 1.9 kg/ha, except for weed control in lowbush blueberries.

STORE IN A DRY PLACE.

FIRST AID: If swallowed, immediately give large quantities of water and induce vomiting. Never give anything by mouth to an unconscious person. Obtain medical attention. If in eyes splash repeatedly with large amounts of clean water. Obtain medical attention. If on skin, remove contaminated clothing immediately. Wash skin thoroughly with soap and water. If irritation occurs, obtain medical attention.

DECONTAMINATION AND DISPOSAL:

1. Follow provincial instructions for any required additional

- cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.
 3. Dispose of the container in accordance with provincial requirements.
 4. For information on the disposal of unused, unwanted product and the cleanup of spills contact the regional office of Conservation and Protection, Environment Canada.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

DUAL and KIL-MOR are registered trademarks of CIBA-GEIGY CANADA LTD.

SUTAN+ is a registered trademark of ICI Americas Ltd.

BLADEX is a registered trademark of SHELL CANADA LTD.

CIBA-GEIGY CANADA LTD. is the registered user.

BANVEL is a registered trademark of SANDOZ CROP PROTECTION CORP.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))