

85.12.31

DITCHLING

SULFARIN PELLETS

RAT AND MOUSE KILLER BAIT  
CONTAINS WARFARIN AND SULFAQUINOXALINE

GUARANTEE:

WARFARIN.....0.025%  
SULFAQUINOXALINE...0.025%

COMMERCIAL

REG.NO. 10883  
PCP ACT

READ LABEL BEFORE USING

DITCHLING CORPORATION LIMITED  
PO BOX 395,  
DON MILLS, ONTARIO  
M3C 2S7

NOTICE TO USER: This control product is to be used only in accordance with directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that

condition.

PRECAUTIONS: Avoid contamination of food and feedstuffs. Wash thoroughly after handling. Keep out of reach of children, pets and domestic animals. This product contains as its active ingredient an anticoagulant chemical which if taken accidentally by humans, animals or pets, may reduce the clotting ability of the blood and serious hemorrhage may result. In case baits are accidentally eaten, induce vomiting with a finger at the back of the throat and repeat until vomit fluid is clear. CALL A PHYSICIAN.

PELLETS for use in Processing Plants(feed, food), Processing Plants(non-food), Food Service Establishments, Storage Areas(feed, food), Storage Areas(non-food), Dwellings, Buildings(outdoor surfaces), Outdoor Living Areas, Residential Areas, Garbage Dumps, Garbage Containers, Outdoor Areas, and Farm Buildings.

DIRECTIONS:

RODENT CONTROL

Rats: Place from 50g to 500g in areas frequented by the rodents. Protect bait placements from children, pets and livestock. Where a continuous source of infestation prevails, permanent bait stations should be used and kept continuously filled with fresh bait.

Mice: To control Mice follow the same general directions as for Rats but baits may be smaller and more placements should be made. Mice are more difficult to control than Rats and complete control may take a longer period of baiting.

IMPORTANT: Ensure that bait is always available to Rodents until lack of feeding is noted. This should be from five to fourteen days.

TOXICOLOGICAL INFORMATION:

When a human has been known to have accidentally ingested Warfarin, blood transfusions combined with intravenous injections

and oral dosages of Vitamin K, are indicated as in case of hemorrhage caused by overdoses of Dicumarol.

CONTAINER DISPOSAL:

Do not reuse empty container. Container should be disposed of by punching holes in it and burying with wastes or by burning(keep out of smoke).

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))