

1995

WHITE ROSE(R) GUARDIAN

Plant and garden pest control products

ANT, EARWIG AND GRUB KILLER DUST

Kills many soil-inhabiting and household insects.

GUARANTEE: Chlorpyrifos 0.5%

REG. NO. 17798.01 P.C.P. ACT

DOMESTIC

READ THE LABEL BEFORE USING

500 g

WHITE ROSE NURSERIES
Unionville, Ontario
L3R 2L5

DIRECTIONS FOR USE: Pests out-of-doors

TURF AND LAWNS: Make no more than 2 applications per season. White Grubs, European chafer larvae, Japanese beetle larvae: apply 400 g per 10 square metres and water into turf.

Ants: apply 200 grams per 10 square metres. Treat ant nests and trails. Chinch Bugs: apply at a rate of 200 g per 10 square metres at first sign of damage. Water thoroughly. Keep off lawn until dry.

ORNAMENTAL SHRUBS AND BUSHES: Make no more than 2 applications per season. Ants, crickets, earwigs, sowbugs, spiders, cutworms, white grubs, European chafer larvae and Japanese beetle larvae. Apply dust to turf or to soil around plants at a rate of 400 g per 10 square metres. For control of the following foliar pests, dust plants lightly; aphids, mealybugs, mites, scales, spittlebugs, thrips, and whiteflies. Apply to give uniform coverage of upper and lower leaf surfaces. Do not use on roses, azaleas, camellias, poinsettias, oxalis, coleus or variegated ivy.

GARDENS: For control of cutworms apply if damage is expected.

Use on broccoli, brussels sprouts, cabbage, chinese cabbage, cauliflower, carrots, celery, cucumber, green pepper, onion, potato, sweet corn, and tomato. Apply dust to soil around plant at a rate of 70g per 10 metres of row. Do not use on green bunching onions.

For control of potato flea beetle and tarnished plant bug on potato apply dust to plant at a rate of 35g per 10 metres of row. Repeat weekly as needed. Do not make more than nine applications during one season.

Days to wait between application and harvest.

Broccoli	32 Days	Celery	70 Days
Brussels Sprouts	32 Days	Cucumber	60 Days
Cabbage	32 Days	Green Pepper	40 Days
Carrot	60 Days	Onion	60 Days
Chinese Cabbage	32 Days	Potato	7 Days
Cauliflower	32 Days	Sweet Corn	70 Days
		Tomato	70 Days

INSECT PESTS OF HOUSEPLANT SOIL AND ON HOUSEPLANTS: If possible remove plants out-of-doors for treatment.

For Soil Insects: Apply dust lightly to soil around plant. Repeat as needed but no more than once every 2 months.

For Foliar Insects: Dust plant to give thorough coverage of upper and lower leaf surfaces. Do not use on roses, azaleas, camellias, poinsettias, geraniums, oxalis, coleus or variegated ivy. Repeat as needed but no more than once every 2 months.

PRECAUTIONS: Do not allow dust to contact food, food preparing surfaces, utensils or drinking water. Wash contaminated food surfaces and utensils with soap and water.

Do not use in poultry houses or other animal buildings.

May cause irritation to skin or eyes; avoid contact. Avoid inhaling dust. Wash thoroughly with soap and water after using.

KEEP OUT OF REACH OF CHILDREN

WARNING: This product is toxic to fish and crustaceans. Avoid contamination of ornamental fish ponds and fish bowls. Dispose of the container away from food, feed or water supplies.

FIRST AID: If swallowed induce vomiting and seek medical attention. Flush contaminated eyes and skin with plenty of water. For eyes get prompt medical attention.

TOXICOLOGICAL INFORMATION: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine by injection only is an antidote.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))