

27-APR-2000

((LABEL))

Freedom™ herbicide

dry flowable

For Use On Imazethapyr Tolerant Canola
(e.g. Canola varieties with the SMART(TM) trait)

AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

GUARANTEE: Thifensulfuron methyl 75%

NET CONTENTS: 320 grams

REGISTRATION NO. 25869

PEST CONTROL PRODUCTS ACT

~~DUPONT CANADA INC.~~
AGRICULTURAL PRODUCTS
BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

E.I. DU PONT CANADA COMOPANY

Notification Change

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.
Avoid breathing spray mist.
Avoid contact with skin, eyes and clothing.
Wear a long sleeved shirt, long pants and
chemical resistant gloves while handling this
product.
Do not contaminate any body of water.

FIRST AID

EYE CONTACT: Flush eyes immediately with plenty of water for 15
minutes. Contact a physician if irritation
persists.

SKIN CONTACT: Wash skin with plenty of soap and water. Remove
and wash contaminated clothing before reuse.

INGESTION: Administer water freely and induce vomiting.

Repeat until vomit fluid is clear. Contact a physician or poison control centre.

For medical emergencies and transportation emergencies such as spills, leaks or fires, call 1-613-348-3616 (24 hours).

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

STORAGE AND DISPOSAL

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

The DuPont Oval® is a registered trademark of E.I. duPont de Nemours and Company. DuPont Canada is a licensee.

FREEDOM™ is a trademark of DuPont Canada Inc.

SMART™ is a trademark of American Cyanamid Company.
A00375971

((DIRECTIONS FOR USE BROCHURE))

Freedom™

Dry Flowable

For Use On Imazethapyr Tolerant Canola
(e.g. Canola varieties with the SMART(TM) trait)

AGRICULTURAL

READ THE LABEL AND THIS BOOKLET BEFORE USING

GUARANTEE: Thifensulfuron methyl 75%

REGISTRATION NO. 25869

PEST CONTROL PRODUCTS ACT

E.I. DU PONT CANADA COMPANY

~~DUPONT CANADA INC.~~
AGRICULTURAL PRODUCTS
BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

Notification Change

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.
Avoid breathing spray mist.
Avoid contact with skin, eyes and clothing.
Wear a long sleeved shirt, long pants and
chemical resistant gloves while handling this
product.
Do not contaminate any body of water.

FIRST AID

EYE CONTACT: Flush eyes immediately with plenty of water for 15
minutes. Contact a physician if irritation
persists.

SKIN CONTACT: Wash skin with plenty of soap and water. Remove
and wash contaminated clothing before reuse.

INGESTION: Administer water freely and induce vomiting.
Repeat until vomit fluid is clear. Contact a
physician or poison control centre.

For medical emergencies and transportation emergencies such as

spills, leaks or fires, call 1-613-348-3616 (24 hours).

IMPORTANT

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water that may be used on other crops.

Imazethapyr tolerant canola (e.g. Canola varieties with the SMART™ trait) must not be harvested, grazed or fed to livestock within 74 days of application.

Carefully observe sprayer cleanup instructions, as spray tank residue may damage crops other than spring wheat, winter wheat, barley, oats, imazethapyr tolerant canola (e.g. canola varieties with the SMART(TM) trait).

DO NOT APPLY BY AIR.

GENERAL INFORMATION

FREEDOM™ Herbicide is recommended for selective postemergence control of broadleaf weeds in, imazethapyr tolerant canola (eg. canola varieties with the SMART™ trait), not underseeded to legumes or grasses. FREEDOM™ is in the form of a dry flowable granule to be mixed in water and applied as a uniform broadcast spray. It is noncorrosive, nonflammable, nonvolatile, and does not freeze.

FREEDOM™ must be applied with a recommended surfactant early postemergence to the main flush of actively growing weeds. Warm, moist growing conditions promote active weed growth and enhance the activity of FREEDOM™ by allowing maximum foliar uptake and contact activity. Weeds hardened off by environmental stress may not be adequately controlled or suppressed and regrowth may occur.

FREEDOM™ may only be applied using ground equipment. For best results, ensure thorough spray coverage of target weeds. Refer to "Directions For Use" section of the label for complete use details.

FREEDOM™ stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending on growing conditions and weed susceptibility. Favourable growing conditions following treatment promote the activity of FREEDOM™ while cold, dry conditions delay the activity. Degree of control and duration of effect depend on weed sensitivity, weed size, crop competition, growing conditions and spray coverage.

DIRECTIONS FOR USE

For use only on:

Imazethapyr Tolerant Canola (eg. Canola Varieties with the SMART (TM) trait).

FOR USE ONLY IN THE PRAIRIE PROVINCES AND THE PEACE RIVER REGIONS OF BRITISH COLUMBIA.

FREEDOM™ Herbicide will control the following broadleaf weeds:

WEEDS CONTROLLED

Chickweed*
Hemp-nettle
Lady's-thumb
Redroot pigweed
Corn spurry
Russian thistle
Green smartweed
Stinkweed
Lamb's-quarters
volunteer rapeseed***
Cow cockle
Wild buckwheat**
Wild mustard

WEEDS SUPPRESSED

Kochia

Suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area.

For best results apply to emerged young actively growing weeds before the canopy closes. Weeds should be less than 10 cm tall or across. Weeds that emerge after treatment will not be controlled.

Note: If rain occurs soon after application control may be reduced. Several hours of dry weather are needed to allow FREEDOM

to be absorbed by weed foliage.

Specific Weed Recommendations

- * Chickweed: Apply FREEDOM™ herbicide plus recommended surfactant when the chickweed is small (1 to 6 leaf) and actively growing but before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.
- ** Wild Buckwheat: Apply FREEDOM™ plus surfactant to actively growing wild buckwheat in the 1 - 3 leaf stage. Under dry conditions control may be reduced. Large plants may regrow after treatment.
- *** Volunteer Rapeseed: FREEDOM™ Will not control imazethapyr Tolerant Canola (eg. canola varieties with the SMART™ trait).

For the control of annual grasses as well as broadleaf weeds, apply a tank mixture of:

FREEDOM™ at 20 grams per hectare
plus
ASSURE® II at 0.38 or 0.5 litres per hectare
plus
CANPLUS® 411 at 0.5-1.0 % v/v
or
SURE-MIX™ at 0.5 % v/v

Apply FREEDOM™ plus ASSURE II® tank mixture when the imazethapyr tolerant canola (eg. canola varieties with the SMART™ trait) is in the 3 to 6 leaf stage of growth but prior to beginning of bolting. Temporary crop yellowing may be observed shortly after application of the FREEDOM™ plus ASSURE II® tank mixture, especially when applied to the growing under environmentally stressful conditions (see USE PRECAUTIONS section of the label).

Application of the FREEDOM™ plus ASSURE II® tank mixture prior to the 3 leaf stage of imazethapyr tolerant canola (eg. canola varieties with the SMART™ trait) could result in crop injury. Avoid application to imazethapyr tolerant canola (eg. canola varieties with the SMART™ trait) fields in which germination is uneven (some plants outside the recommended leaf stage for application), as crop injury may result.

NOTE: DO NOT apply tonon-imazethapyr tolerant canola as severe crop injury or death of the plant may occur.

CAUTION: Application of the FREEDOM™ plus ASSURE® II tank mixture on imazethapyr tolerant canola (eg. Canola varieties with the

SMART™ trait) may result in a delay in flowering.

This tankmix will control the above broadleaf weeds as well as the grass weeds listed below:

RATE	WEEDS CONTROLLED	WEED LEAF STAGE
0.38 or 0.5 L/ha	Barnyard grass	2 leaf to early tillering*
	Green foxtail	2 leaf to early tillering
	Volunteer barley	2 leaf to early tillering
	Volunteer corn	2-6 leaf
	Volunteer oats	2 leaf to early tillering**
	Volunteer wheat	2 leaf to early tillering
	Wild Oats	1-5 leaf without tillers (0.38 L/ha) 2 leaf to early tillering** (0.5 L/ha)
	WEEDS SUPPRESSED	
	Quackgrass	2-6 leaf*

* Use the higher rate of ASSURE® II (0.5 L/ha) for control of barnyard grass, as well as suppression of quackgrass.

** Best results on volunteer/wild oats if application is made before tillering begins.

Use the higher rate of CANPLUS® 411 or SURE-MIX™ when wild oats or quackgrass are the main target weeds, or under conditions not conducive to good growth.

NOTE: Consult the ASSURE II® label for additional instructions regarding grassy weed size and/or height at time of application as well as other use precautions, and restrictions.

MIXING INSTRUCTIONS

1. Select a spray volume that will ensure thorough coverage and uniform spray pattern. Use a minimum of 100 L of spray volume per hectare.
2. Fill the spray tank 1/4 to 1/3 full with fresh water.
3. Add the recommended amount of FREEDOM™ Herbicide to the spray tank while the agitator is running. Make sure all of the FREEDOM™ is in suspension. Continuous agitation is required to keep FREEDOM™ Herbicide in suspension.
4. After FREEDOM™ Herbicide is in suspension, other tank mix components may be added while continuing to the tank with water. The proper sequence for tank mix components is:
 - I. dry formulations
 - II. emulsifiable concentrates (i.e. ASSURE® II)
 - III. surfactants or adjuvants
(recommended adjuvants, for use with ASSURE® II, tank only CANPLUS® and SURE-MIX™).

Note: Refer to the appropriate tank mix section of this label for more detailed recommendations on the specific surfactant or adjuvant required. A surfactant or adjuvant recommended on this label must be added to improve wetting and contact activity. Anti-foaming agents may be required.

Ensure that the herbicide is completely mixed before proceeding to the next step.

5. For repeat loads, reduce the tank heel to 10% or less of the previous load. Fill tank with fresh water as described in step 2, and continue as directed. Emulsifiable concentrates may make dispersion of FREEDOM™ Herbicide more difficult. For that reason, tank heels of the previous tank Mix should be kept to 10% or less of the spray tank volume.
6. Do not allow the spray mixture to remain in the tank for more than 24 hours before spraying or the effectiveness may be reduced. If the spray mixture has been allowed to stand in the tank, use vigorous agitation to thoroughly disperse before resuming spraying.
7. Use the spray preparation of FREEDOM™ Herbicide within 24 hours of product degradation may occur. If spray solution is left standing without agitation, thoroughly agitate and re-suspend any settled FREEDOM™ Herbicide before resuming the spraying operation.

The volumetric measuring cylinder for FREEDOM™ is to be used only as a guide. For precise measurement, use scales calibrated in grams.

EQUIPMENT-SPRAY VOLUMES

Apply the spray mixture uniformly with properly calibrated ground equipment only. Ensure thorough coverage and a uniform spray pattern. Flat fan nozzles are recommended (minimum 100 litres spray volume per hectare). When using flood jet or "Raindrop" nozzles, apply 225 litres per hectare total spray volume to ensure thorough coverage. Use 50 mesh filter screens or larger (metal or nylon).

Continuous agitation is required to keep FREEDOM™ in suspension. Apply uniformly and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over application.

SPRAYER CLEANUP

To avoid subsequent injury to crops other than wheat, barley, oats, imazethapyr tolerant canola varieties with the SMART(TM) trait) immediately after and prior to spraying other crops thoroughly remove all traces of FREEDOM(TM) from mixing and spray equipment as follows:

- 1) Drain tank and flush tank, boom and hoses with clean water for a minimum of ten minutes. Visually inspect tank to assure removal of all visible residues of FREEDOM(TM). If necessary, repeat step 1. DO NOT CLEAN SPRAYER NEAR WELL OR WATER SOURCE OR NEAR TO DESIRABLE VEGETATION.
- 2) Fill tank with clean water while adding 1 litre household ammonia (containing a minimum 3% ammonia) per 100 litres of water. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation. Again flush the hoses, boom and nozzles with the cleaning solution and drain tank.
- 3) Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
- 4) Repeat Step 2.
- 5) Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the hoses and boom.

CAUTION: Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour which may cause eye, nose, throat, and lung irritation. Do not clean equipment in an enclosed area.

USE PRECAUTIONS

Do not apply to wheat, barley, oats, imazethapyr tolerant canola (eg. canola varieties with the SMART(TM) trait that are or have been stressed prior to application by severe weather conditions, frost, low fertility, drought, water saturated soil, disease or insect damage as crop injury may result. Do not apply to registered crops if the above stress conditions are expected within 3 days after application as crop injury may result.

Do not apply within 15 metres of a body of water or wetland area.

Do not apply to irrigated land where tail water will be used to irrigate crop land. Do not contaminate irrigation water.

Under certain environmentally stressful conditions (such as heavy rainfall, prolonged cool or hot weather, frost conditions, wide fluctuations in day/night temperatures), temporary lightening in crop color and occasionally a slight reduction in crop height may occur.

Do not apply to crops underseeded to legumes, or grasses as injury to forage may result.

Because crop varieties differ in their tolerance to herbicides, limit first use of FREEDOM(TM) to a small area of each variety prior to adoption as a field practice.

Do not exceed a total of 20 g per hectare per crop year.

Do not plant any crop other than wheat, barley for 30 days after application of FREEDOM™.

Some naturally occurring biotypes of certain weeds listed on this label may not be controlled by ACCENT®, ELIM®, ULTIM®, REFINE®, ALLY®, AMBER®, EXPRESS PACK®, MUSTER®, PURSUIT®, REFINE EXTRA®, CLASSIC®, PINNACLE®, RELIANCE STS®, FREEDOM(TM), TBM 75®, the CROSS Toss-N-Go component of CROSSFIRE®, the EXTRA® component of CHAMPION EXTRA®, the REFINE component of REFINE® wo, FREEDOM(TM), the DF® component of LASER DF®, or the PLUS® component of TRIUMPH PLUS® and CHAMPION PLUS®. These populations are able to survive a use rate several times higher than needed to control susceptible populations. To delay selection for resistant populations rotate the use of ACCENT®, ELIM®, ULTIM®, REFINE®, ALLY®, AMBER®, EXPRESS PACK®, MUSTER®, PURSUIT®, REFINE EXTRA®, CLASSIC®, PINNACLE®, RELIANCE® STS®, FREEDOM®, TBM75®, CROSSFIRE®, CHAMPION EXTRA®, REFINE wo, LASER DF®, TRIUMPH PLUS and CHAMPION PLUS® with other herbicides effective on the same weeds. Avoid use of ACCENT®, ELIM®, ULTIM®, REFINE®, ALLY®, AMBER®, EXPRESS PACK®, LASER DF®, MUSTER®, PURSUIT®, REFINE EXTRA®, CLASSIC®, PINNACLE®,

RELIANCE STS®, FREEDOM(TM), TBM75®, CROSSFIRE®, CHAMPION EXTRA®, REFINE wo, TRIUMPH PLUS and CHAMPION PLUS repeatedly in the same field. Contact your local DuPont representative or the DuPont hotline at 1-800-667-3925 for further information.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe any Canadian patent.

STORAGE AND DISPOSAL

Storage:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

Disposal:

- 1) Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
- 2) Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
- 3) Make the empty container unsuitable for further use.
- 4) Dispose of the container in accordance with provincial requirements.
- 5) For information on the disposal of unused, unwanted product, and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

The DuPont Oval®, ACCENT®, ULTIM®, REFINE®, REFINE EXTRA®, ALLY, EXPRESS PACK®, ASSURE®, MUSTER®, and SURE-MIX® are trademarks of E.I. du Pont de Nemours and Company. DuPont Canada Inc. is a

licensee.

FREEDOM™ and ELIM® are trademarks of DuPont Canada Inc.

AMBER® is a registered trademark of Novartis Crop Protection Canada Inc.

PURSUIT® and SMART™ are trademarks of American Cyanamid Company.

CANPLUS® 411 is a registered trademark of ICI Americas Inc.

CITOWETT® PLUS is a registered trademark of BASF AG used under license by BASF Canada Inc.

LASER® and TRIUMPH® Plus are registered trademarks of Hoechst Schering AgrEvo GmbH, Germany.

27-APR-2000

DUPONT®
DuPont Canada

SUPPLEMENTAL LABEL

FREEDOM™ HERBICIDE

DRY FLOWABLE

Tank Mixes of: Freedom™ + Lontrel™ 360 EC
Freedom™ + Lontrel™ 360 EC + Assure® II

For Use on Imazethapyr Tolerant Canola (e.g. Canola varieties
with the SMART trait)

FOR USE ONLY IN THE PRAIRIE PROVINCES AND THE PEACE RIVER REGIONS
OF BRITISH COLUMBIA

AGRICULTURAL

READ THE LABEL AND THIS BOOKLET BEFORE USING

GUARANTEE: Thifensulfuron methyl 75%

REGISTRATION NO. 25869 PEST CONTROL PRODUCTS ACT

DUPONT CANADA INC.

AGRICULTURAL PRODUCTS
BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN.

Avoid breathing spray mist.

Avoid contact with skin, eyes and clothing.

Wear a long sleeved shirt, long pants and chemical resistant gloves while handling this product.

Do not contaminate any body of water.

FIRST AID

EYE CONTACT: Flush eyes immediately with plenty of water for 15 minutes. Contact a physician or a poison control centre.

SKIN CONTACT: Wash skin with plenty of soap and water. Remove and wash contaminated clothing before reuse.

INGESTION: Administer water freely and induce vomiting. Repeat until vomit fluid is clear. Contact a physician or poison control centre.

For medical emergencies and transportation emergencies such as spills, leaks or fires, call 1-613-348-3616 (24 hours).

DIRECTIONS FOR USE

FREEDOM™ + LONTREL™ 360 EC

For control of annual broadleaf weeds plus top growth control of Canada Thistle, apply a tank mixture of the following:

FREEDOM at 20 grams per hectare

plus

LONTREL 360 EC at 0.42 litres per hectare

plus

Non-ionic Surfactant at 0.2 % v/v

or

SURE-MIX at 0.5 % v/v or 5 litres per 1000 litres spray solution

Apply the FREEDOM™ plus LONTREL™ 360 tank mixture when the Imazethapyr tolerant canola (e.g. canola varieties with the SMART® trait) is in the 3 to 6-leaf stage of growth but prior to beginning of bolting. Application of the FREEDOM™ plus LONTREL™

360 tank mixture prior to the 3-leaf stage of Imazethapyr tolerant canola (e.g. canola varieties with the SMART® trait) could result in crop injury.

Weeds Controlled or Suppressed:

This tank mix will control or suppress the broadleaf weeds listed above with FREEDOM™ used alone, PLUS provide top growth control of Canada Thistle. Consult the LONTREL™ 360 EC label for additional instructions regarding Canada Thistle size at the time of application, as well as other use precautions.

FREEDOM + LONTREL 360 EC + ASSURE II

For top growth control of Canada Thistle as well as control of annual grasses and broadleaf weeds, apply a tank mixture of the following:

FREEDOM™	at 20 grams per hectare
plus	
LONTREL™ 360 EC	at 0.42 litres per hectare
plus	
ASSURE®™ II	at 0.38 or 0.5 litres per hectare
plus	
SURE-MIX	at 0.5 % v/v or 5 litres per 1000 litres spray solution

Or

CANPLUS® 411	at 0.5 - 1.0 % v/v
--------------	--------------------

Apply the FREEDOM™ plus LONTREL™ 360 plus ASSURE II tank mixture when the Imazethapyr tolerant canola (e.g. canola varieties with the SMART® trait) is in the 3 to 6-leaf stage of growth but prior to beginning of bolting. Application of the FREEDOM™ plus LONTREL™ 360 plus ASSURE® II tank mixture prior to the 3-leaf stage of Imazethapyr tolerant canola (e.g. canola varieties with the SMART® trait) could result in crop injury.

Weeds Controlled or Suppressed:

This tank mix will control or suppress the annual grasses & broadleaf weeds listed with the FREEDOM™ + ASSURE® II tank mix, PLUS provide top growth control of Canada Thistle. Consult the ASSURE® II and LONTREL™ 360 EC labels for additional instructions regarding weed size &/or height at the time of application, as well as other use precautions.

MIXING INSTRUCTIONS

1. Select a spray volume that will ensure thorough coverage and uniform spray pattern. Use a minimum of 100 L of spray volume per hectare.
2. Fill the spray tank 1/4 to 1/3 full with fresh water.
3. Add the recommended amount of FREEDOM™ Herbicide to the spray tank while the agitator is running. Make sure all of the FREEDOM™ is in suspension. Continuous agitation is required to keep FREEDOM™ Herbicide in suspension.
4. After FREEDOM™ Herbicide is in suspension, other tankmix components may be added while continuing to fill the tank with water. The proper sequence for tank mix components is:
 - I. dry formulations
 - II emulsifiable concentrates (i.e. LONTREL™ 360 EC, or ASSURE® II)
 - III Surfactants or adjuvants (recommended adjuvants, for use with ASSURE® II, LONTREL™ 360 EC or ASSURE® II + LONTREL™ 360 EC tank mixes are CANPLUS® 411 and SURE-MIX®).

Note: Refer to the appropriate tank mix section of this label for more detailed recommendations on the specific surfactant or adjuvant required. A surfactant or adjuvant recommended on this label must be added to improve wetting and contact activity. Anti-foaming agents may be required. Ensure that the herbicide is completely mixed before proceeding to the next step.

5. For repeat loads, reduce the tank heel to 10% or less of the previous load. Fill tank with fresh water as described in step 2, and continue as directed. Emulsifiable concentrates may make dispersion of FREEDOM™ Herbicide more difficult. For that reason, tank heels of the previous tank mix should be kept to 10% or less of the spray tank volume.
6. Do not allow the spray mixture to remain in the tank for more than 24 hours before spraying or the effectiveness may be reduced. If the spray mixture has been allowed to stand in the tank, use vigorous agitation to thoroughly disperse before resuming spraying.
7. Use the spray preparation of FREEDOM™ Herbicide within 24 hours or product degradation may occur. If spray solution is left standing without agitation, thoroughly agitate and re-suspend any settled FREEDOM™ Herbicide before resuming the spraying operation.

The volumetric measuring cylinder for FREEDOM™ is to be used only

as a guide. For precise measurement, use scales calibrated in grams.

STORAGE AND DISPOSAL

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe any Canadian patent.

The DuPont Oval and ASSURE® are registered trademarks of E.I. du Pont de Nemours and Company.

DuPont® Canada Inc. is a licensee.

SURE-MIX™ is a trademark of E.I. du Pont de Nemours and Company. DuPont Canada Inc. is a licensee.

FREEDOM™ is a trademark of DuPont Canada Inc.

SMART® is a registered trademark of American Cyanamid Company.

LONTREL™ is a trademark of Dow AgroSciences LLC.

CANPLUS® 411 is a registered trademark of ICI Americas Inc.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))