

96.01.23

Allstar
DC Disinfectant Cleaner
DISINFECTANT-CLEANER-SANITIZER-FUNGICIDE-MILDEWSTAT-VIRUCIDE
DEODORIZER

With organic soil tolerance for Hospitals, Institutional and
Industrial use

COMMERCIAL

GUARANTEE:

Alkyl (C14, 50%;C12, 40%;C16,10%)dimethyl benzyl ammonium
chloride.....1.520%
Octyl decyl dimethyl ammonium chloride.....1.140%
Dioctyl dimethyl ammonium chloride.....0.456%
Didecyl dimethyl ammonium chloride.....0.684%

CAUTION

POISON

CORROSIVE

READ THE LABEL BEFORE USING

REGISTRATION NO. 24287 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 11 Litres

Manufactured by:
Unisource Canada Inc.
1475 Courtneypark Drive East
Mississauga, Ontario
L5T 2R1

For chemical emergencies only, call:
Infotrac 1-800-535-5053 (US)
Canutec 1-613-996-6666 (Canada)

Allstar DC Disinfectant Cleaner has been formulated to aid in the reduction of cross-contamination in institutions, schools and hospitals. Allstar DC Disinfectant Cleaner is a one-step disinfectant cleaner that is effective against a broad spectrum of bacteria, is virucidal, fungicidal and will prevent the growth of mold and mildew and their odors when used as directed.

Allstar DC Disinfectant Cleaner is an effective one-step sanitizer-cleaner for use on non-food contact surfaces. When used as directed, Allstar DC Disinfectant Cleaner will deodorize surfaces in toilet areas, behind and under sinks and counters, garbage cans and garbage storage areas and other places where bacterial growth can cause malodors.

PRECAUTIONARY STATEMENTS:

KEEP OUT OF REACH OF CHILDREN. DANGER-CORROSIVE TO EYES. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. HARMFUL IF SWALLOWED. AVOID CONTAMINATION OF FOOD.

FIRST-AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician or poison control centre immediately.

TOXICOLOGICAL INFORMATION: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to treatment site.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal or reconditioning.
3. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Environment Canada.
5. For small spills. Recover free liquid. Soak up residue with solids absorbent. DO NOT FLUSH INTO SURFACE STREAMS. INFORM ENVIRONMENT CANADA AS IN #4.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

DIRECTIONS FOR USE

Allstar DC Disinfectant Cleaner is a proven "one-step" cleaner-disinfectant-sanitizer-fungicide-mildewstat-virucide in the presence of moderate amounts of organic soil. Apply Allstar DC Disinfectant Cleaner to walls, floors and other hard (inanimate) non-porous surfaces such as tables, chairs,

countertops, sinks and bedframes with a cloth, or mop so as to thoroughly wet surfaces. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution daily or when use solution becomes visibly dirty.

DISINFECTATION - to disinfect hard, non-porous surfaces, add 17 mL/L of water. Treated surfaces must remain wet for 10 minutes. At this use level, Allstar DC Disinfectant Cleaner is also virucidal, fungicidal and mildewstatic. At 9 mL/L; Allstar DC Disinfectant Cleaner will disinfect hard non-porous surfaces in schools, industry and non-medical institutions.

SANITIZING - to sanitize porous or non-porous, non-food contact surfaces, add 5 mL/L of water. Treated surfaces must remain wet for 60 seconds.

For hospitals and nursing homes, add 17 mL/L of water. The broad spectrum activity of Allstar DC Disinfectant Cleaner has been evaluated in the presence of 5% serum and found to be effective against the following organisms by the AOAC Use-Dilution test: *Pseudomonas aeruginosa*, *Salmonella choleraesuis*, *Serratia marcescens*, *Enterobacter aerogenes*, *Streptococcus faecalis*, *Brevibacterium ammoniagenes*, *Staphylococcus aureus*, *Escherichia coli*, *Klebsiella pneumoniae*, *Salmonella schottmuelleri*, *Shigella dysenteriae*.

AOAC Fungicidal Test: At 9 mL/L Allstar DC Disinfectant Cleaner is an effective fungicide against *Trichophyton mentagrophytes* (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas, exercise facilities, etc.

Mold and Mildew Control: at 4 mL/L Allstar DC Disinfectant Cleaner will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, non-porous surfaces (as indicated in general instructions above). Allow to dry on surface and repeat when mildew growth returns.

Virucidal Performance: at 9 mL/L use level, Allstar DC Disinfectant Cleaner was evaluated in the presence of 5% serum and found to be effective against the Influenza A/Brazil Virus on inanimate non-porous environmental surfaces.

For schools, non-medical institutional and industrial uses: at 9 mL/L of water, Allstar DC Disinfectant Cleaner delivers excellent cleaning and germicidal effectiveness. It is effective against *Staphylococcus aureus*, *Salmonella choleraesuis* and *Escherichia coli*. The same AOAC tests used above to confirm suitable performance for hospitals were used.

SANITIZING - Non-Food Contact Surfaces (such as floors, walls, tables etc.): at 5 mL/L use-level, Allstar DC Disinfectant Cleaner is an effective sanitizer against *Staphylococcus aureus* and *Klebsiella pneumoniae* in the presence of 5% serum on porous and non-porous environmental surfaces.

Instructions for Cleaning and Decontaminating Against HIV-1 on Surfaces and Objects Soiled with Blood or Bodily Fluids:
This product is effective against HIV-1 (Human Immunodeficiency Virus) in the presence of organic soil (5% blood serum)

Personal Protection: When handling items soiled with blood or bodily fluids, wear disposable latex gloves, gowns, masks, and eye coverings.

Cleaning Procedure: Thoroughly clean blood or bodily fluids from surfaces before applying Allstar DC Disinfectant Cleaner.
Contact Time: To kill HIV-1 apply Allstar DC Disinfectant Cleaner at 9 mL per litre of water and allow surface to remain wet for 10 minutes.

Disposal of Infectious Material: Blood, body fluids, cleaning materials and clothing should be autoclaved and disposed of according to local regulations for infectious waste disposal.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))