

85.12.31

ENTEX (R)

OIL SOLUBLE CONCENTRATE

PCO INSECTICIDE

Contains fenthion

FOR USE BY QUALIFIED PEST CONTROL OPERATORS ONLY

COMMERCIAL

FOR CONTROL OF HOUSEHOLD PESTS

GUARANTEE: Fenthion.....480 g/L

NET CONTENT \_\_\_\_\_ LITRES

REGISTRATION NO. 8090  
PEST CONTROL PRODUCTS ACT

KEEP OUT OF REACH OF CHILDREN  
(See rear panel for first aid and danger statements)

READ THE LABEL BEFORE USING

CHEMAGRO LIMITED  
256 Britannia Road East  
Mississauga, Ontario, Canada L4Z 1S6

ENTEX Oil Soluble Concentrate

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

#### DIRECTIONS FOR USE

ENTEX Oil Soluble Concentrate is recommended for residual control of cockroaches, mosquitoes, flies, crickets, and wasps which are a problem in and around the home and other buildings such as restaurants, stores, warehouses, and industrial structures.

ENTEX may be used around the outside premises and in the inedible products departments of federally inspected meat and poultry processing plants. Before treatment, remove all exposed food products, packaging materials, and equipment and utensils used for handling edible products which may be in these inedible products areas.

For use inside buildings, application should be made as a coarse spray or with a paint brush to wallboards, cracks, crevices, and other hiding places where these insects have been seen running, resting, hiding, or nesting. Also apply to adjacent exposed surfaces where the insects will crawl when they come out of hiding. Repeat as necessary to maintain effective control.

For exterior use apply as a coarse spray or with a paint brush to selected surfaces such as doors, screens, window frames, door sills, and areas of porches where these insects alight or congregate. Apply to the nests of wasps after dark when all of the insects have returned to the nest. Repeat as necessary to maintain effective control.

Mix 250 to 375 mL of the oil soluble concentrate in 8 L of deodorized insecticide base oil for application to approximately 100 square meters of area or to run-off. Use the higher concentration for the initial treatment or for heavy cockroach infestations. For subsequent periodic applications and for control of flies and mosquitoes, use the lower dosage. The lower dosage may also be used in dairy, beef, and hog barns and poultry houses, but do not apply directly to animals.

#### MIXING CHART

Concentration of ENTEX Required	Amount of ENTEX OIL Soluble Concentrate Needed For:		
		1 L	5 L

1.5%	35 mL	100 mL	315 mL
2.25%	50 mL	235 mL	470 mL

#### RESTRICTIONS

Do not use as a space spray. Avoid excessive wetting of plastic, tile, rubber, and plastic materials. Do not spray on plants. Do not use in areas where children or pets can come into direct contact with treated surfaces. Remove fish or other pets from room before spraying. Do not contaminate feed, drinking water, milk or milk handling equipment.

#### DANGER

May be fatal if swallowed, inhaled, or absorbed rapidly through the skin. Do not get in eyes, on skin, or on clothing. Do not breathe spray mist. Wash thoroughly with soap and warm water after handling. Wash clothing with soap and hot water before reuse.

Avoid contamination of feed, food, or food-handling equipment. If food-handling equipment becomes contaminated, clean thoroughly before reuse. Keep out of reach of children.

FIRST AID: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, vomiting should be induced. Administer milk or water freely and induce vomiting by giving one dose (15 mL) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. The patient should be lying down with the head below the foot level and facing down or to one side. Professional medical assistance should be secured immediately. DO NOT INDUCE VOMITING TO AN UNCONSCIOUS PERSON OR TO PERSONS IN A CONVULSIVE STATE. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash immediately with flowing water for at least 15 minutes.

TOXICOLOGICAL INFORMATION: The compound inhibits cholinesterase resulting in stimulation of the central nervous system. This results in a sense of "tightness" in the chest, sweating, contracted pupils, stomach pains, vomiting and diarrhea. In case of poisoning call a physician immediately. The antidote is atropine sulfate administered in large therapeutic doses repeated as necessary to the point of tolerance, 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning

even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

NOTICE

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising, from the use or handling of this product and accepts the product on that condition.

ENTEX is a Reg. TM of Bayer AG.

Printed in U.S.A.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))