

91.11.12

RHONE-POULENC

NET WEIGHT/POIDS NET 2 KILOGRAMS

FIXED COPPER
50-W

CUIVRE FIXE
50-W

COPPER OXYCHLORIDE FUNGICIDE
FUNGICIDE A OXYCHLORURE DE CUIVRE
WETTABLE POWDER/POUDRE MOUILLABLE

AGRICULTURAL

GUARANTEE:

Copper 50%
REGISTRATION NO. 14741

PEST CONTROL PRODUCTS ACT
READ THE LABEL BEFORE USING

AGRICOLE

GARANTIE:

Cuivre 50%

NUMERO D'HOMOLOGATION 14741
LOI SUR LES PRODUITS ANTIPARASITAIRES

LIRE L'ETIQUETTE AVANT L'EMPLOI

RHONE-POULENC

RHONE-POULENC CANADA INC.
2000 ARGENTIA ROAD PLAZA 3
SUITE 400

MISSISSAUGA, ONTARIO L5N 1V9

TEL: (416) 821-4450 FAX: (416) 821-9339 TELEX 06-217701

In case of spills, poisoning or fire
telephone emergency response number collect
(416) 634-2359 (24 hours a day)

En cas de renversement, d'empoisonnement
ou incendie, appeler à frais virés le numéro
de secours suivant:
(416) 634-2359 (24 heures par jour)

DIRECTIONS FOR USE

GENERAL INFORMATION: Rates per 1000 litres are based on a dilute spray mixture. FIXED COPPER 50-W may be used in low volume or concentrate sprays by increasing the dosage to apply the same quantity of FIXED COPPER 50-W per hectare as would normally be applied when spraying dilute mixtures. Apply in sufficient water for thorough coverage of all plant parts; this volume depends on spray equipment and size of plants.

CROP	DISEASE	RATE AND METHOD
CUCUMBER, MELON, PUMPKIN, SQUASH	Angular Leaf Spot Anthracnose Bacterial Wilt Downy Mildew Fungus Leaf Spot	Use 4.5 kg in 1000 L of water per hectare. Spray at weekly intervals. For Bacterial Wilt Control it is necessary to control cucumber beetles. May be used up to each picking.

POTATO, TOMATO, CELERY	Early Blight Late Blight Septoria Leaf Spot Bacterial Canker (on greenhouse tomatoes)	Use 4.5 kg in 1000 L of water per hectare. Start early in the growing season with regular applications at 7-10 day intervals. May be used up to each day of harvest. For Bacterial Canker use at the rate of 3.25 kg in 1000 L/ha.
GRAPES	Downy Mildew Powdery Mildew	Mix 3 kg plus 6 kg of hydrated lime in 1000 L. Apply at Pre-bloom at 1500 L/ha and Post-bloom and in Cover sprays at 2000 L/ha until mid-August. Do not apply on Seibel varieties.
SOUR, CHERRY	Brown Rot Leaf Spot	Use 2 kg plus 4 kg hydrated lime per 1000 L of dilute spray. Consult local spray calendar for timing of application.
PEACH, APRICOT (British Columbia)	Coryneum Blight Peach Leaf Curl	Apply 9 kg/ha in September AFTER the harvest is completed.

COTON- EASTER, CRABAPPLE, HAWTHORN, MOUNTAIN ASH, QUINCE	Fireblight (see below)	Apply 1.25 kg/1000 L. Spray several times during blossoming; also repeat during summer when bacteria spread rapidly in warm, rainy weather.
ARBORATIVE, CEDAR, FIR, JUNIPER, PINE, SPRUCE	Leafblight Needlecast Tipblight Twigblight	Apply 4kg/1000 L of water when new growth starts in spring and repeat 3 to 4 times at 10 day intervals. Further sprays are advisable during prolonged rainy periods.

FIREBLIGHT: May be spread to healthy plants by insects such as aphids and leafhoppers. It is therefore, of utmost importance that regular applications with an effective insecticide such as Thioden, are made. Diseased twigs which appear to be "scorched" must be cut off 15 cm below the infected part and burned; disinfectant pruning shears after each cut in a solution of 1 part "Javex" to 4 parts water.

COMPATIBILITY: FIXED COPPER with lime can be mixed with most insecticides, Super-Spred Oil. FIXED COPPER with lime cannot be mixed with insecticide wettable powders. When adding oil or an insecticide, observe the cautions and intervals before harvest on edible crops printed on the insecticide label.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Hazardous if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes and clothing. After handling, wash thoroughly with soap and water before eating or smoking and change to clean clothing. Do not apply or allow to drift to areas occupied by unprotected persons and animals or to streams, lakes or ponds to protect wildlife. Avoid contamination of feed or food, including such crops on which residue is unsafe.

FIRST AID: If swallowed, induce vomiting by touching the back of the throat with a finger or blunt object, repeat until vomit fluid is clear. Bring patient to nearest hospital taking this labelled container with you. In case of skin contact, remove contaminated clothing and wash skin with soap and water. If in eyes, flush with plenty of water for 15 minutes and go to nearest hospital at once.

STORAGE: Store in a cool, dry place away from food or feed.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the emptied container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of the unused, unwanted product and the cleanup of spills contact the regional office of Conservation and Protection, Environment Canada.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this

service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))