

08-AUG-2001

((Label))

**DIVIDEND® 36FS**

FUNGICIDE

AGRICULTURAL

Seed Treatment for control of certain diseases of wheat

**GUARANTEE:**

difenoconazole.....3.15 %

**READ THE LABEL AND BOOKLET BEFORE USING  
KEEP OUT OF THE REACH OF CHILDREN**

REGISTRATION NUMBER: **25775**  
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **10L**

**Syngenta Crop Protection Canada, Inc.**  
140 Research Lane  
Research Park, University of Guelph  
Guelph, Ontario  
N1G 4Z3

Label

**NOTICE TO USER:**

This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

**NOTICE TO BUYER:**

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

**FIRST AID:**

**IN CASE OF POISONING,** contact a physician or a poison control centre immediately. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**If swallowed,** induce vomiting. Obtain prompt medical attention or contact a poison control centre.

**If in eyes,** splash repeatedly with large amounts of clean water for at least 15 minutes. Obtain medical attention or contact a poison control centre.

**If on skin,** wash thoroughly with soap and water. If irritation occurs, obtain medical attention or contact a poison control centre.

**If Inhaled,** move patient to fresh air.

**TOXICOLOGICAL INFORMATION:**

If ingested, induce emesis or lavage stomach. Treat symptomatically.

**PRECAUTIONS:**

KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed. Avoid contact with eyes or prolonged contact with skin. Avoid inhalation of vapours or spray mist. Wash hands and face after handling and before eating or smoking. Wear a suitable dust mask when transferring seed to a storage bin.

For On-Farm Use: When handling DIVIDEND 36FS, contaminated equipment or seed treated with 36FS, wear long pants, a long-sleeved shirt and chemical resistant gloves.

For Commercial Seed Treatment Plant Use: When handling DIVIDEND 36FS, contaminated equipment or seed treated with DIVIDEND 36FS, wear long-sleeved coveralls over normal work clothing and chemical resistant gloves. In addition, wear a suitable dust mask when bagging or sewing bags of treated seed or when transferring seed to a storage bin.

This product is toxic to fish and other aquatic invertebrates. Do not apply directly to water or to areas where surface water is present. Do not contaminate food, feed, domestic or irrigation water supplies, lakes, streams and ponds. If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion.

#### **DECONTAMINATION AND DISPOSAL:**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

#### **CONTAINER DISPOSAL:**

##### **For refillable containers:**

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

##### **For recyclable containers:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

**IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING  
CALL 1-800-267-6351**

® DIVIDEND is a registered trademark of a Syngenta company

((Booklet))

**DIVIDEND® 36FS**

FUNGICIDE

AGRICULTURAL

Seed Treatment for control of certain diseases of wheat

**GUARANTEE:**

difenoconazole.....3.15 %

**READ THE LABEL AND BOOKLET BEFORE USING  
KEEP OUT OF THE REACH OF CHILDREN**

REGISTRATION NUMBER: **25775**

PEST CONTROL PRODUCTS ACT

**Syngenta Crop Protection Canada, Inc.**

140 Research Lane  
Research Park, University of Guelph  
Guelph, Ontario  
N1G 4Z3

BOOKLET

**NOTICE TO USER:**

This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

**NOTICE TO BUYER:**

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

**FIRST AID:**

**IN CASE OF POISONING**, contact a physician or a poison control centre immediately. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**If swallowed**, induce vomiting. Obtain prompt medical attention or contact a poison control centre.

**If in eyes**, splash repeatedly with large amounts of clean water for at least 15 minutes. Obtain medical attention or contact a poison control centre.

**If on skin**, wash thoroughly with soap and water. If irritation occurs, obtain medical attention or contact a poison control centre.

**If Inhaled**, move patient to fresh air.

**TOXICOLOGICAL INFORMATION:**

If ingested, induce emesis or lavage stomach. Treat symptomatically.

**PRECAUTIONS:**

KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed. Avoid contact with eyes or prolonged contact with skin. Avoid inhalation of vapours or spray mist. Wash hands and face after handling and before eating or smoking. Wear a suitable dust mask when transferring seed to a storage bin.

For On-Farm Use: When handling DIVIDEND 36FS, contaminated equipment or seed treated with 36FS, wear long pants, a long-sleeved shirt and chemical resistant gloves.

For Commercial Seed Treatment Plant Use: When handling DIVIDEND 36FS, contaminated equipment or seed treated with DIVIDEND

36FS, wear long-sleeved coveralls over normal work clothing and chemical resistant gloves. In addition, wear a suitable dust mask when bagging or sewing bags of treated seed or when transferring seed to a storage bin.

This product is toxic to fish and other aquatic invertebrates. Do not apply directly to water or to areas where surface water is present. Do not contaminate food, feed, domestic or irrigation water supplies, lakes, streams and ponds. If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion.

**DECONTAMINATION AND DISPOSAL:**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**CONTAINER DISPOSAL:**

**For refillable containers:**

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**For recyclable containers:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING***

***CALL 1-800-267-6351***

**GENERAL INFORMATION:**

DIVIDEND 36FS is a ready-to-use systemic seed dressing which controls or suppresses certain seed-borne, soil-borne and early season foliar disease of spring and winter wheat. The formulation contains a colouring agent which will colour the treated seed.

**Note:** Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigour and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigour. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of the seed before committing the total seed lot to a selected seed treatment. Due to seed quality and seed storage conditions beyond the control of Syngenta Crop Protection Canada, Inc., no claims are made to guarantee the germination of carry-over seed.

**DIRECTIONS FOR USE:**

DIVIDEND 36FS is a ready-to-use formulation for use in commercial seed treatment plants, and for on-farm treatment using standard gravity flow or mist type seed treatment equipment which accurately meters and mixes a flowable seed treatment. DIVIDEND 36FS may also be used in a treat-on-the-go air seeder. The equipment must provide uniform coverage of DIVIDEND 36FS on the seed. Uneven seed coverage may not give the desired level of disease control.

This product does not require the addition of water for application. Consult the manufacturer of the application equipment planned to be used for suitability for this application and for instructions on operation and calibration of the equipment.

**WHEAT:**

Apply DIVIDEND 36FS to wheat seed at the rates given in the following tables for control of common bunt (seed-borne and soil-borne), dwarf bunt (including soil-borne), loose smut, seed-borne *Fusarium* and *Septoria*, seed rots (caused by *Penicillium* and *Aspergillus*) and early season *Septoria* leaf blotch; and for suppression of take-all and common root rot.

**WINTER WHEAT:**

Rate/100 kg of seed	Diseases Controlled	Early Season Foliar Diseases Controlled <sup>1</sup>	Diseases Suppressed <sup>2</sup>
650 mL	Common Bunt <sup>3</sup> , Dwarf Bunt, Loose Smut, Seed-Borne <i>Septoria</i> , Seed-Borne <i>Fusarium</i> , General Seed Rots <sup>4</sup>	<i>Septoria</i> Leaf Blotch	Common Root Rot ( <i>Cochliobolu</i> <i>s</i> spp.), Take-All
325 mL	Common Bunt <sup>3</sup> , Dwarf Bunt, Loose Smut, Seed-Borne <i>Fusarium</i> , General Seed Rots <sup>4</sup>		Common Root Rot ( <i>Cochliobolu</i> <i>s</i> spp.), Take-All
163 mL	Common Bunt		

1. DIVIDEND 36FS provides control of early season *Septoria* leaf blotch in winter wheat for the first 6 weeks after planting. For full season control, apply a foliar fungicide according to label directions.
2. Suppression means consistent control at a level which is not optimal but is still of commercial benefit.
3. DIVIDEND 36FS controls both seed-borne and soil-borne common bunt.
4. General seed rots controlled include those caused by saprophytic organisms such as *Penicillium* and *Aspergillus*.

**SPRING WHEAT:**

Rate/100 kg of Seed	Diseases Controlled	Diseases Suppressed <sup>1</sup>
650 mL	Common Bunt <sup>2</sup> , Loose Smut, Seed-Borne <i>Septoria</i> , Seed-Borne <i>Fusarium</i> ,	Common Root Rot ( <i>Cochliobolus</i> spp.), Take-All



	<b>General Seed Rots<sup>3</sup></b>	
<b>325 mL</b>	<b>Common Bunt<sup>2</sup>, Loose Smut, Seed-Borne <i>Fusarium</i>, General Seed Rots<sup>3</sup></b>	<b>Common Root Rot (<i>Cochliobolus</i> spp.), Take-All</b>
<b>163 mL</b>	<b>Common Bunt</b>	

1. Suppression means consistent control at a level which is not optimal but is still of commercial benefit.
2. DIVIDEND 36FS controls both seed-borne and soil-borne common bunt.
3. General seed rots controlled include those caused by saprophytic organisms such as *Penicillium* and *Aspergillus*.

**USE LIMITATIONS:**

1. Do not use treated seed for food, feed or oil purposes.
2. Do not graze, feed green forage or cut for hay within 55 days of planting.
3. Do not plant any crop other than wheat within 30 days to fields in which treated seeds were planted.

**NOTE:**

Seed treated with DIVIDEND 36FS must be labeled with the following statement: "This seed has been treated with difenoconazole fungicide. Do not use for food, feed or oil purposes. Store away from feeds and foodstuffs. When handling treated seed, use chemical resistant gloves. Do not graze, feed green forage or cut for hay within 55 days of planting. Do not plant any crop other than wheat within 30 days to fields in which treated seeds were planted."

® DIVIDEND is a registered trademark of a Syngenta company

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore

not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+) )