

93.08.16

REUSABLE/REUTILISABLE

PREBAITED/PRE-APPATE

TOMCAT(R)

MOUSE BAIT STATION

STATION D'APPATS POUR SOURIS

KILLS MICE

TUE LES SOURIS

REG.NO./NO.D'ENR. 22406 P.C.P. ACT/L.P.A.

DOMESTIC

DOMESTIQUE

GUARANTEE/GARANTIE:

DIPHACINONE.....0.005%

READ THE LABEL BEFORE USING

(SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS)

KEEP OUT OF REACH OF CHILDREN

LIRE L'ETIQUETTE AVANT USAGE

(VOIR A L'ARRIERE POUR PRECAUTIONS SUPPLEMENTAIRES)

TENIR HORS DE PORTEE DES ENFANTS

NET CONTENTS:

1 PREBAITED MOUSE BAIT STATION WITH TOMCAT BAIT CHUNX
(28g)

POIDS NET:

1 STATION PRE-APPATEE POUR SOURIS AVEC TOMCAT BAIT CHUNX
(28g)

MANUFACTURED BY/FABRIQUE PAR:

MOTOMCO LTD.
MADISON, WI 53708 U.S.A.

DIRECTIONS FOR USE

USE RESTRICTIONS:

Use TOMCAT Mouse Bait Station to control the house mouse (*Mus musculus*) in and around dwellings.

SELECTION OF TREATMENT AREAS:

Determine areas where mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, beside burrows, or in corners and concealed places.

APPLICATION DIRECTIONS:

HOUSE MICE: Place bait station at intervals of 2.5 to 3 meters along walls or in corners where mice live or travel. The bait station can be secured to the floor with a screw or double-sided tape. When bait is consumed, replace with one TOMCAT Bait Chunx.

TO REFILL: (See illustrations)

1. Fully insert 2-pronged key in holes on top of TOMCAT Mouse Bait Station.
2. Bend key away from station.
3. Gently lift up.
4. Place one TOMCAT Bait Chunx in the triangular area next to the hinge.
5. Close lid until it snaps securely. Repeat steps for easy rebaiting. To prevent reinfestation, eliminate food, water and harborage as much as possible. If reinfestation does occur, repeat treatment.

DISPOSAL: Wrap and dispose of empty container in household garbage.

PRECAUTIONARY STATEMENTS: Hazards to humans and domestic animals.

PRECAUTIONS: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. If swallowed, call a physician or Poison Control Centre at once.

TOXICOLOGICAL INFORMATION: Vitamin K1 in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times(P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

Net Contents: 28 gms(1 oz.)

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))