

93.08.17

VAPORETTE

INTEGRAL BUCKLE
FLEA COLLAR FOR CATS

Kills Fleas Up To 4 Months

DOMESTIC

READ THE LABEL BEFORE USING

GUARANTEE: Carbaryl.....8.5%

REGISTRATION NO. 13607 PEST CONTROL PRODUCTS ACT

SANDOZ AGRO CANADA INC.
Suite 302, Plaza 4, 2000 Argentia Road
Mississauga, Ontario L5N 1W1

Vaporette is a Registered trademark of Sandoz Ltd.

For Use in the control of cat and dog fleas on cats.

DIRECTIONS: Thread flat end of collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle. Collar must be worn loosely, but securely enough to prevent easy removal. Cut off any excess length and discard in trash. As the plastic buckle is moulded into the collar, the probability of neck irritation and buckle failure is reduced. To release the collar, squeeze buckle between thumb and finger and pull sharply.

The fleas on the cat will be killed and new ones, which may temporarily appear on the cat will also be killed during the 4 months (17 weeks) the collar is worn. FOR CONTINUOUS PROTECTION, REPLACE COLLAR WHEN EFFECTIVENESS DIMINISHES.

PRECAUTIONS: Do not open inner envelope until ready to use. DO

NOT ALLOW CHILDREN TO HANDLE UNATTACHED COLLAR. This collar is intended for use as an insecticide generator and is not be taken internally by man or animal. Not intended for use on humans. Hands should be washed following application.

Some animals may be sensitive to this collar. Any collar, when fastened too tightly, may cause skin irritation. If irritation persists after proper adjustment, remove the collar and consult a veterinarian.

Do not use on sick or convalescing animals. Do not use other pesticides on cat or administer other cholinesterase inhibitors, (e.g. oral parasiticides), while collar is worn or for a few days after removing collar. Do not use on kittens under 12 weeks of age or on nursing females. Consult your veterinarian before using this product on pregnant animals.

Dispose of any excess collar, packaging and expired collars with garbage to keep out of reach of children.

FIRST AID: In case of poisoning, call a physician immediately. If dust from collar contacts skin, wash thoroughly with soap and water. If dust contacts eyes, flush eyes with water and get medical attention.

TOXICOLOGICAL INFORMATION: Dust in eyes will cause temporary pupillary constriction. Dust release from collar is a cholinesterase inhibitor.

Atropine and Homatropine are antidotal.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))