

1998/2003

DowElanco

Dursban*

Home and Garden Insecticide

Granules

Contains Chlorpyrifos

DOMESTIC

GUARANTEE:

chlorpyrifos 0.5%

REGISTRATION NO. 15888

PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 500 g

DowElanco Canada Inc.
17705 Leslie Street
Newmarket, Ontario
L3Y 3E3
(905) 836-0436

* Trademark of DowElanco
DowElanco Canada Inc. is a licensed user

PRECAUTIONS

Do not store granules with food or feed or drinking water. Do not use in poultry houses or other animal buildings. May cause irritation to skin or eyes, avoid contact. Avoid inhaling dust from granules. Wash thoroughly with soap and water after using.

KEEP OUT OF REACH OF CHILDREN.

FIRST AID

If swallowed, induce vomiting and seek medical attention. Flush contaminated eyes and skin with plenty of water. For eyes get prompt medical attention.

TOXICOLOGICAL INFORMATION

Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine by injection only is an antidote.

WARNING

This product is toxic to fish and crustaceans. Avoid contamination of ornamental fish ponds and fish bowls.

DISPOSAL

Do not reuse container.
Dispose of empty container with household garbage.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

Do Not Ship or Store with Food, Feeds, Drugs or Clothing.

DIRECTIONS FOR USE

Turf and Lawns: Make no more than 2 applications per season. White grubs, European chafer larvae, Japanese beetle larvae. Apply 400 g per 10 m² and water into turf. Apply when grubs are small and feeding near soil surface, usually in late July or August.

Ants - Apply 200 g/10 m². Treat ant nests and trails. Chinch Bugs - Apply 200 g/10 m² at first sign of damage. Water thoroughly. Keep off lawn until dry. Sod Webworms - Apply 200 g/10 m² at first sign of damage. Water thoroughly. Keep off lawn until dry.

Ornamental Shrubs and Bushes: Ants, crickets, earwigs, sowbugs, spiders, cutworms, white grubs, European chafer larvae and Japanese beetle larvae. Apply to turf or to soil around plants at a rate of 400 g/10 m². Make no more than 2 applications per season.

Gardens: For control of cutworms and root maggots apply if damage is expected. Use on broccoli, Brussels sprouts, cabbage,

Chinese cabbage, cauliflower, carrots, celery, cucumber, green pepper, onion, potato, sweet corn and tomato. Apply 200 g/10 m2 of soil surface before planting. Do not make more than one application per year. Rake soil lightly to incorporate granules into top 5 cm of soil. Do not use on green bunching onions.

Days to wait between application and harvest:

Crop	Days	Crop	Days
Broccoli	32	Cucumber	60
Brussels sprouts	32	Green Pepper	40
Cabbage	32	Onion	60
Carrot	60	Potato	7
Chinese cabbage	32	Sweet corn	70
Cauliflower	32	Tomato	70
Celery	70		

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))