

86.05.30

Empire  
DISINFECTANT

VIREX

Cleaner, Detergent, Deodorizer, Sanitizer,  
Mildewstat (hard surface), Disinfectant, Fungicide

INDUSTRIAL

GUARANTEE

N-Alkyl (5% C12, 60% C14, 30% C16, 5% C18)	
Dimethyl Benzyl Ammonium Chlorides	0.8%
N-Alkyl (68% C12, 32% C14)	
Dimethyl Ethylbenzyl Ammonium Chlorides	0.8%

CAUTION  
CORROSIVE

READ THE LABEL BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Causes eye and skin irritation.  
Do not get in eyes or on skin.  
See side panel for additional precautions.

REGISTRATION NO. 19251  
PEST CONTROL PRODUCTS ACT

D.I.N. 664057

Net Contents                      Contenu Net

MADE IN CANADA

EMPIRE INTERNATIONAL  
Box 695, Streetsville P.S.  
Mississauga, Ontario L5M 2C2  
ATLANTA, GEORGIA; E. BRUNSWICK, NEW JERSEY;  
U.S.A.; SAN JUAN, PUERTO RICO; SINGAPORE

VIREX has been designed specifically for hospitals, offices, motels, nursing homes, schools, food processing plants and other institutions where housekeeping is of prime importance.

DISINFECTION - At 35.0 mL Virex/litre of water (5.6 oz. Virex/gallon of water) VIREX exhibits effective disinfectant activity against the organisms: Salmonella choleraesuis, Staphylococcus aureus and E. Coli. At this level VIREX is also fungicidal against the pathogenic fungi Trichophyton mentagraphytes.

To disinfect hard inanimate surfaces (such as walls, floors, table tops), apply solution with mop, cloth, sponge or mechanical sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes and then let air dry. DILUTION RATES: General Disinfection - Add 35.2 mL Virex/litre of water (5.6 oz. Virex/gallon). Hospital Disinfection - At 46.9 mL Virex/litre of water (7.5 oz. Virex/gallon of water) VIREX is an effective hospital disinfectant and demonstrates disinfectant activity against Pseudomonas aeruginosa.

To clean walls, floors, tables and other hard surfaces in operating rooms, patient quarters, washrooms etc., add 46.9 mL Virex/litre of water (7.5 oz. Virex/gallon of water). To disinfect toilet bowls, flush toilet, add 35.2 mL (1.2 oz) of Virex directly to the bowl water. Swab the bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand for 10 minutes and flush. For heavily soiled areas, a pre-cleaning step is required and recommended. Prepare a fresh solution for each use.

MILDEWSTAT - VIREX prevents and controls mold and mildew and the odours they cause on hard non-porous inanimate surfaces. Clean surfaces thoroughly, then add 35.2 mL Virex/litre of water (5.6 oz. Virex/gallon of water). Apply with a cloth, mop or sponge to surfaces to be treated making sure to wet thoroughly. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth reappears.

SANITIZATION - When used for sanitization of previously cleaned and rinsed non-porous food contact surfaces, prepare 200 ppm active quaternary VIREX solution by adding 12.5 mL Virex/litre of water (2 oz. Virex/gallon of water). To sanitize immobile items such as tanks, chopping blocks and counter tops, flood the area with a 200 ppm VIREX solution or apply with a cloth or sponge, making sure to wet all surfaces completely for at least 60 seconds. Follow cleaning with a complete rinse with potable water. Allow to air dry. Prepare a fresh solution for each use. To sanitize mobile items, such as drinking glasses and eating utensils, immerse in a 200 ppm VIREX solution for at least 60 seconds making sure to immerse completely. All cleaning must be followed by a potable water rinse. Allow to air dry. Prepare a fresh solution daily or more frequently as soil is apparent.

PRECAUTION

KEEP OUT OF REACH OF CHILDREN

Causes eye and skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food.

Not to be used as a glass cleaner. Do not allow to dry on surface - all cleaning must be followed immediately with potable water rinse.

FIRST AID - In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before re-use. If swallowed, drink milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician.

NOTICE TO USER - This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

FOR INDUSTRIAL USE ONLY  
NOT FOR RESALE

E405

R0286

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))