

Hartz® 2 in 1 Flea and Tick Collar With Deodorant for Cats
Bright White, Bright Blue, Bright Pink, Bright Purple, Bright Red

DOMESTIC

Kills Fleas Up To 5 Months
Kills Ticks

GUARANTEE:

Tetrachlorvinphos.....14.55%

REG. NO. 25623 P.C.P. ACT

OPEN CARTON ON SIDE SEAM AND READ LABEL INSIDE
BEFORE USING

KEEP OUT OF REACH OF CHILDREN
FOR USE ONLY ON CATS

NET CONTENTS: 15 g

Made in U.S.A. for;
HARTZ CANADA INC.,
St. Thomas, Ontario
N5P 3W7
1-800-275-1414

UNCONDITIONAL MONEY BACK GUARANTEE by HARTZ®

To activate, hold collar at each end and stretch.

Hartz® products are designed to compliment each other in an integrated program for flea & tick management. To assure proper treatment, we recommend use of Hartz® products for flea and tick management.

The Hartz® 2 in 1® Flea & Tick Collar for Cats will begin immediately to kill fleas and ticks including black legged ticks (Deer Ticks), which may carry the organism which causes Lyme Disease, and Rocky Mountain Wood Ticks, which may carry the organism which causes Rocky Mountain Spotted Fever. For continuous protection against fleas under light pest pressures replace the collar every five months, for ticks every four months. Under conditions where cats are exposed to severe flea or tick infestations it may be necessary to replace the collar more frequently. The flea collar may be worn with a regular collar. Wetting will not impair the collar's effectiveness or pet's protection. However, the collar should be removed if the cat is bathed with soap or detergent.

DIRECTIONS: Do NOT remove the collar from its package until ready to use. Remove the collar from the package, unroll and STRETCH to activate. This releases the active ingredient at time of use and assures full activity. Place the flea collar around the cat's neck allowing a spacing of 2 fingers between the collar and neck and buckle in place. Leave 5 to 8 cm on the collar for extra adjustment. Cut off any excess length, wrap and dispose of it in the garbage. Wash hands with soap and water after handling collar.

When collar is first worn, observe the pet's neck area every few days for irritation or adverse reaction. Do NOT use on cats under 12 weeks of age. Consult a veterinarian before using on sick, aged, pregnant, or nursing animals or animals receiving drug or other pesticide treatment. Do not use this product on cats at the same time or within 30 days before or after treatment with, or exposure to, cholinesterase-inhibiting drugs, pesticides, or chemicals. Any irritation usually indicates the collar has been affixed too tightly. If irritation occurs, loosen the collar. If irritation continues, remove the collar from the animal and consult your veterinarian. Replace collar no more than once every 4 months for continual pest control in situations where there is an ongoing risk of pest infestation. RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE

For effective flea and tick control, treatment of the pet should be combined with sanitation of any area used by the pet. Vacuum floors, carpets and furniture, (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problem persists, an insecticidal treatment of the premises may be required. A veterinarian should be consulted when flea infestation continues to be a problem on the pet.

PRECAUTIONS:

HUMANS: KEEP OUT OF REACH OF CHILDREN. DO NOT ALLOW CHILDREN TO HANDLE THIS COLLAR. Do not unroll collar until ready to use. This collar is intended for use only as an insecticide generator cat collar. Dust will form on the collar during storage. Dust may irritate eyes, nose and throat. Harmful if collar is sucked or chewed, or if dust is inhaled or absorbed through skin. Avoid contact with mouth, eyes, skin and clothing. Avoid breathing dust from collar. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Avoid product contact with food, food utensils or places where food is prepared or stored.

PETS: Dust may irritate eyes, nose and throat. Harmful if collar is chewed or swallowed. Avoid contact with the cat's mouth, face, eyes and genitals. Avoid letting the pet inhale the dust from the collar.

Consult a veterinarian before treating nursing animals, pregnant or sick animals, or animals under drug or other pesticide treatment. It is not advisable to use this product or similar pesticides on nursing animals or kittens less than 12 weeks old. Any irritation usually indicates the collar has been affixed too tightly. If irritation occurs, loosen the collar. If irritation continues, remove the collar from the animal and consult your veterinarian.

FIRST AID:

HUMANS: IF SUCKED OR CHEWED, obtain medical attention or call a poison control centre immediately. IF UNINTENTIONAL SKIN CONTACT OCCURS, wash affected areas with soap and water. IF DUST FROM COLLAR GETS IN EYES, flush eyes with plenty of water. If skin or eye irritation persists, obtain medical attention. IF DUST FROM COLLAR IS INHALED, seek medical attention.

PETS: Sensitivities may occur after using ANY pesticide for pets. If signs of sensitivity occur remove collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. If swallowed or chewed, consult a veterinarian.

Take container, label, or product name and P.C.P. Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION: This product contains a pesticide that is a cholinesterase inhibitor (anti-cholinesterase compound). Symptoms of human poisoning may include headache, weakness, sweating, blurred vision, nausea and diarrhea. Obtain medical attention or call a poison control centre at once. Atropine is antidotal. Acute symptoms of overdoses in dogs and cats include diarrhea, salivation, vomiting, muscular tremors and weakness. Contact a veterinarian immediately.

STORAGE: Store in original, unopened container at room temperature, away from food.

DISPOSAL: Dispose of any excess collar, packaging and unwanted (used) collars in household garbage.

Hartz Canada Inc., St. Thomas, Ontario N5P 3W7

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.