

96.08.29

NET WEIGHT 2 kg

WILBUR-ELLIS  
IDEAS TO GROW WITH (R)

METHOXYCHLOR 50-W  
INSECTICIDE - WETTABLE POWDER

AGRICULTURAL

GUARANTEE: Methoxychlor 50%

REGISTRATION NO. 14757  
PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

WILBUR-ELLIS COMPANY  
P.O. BOX 16458  
FRESNO, CA  
93755

DIRECTIONS FOR USE

METHOXYCHLOR 50-W may be used in low volume or concentrate sprays by increasing the dosage to apply the same quantity of Methoxychlor per hectare as normally would be applied when spraying dilute mixtures. Apply in sufficient water for thorough coverage of all plant parts; this volume depends on spray equipment and size of plants. Use the lower rate when vegetable plants are small and the higher rate when crop is full grown or insect population is heavy.

CROP/INSECT	METHOD	RATE
-------------	--------	------

APPLE, PEAR, PEACH, CHERRY, APRICOT, PLUM Plum Curculio, Cherry Fruitfly	Starting at Petal-fall, make 2 to 3 applications at 7-14 day intervals. Follow provincial recommendations for proper timing. Do not apply during Bloom. Do not apply later than the following intervals before harvest; apple, pear, plum, cherry - 7 days; peach, apricot - 21 days	6.75 - 13.25 kg/ha
STRAWBERRY, GOOSEBERRY, CURRANT Strawberry Weevil Currant Fruitfly	Apply in areas where insects are a problem; follow provincial recommendations for proper timing. Use 2000 L/ha. Do not apply during Bloom and not later than 14 days before harvest.	4.5 kg/ha
ASPARAGUS Asparagus Beetle	Apply when insects are noticed and repeat as necessary but not later than 3 days before each cutting.	1.75- 3.25 kg/ha
CUCUMBER, MELON, SQUASH, PUMPKIN, EGGPLANT, PEPPER, TOMATO Leafhoppers, Flea Beetles, Cucumber Beetle, Blister Beetle, Tomato Fruitworm, Fall Armyworm	Apply when insects are noticed and repeat as necessary but not later than 1 day before each picking. Do not feed refuse to livestock.	2.25-4.5 kg/ha
BEANS, PEAS Bean Weevil, Bean Leaf Beetle, Mexican Bean Beetle, Leafhoppers, Pea Weevil	Apply when insects are first noticed and repeat as necessary. Do not apply later than 7 days before harvest and do not feed bean and pea foliage or straw to livestock.	2.25- 4.5 kg/ha
ONION Thrips	Apply when insects are first noticed and repeat as necessary. Do not apply on young lima beans. Do not apply later than 7 days before harvest and do not feed bean and pea foliage or straw to livestock.	2.25-4.5 kg/ha

<p>BEETS, CAULIFLOWER, CABBAGE, CARROTS, BROCCOLI, LETTUCE, BRUSSELS SPROUTS, RADISH, SPINACH, TURNIP, POTATO flea beetle, leafhoppers, cabbage worm, blister beetle, Colorado potato beetle</p>	<p>Apply when insects are first noticed and repeat as necessary. Do not feed treated crop or crop refuse to livestock. Do not apply later than the following intervals before harvest; beet, carrot, cabbage, cauliflower, radish, turnip, potato - 7 days; broccoli, lettuce, spinach, Brussels sprouts - 14 days.</p>	<p>2.25 - 4.5 kg/ha</p>
<p>MUSHROOM HOUSES Phorid Flies Sciarid Flies</p>	<p>BETWEEN CROPS - When house is empty spray inside walls, floors, windows and woodwork to the point of run-off at the rate of 5 L / 100 m<sup>2</sup>. Repeat monthly until fall filling begins. PRE-SPAWNING - Repeat application during cool-down when air temperature drops to 32 Celsius. Do not spray ceilings or beds. PRE-CASING - Repeat 14 days after pre-spawning spray, no later than 4 days before casing. Do not spray ceilings or beds. PRE-PRODUCTION - Repeat again 7 days after casing. Do not spray ceilings or beds but apply on side boards with a brush.</p>	<p>10kg/ 1000 L</p> <p>5kg/ 1000 L</p>
<p>ALFALFA, CLOVER Alfalfa Weevil Sweet Clover Weevil Flea Beetle, Leafhoppers Grasshoppers, Armyworm</p>	<p>Apply when insects are first noticed and repeat as necessary. Do not spray when crop is in Bloom. Do not apply within 7 days of harvest or grazing. Follow your provincial recommendation as to the necessity and proper timing of the application.</p>	<p>2.25 kg/ha</p> <p>For Armyworm use 4.5 kg/ha</p>

ORNAMENTAL TREES, SHRUBS AND FLOWERS Thrips, Lace Bug, Leafhoppers, Chafer Weevils, Caterpillars, Elm and Rose Leaf Beetle, Webworms	Apply when insects are first noticed and repeat as necessary. Use a hydraulic sprayer with hose and gun. Avoid contamination of those fruits and vegetables for which the use is different from the directions on this label.	2 kg /1000 L
---	--	-----------------

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Do not inhale dust or spray mist. Avoid contact with skin and eyes. After handling, wash thoroughly with soap and water before eating or smoking. Do not contaminate ponds, streams or other bodies of water. Avoid contamination of feed or food.

FIRST AID: If swallowed, induce vomiting by sticking a finger or blunt object in the back of the throat; repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person. Take patient to nearest hospital taking this labelled container with you. In case of skin contact, remove contaminated clothing and wash skin with soap and water. If in eyes, flush with plenty of water for 15 minutes and get medical attention.

STORAGE AND DISPOSAL: Store in a cool, dry place away from food or feed. Thoroughly empty contents of bag into application equipment. Follow provincial instructions for any required additional cleaning of bag prior to its disposal. Make the empty bag unsuitable for further use. Dispose of the bag in accordance with provincial requirements. For information on the disposal of unused unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label

information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))